

=&gt; d his

(FILE 'HOME' ENTERED AT 11:24:56 ON 16 DEC 2004)

FILE 'HCAPLUS' ENTERED AT 11:25:27 ON 16 DEC 2004

E WO2000-CA811/AP, PRN

L1 1 WO2000-CA811/AP, PRN

L2 1 US6391313/PN

L3 1 L1-2

FILE 'REGISTRY' ENTERED AT 11:26:35 ON 16 DEC 2004

FILE 'HCAPLUS' ENTERED AT 11:26:37 ON 16 DEC 2004

L4 TRA L3 1- RN : 4 TERMS

FILE 'REGISTRY' ENTERED AT 11:26:37 ON 16 DEC 2004

L5 4 SEA L4

FILE 'WPIX' ENTERED AT 11:26:42 ON 16 DEC 2004

E WO2000-CA811/AP, PRN

L6 1 WO2000-CA811/AP, PRN

L7 1 US6391313/PN

L8 1 L6-7

=&gt; b hcap

FILE 'HCAPLUS' ENTERED AT 11:27:25 ON 16 DEC 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 Dec 2004 VOL 141 ISS 25

FILE LAST UPDATED: 15 Dec 2004 (20041215/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=&gt; d all l3

L3 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 2001:63846 HCAPLUS

DN 134:120915

ED Entered STN: 26 Jan 2001

TI Multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis

IN Loosmore, Sheena M.; Yang, Yan-Ping; Klein, Michel H.; Sasaki, Ken

PA Connaught Laboratories Limited, Can.

SO PCT Int. Appl., 58 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K039-00

CC 63-3 (Pharmaceuticals)

Section cross-reference(s): 15

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001005424	A2	20010125	WO 2000-CA811	20000711 <--
WO 2001005424	A3	20010802		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

Search done by Noble Jarrell

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,  
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,  
 CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

US 6391313 B1 20020521 US 1999-353617 19990715 <--  
 CA 2378862 AA 20010125 CA 2000-2378862 20000711 <--  
 EP 1200122 A2 20020502 EP 2000-945494 20000711 <--  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL

AU 767096 B2 20031030 AU 2000-59586 20000711 <--  
 PRAI US 1999-353617 A 19990715  
 WO 2000-CA811 W 20000711 <--

## CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 2001005424	ICM	A61K039-00
US 6391313	ECLA	A61K039/116

AB A multi-valent immunogenic composition confers protection on an immunized host against infection caused by both Haemophilus influenzae and Moraxella catarrhalis. Such composition comprises at least four antigens comprising at least one antigen from Haemophilus influenzae, and at least one antigen from Moraxella catarrhalis. Three of the antigens are adhesins. High mol. weight (HMW) proteins and Haemophilus influenzae adhesin (Hia) proteins of non-typeable Haemophilus and a 200 kDa outer membrane protein of Moraxella catarrhalis comprise the adhesin components while the other antigen is a non-proteolytic analog of Hin47 protein. Each component does not impair the immunogenicity of the others. The multi-valent immunogenic composition may be combined with DTP component vaccines, which may also include non-virulent poliovirus and PRP-T, to provide a component vaccine without impairment of the immunogenic properties of the other antigens.

ST adhesin antigen vaccine Haemophilus Moraxella

IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (HMW1; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (HMW2; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (Hin47; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (Hsf (Haemophilus surface fibril); multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (OMP (outer membrane protein); multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Immunostimulants  
 (adjuvants; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Agglutinins and Lectins  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
 (agglutinogens; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Adhesins  
RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
(antigenic; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Toxoids  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(diphtheria; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Organelle  
(fibril, surface; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Hemagglutinins  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(filamentous; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Chinchilla  
Haemophilus influenzae  
Molecular cloning  
Molecular weight distribution  
Moraxella catarrhalis  
Polyacrylamide gel electrophoresis  
Vaccines  
(multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Antigens  
RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
(multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Heat-shock proteins  
RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
(non-proteolytic; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Human poliovirus  
(non-virulent; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Ear  
(otitis, otitis media; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Agglutinins and Lectins  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(pertactins; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Toxoids  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(pertussis; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Mutation  
(substitution; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Toxoids  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(tetanus; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT 9001-92-7, Proteinase  
RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence)

(activity levels; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT 7784-30-7, Aluminum phosphate 21645-51-2, Aluminum hydroxide, biological studies  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses) (adjuvant; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT 151-21-3, Sds, analysis  
RL: ARU (Analytical role, unclassified); ANST (Analytical study) (multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

=> b reg

FILE 'REGISTRY' ENTERED AT 11:27:31 ON 16 DEC 2004  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2004 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 14 DEC 2004 HIGHEST RN 797749-23-6  
DICTIONARY FILE UPDATES: 14 DEC 2004 HIGHEST RN 797749-23-6

TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=> d ide l5 tot

L5 ANSWER 1 OF 4 REGISTRY COPYRIGHT 2004 ACS on STN  
RN 21645-51-2 REGISTRY  
CN Aluminum hydroxide (Al(OH)3) (9CI) (CA INDEX NAME)  
OTHER CA INDEX NAMES:  
CN Aluminum hydroxide (6CI, 8CI)  
OTHER NAMES:  
CN 42STE  
CN A 3011  
CN AC 400  
CN AC 400 (hydroxide)  
CN AC 450  
CN AC 714KC  
CN AE 107  
CN AF 260  
CN AKP-DA  
CN Alcan SF 4  
CN Alcoa 331  
CN Alcoa 710  
CN Alcoa A 325  
CN Alcoa AS 301  
CN Alcoa C 30BF  
CN Alcoa C 31  
CN Alcoa C 33  
CN Alcoa C 330  
CN Alcoa C 331  
CN Alcoa C 333  
CN Alcoa C 385  
CN Alcoa H 65  
CN Alcoa OC 1000  
CN Alhydrogel  
CN Alolt 50AF  
CN Alolt 59  
CN Alolt 60FLS  
CN Alolt 8  
CN Alolt 80

Search done by Noble Jarrell

CN Alolt 90  
 CN Alternagel  
 CN Alugel  
 CN Alugelibys  
 CN Alumigel  
 CN Alumina trihydrate  
 CN Aluminic acid (H<sub>3</sub>AlO<sub>3</sub>)  
 CN Aluminum oxide (Al<sub>2</sub>O<sub>3</sub>), trihydrate  
 CN Aluminum oxide trihydrate  
 CN Aluminum trihydroxide  
 CN Alusal  
 CN Amberol ST 140F  
 CN Amphogel  
 CN Amphojel  
 CN Antipollon HT  
 CN Apyral  
 CN Apyral 120  
 CN Apyral 120VAW  
 CN Apyral 15  
 CN Apyral 2

ADDITIONAL NAMES NOT AVAILABLE IN THIS FORMAT - Use FCN, FIDE, or ALL for  
 DISPLAY

DR 546141-62-2, 12252-70-9, 13783-16-9, 8012-63-3, 8064-00-4, 1302-29-0,  
 128083-27-2, 106152-09-4, 51330-22-4, 151393-94-1, 159704-77-5

MF Al H3 O3

CI COM

LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BIOBUSINESS, BIOSIS, BIOTECHNO,  
 CA, CABA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB, CEN, CHEMCATS,  
 CHEMLIST, CHEMSAFE, CIN, CSCHM, CSNB, DDFU, DETHERM\*, DIOGENES, DIPPR\*,  
 DRUGU, EMBASE, ENCOMPLIT, ENCOMPLIT2, ENCOMPPAT, ENCOMPPAT2, GMELIN\*,  
 HSDB\*, IFICDB, IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, MSDS-OHS, NIOSHTIC,  
 PDLCOM\*, PIRA, PROMT, PS, RTECS\*, TOXCENTER, TULSA, USAN, USPAT2,  
 USPATFULL, VETU, VTB

(\*File contains numerically searchable property data)

Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*

(\*\*Enter CHEMLIST File for up-to-date regulatory information)


DT.CA Caplus document type: Book; Conference; Dissertation; Journal; Patent;  
 Report

RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study);  
 FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU  
 (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT  
 (Reactant or reagent); USES (Uses); NORL (No role in record)

RLD.P Roles for non-specific derivatives from patents: ANST (Analytical  
 study); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation);  
 PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES  
 (Uses)

RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological  
 study); CMBI (Combinatorial study); FORM (Formation, nonpreparative);  
 MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC  
 (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses);  
 NORL (No role in record)

RLD.NP Roles for non-specific derivatives from non-patents: ANST (Analytical  
 study); BIOL (Biological study); FORM (Formation, nonpreparative); MSC  
 (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process);  
 PRP (Properties); RACT (Reactant or reagent); USES (Uses)



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

22458 REFERENCES IN FILE CA (1907 TO DATE)  
 374 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 22498 REFERENCES IN FILE CAPLUS (1907 TO DATE)  
 1 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

L5 ANSWER 2 OF 4 REGISTRY COPYRIGHT 2004 ACS on STN

RN 9001-92-7 REGISTRY

CN Proteinase (9CI) (CA INDEX NAME)

OTHER NAMES:

CN .alpha.-N-Benzoyl-DL-arginine-p-nitroanilide hydrolase

CN 537 Acidic protease

Search done by Noble Jarrell

CN Actinase  
 CN Alcalase 2.5L DX  
 CN Alkalase 2.4L FG  
 CN Alkalase 2.5L Type DX  
 CN Alkalase 2.5L type X  
 CN Alkaline protease-L FG  
 CN ALP 901  
 CN Alphamalt BK 5020  
 CN Alphamalt LQ 4020  
 CN AO protease  
 CN APL 901  
 CN Aquatinase E  
 CN Arginine esterase  
 CN AS 1.398  
 CN AS 10  
 CN Azocaseinase  
 CN BAPAase  
 CN BAPNAase  
 CN Benzoyl arginine arylamidase  
 CN Benzoyl-DL-arginine-p-nitroanilide hydrolase  
 CN Biopraxe 30L  
 CN Biopraxe SP 4FG  
 CN Bioprotease A  
 CN Bioprotease N 100P  
 CN Biopurase  
 CN Biosoft PW  
 CN Carbonyl hydrolase  
 CN Casein endopeptidase  
 CN Caseinase  
 CN CL-5PG  
 CN Cleanase AP 100-PWC  
 CN Corolase 7089  
 CN Corolase L 10  
 CN DA 10  
 CN DA 10 (enzyme)  
 CN Denapsin 10P  
 CN Denatyme AP  
 CN Deozyme  
 CN Deterzyme L-600  
 CN Durazyme 16.0L  
 CN Endopeptidase  
 CN Endopeptidase O  
 CN Endoprotease  
 CN Endoproteinase  
 CN Enzeco fungal acid protease  
 CN Enzylase K 40  
 CN Enzylon SAL  
 CN Enzylon SAL 300

ADDITIONAL NAMES NOT AVAILABLE IN THIS FORMAT - Use FCN, FIDE, or ALL for  
DISPLAY

DR 9001-93-8, 9012-23-1, 9040-76-0, 125498-72-8, 125752-86-5, 123779-18-0,  
 124041-97-0, 120038-39-3, 120038-40-6, 105913-13-1, 118901-82-9,  
 144906-30-9, 143404-30-2, 143404-41-5, 80804-52-0, 116267-38-0,  
 117278-03-2, 117698-27-8, 118390-80-0

MF Unspecified

CI COM, MAN

LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BIOBUSINESS, BIOSIS, BIOTECHNO,  
 CA, CAPLUS, CASREACT, CBNB, CEN, CHEMCATS, CHEMINFORMRX, CHEMLIST, CIN,  
 CSCHM, CSNB, DDFU, DIOGENES, DRUGU, EMBASE, IFICDB, IFIPAT, IFIUDB,  
 IPA, MSDS-OHS, NAPRALERT, NIOSHTIC, PIRA, PLASPEC\*, PROMT, RTECS\*,  
 TOXCENTER, TULSA, USPAT2, USPATFULL, VTB  
 (\*File contains numerically searchable property data)

Other Sources: EINECS\*\*, TSCA\*\*

(\*\*Enter CHEMLIST File for up-to-date regulatory information)

DT.CA Caplus document type: Book; Conference; Dissertation; Journal; Patent;  
Report

RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study);  
 CMBI (Combinatorial study); FORM (Formation, nonpreparative); MSC  
 (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process);  
 PRP (Properties); RACT (Reactant or reagent); USES (Uses); NORL (No role  
 in record)

RLD.P Roles for non-specific derivatives from patents: ANST (Analytical  
 study); BIOL (Biological study); MSC (Miscellaneous); OCCU (Occurrence);  
 PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or  
 reagent); USES (Uses)

RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological

study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses); NORL (No role in record)  
RLD.NP Roles for non-specific derivatives from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*


\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

40754 REFERENCES IN FILE CA (1907 TO DATE)  
491 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
40811 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L5 ANSWER 3 OF 4 REGISTRY COPYRIGHT 2004 ACS on STN  
RN 7784-30-7 REGISTRY  
CN Phosphoric acid, aluminum salt (1:1) (8CI, 9CI) (CA INDEX NAME)  
OTHER CA INDEX NAMES:  
CN Aluminum phosphate (Al(PO4)) (7CI)  
OTHER NAMES:  
CN ALPO  
CN AlPO 11  
CN AlPO 5  
CN aluminophosphate (AlPO4)  
CN Aluminum monophosphate  
CN Aluminum orthophosphate  
CN Aluminum phosphate  
CN Aluminum phosphate (1:1)  
CN Aluphos  
CN Fabutit 320  
CN Fabutit 748  
CN FB 67  
CN FFB 32  
CN Fosfalugel  
CN Fosfalumina  
CN K-Bond 90  
CN Monoaluminum phosphate  
CN Phosphaljel  
CN Phosphalugel  
CN Phosphalujel  
CN Phosphalutab  
CN Phosphaluvet  
CN Ulcoid  
CN VPI  
AR 98499-64-0  
DR 13765-93-0, 8022-59-1, 135151-77-8, 51668-55-4, 36201-72-6, 37324-42-8, 93237-81-1, 89686-54-4, 52350-11-5  
MF Al . H3 O4 P  
CI COM  
LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CABA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB, CHEMCATS, CHEMINFORMRX, CHEMLIST, CIN, CSCHM, DDFU, DETHERM\*, DIPPR\*, DRUGU, EMBASE, ENCOMPLIT, ENCOMPLIT2, ENCOMPPAT, ENCOMPPAT2, GMELIN\*, IFICDB, IFIPAT, IFIUDB, IMSCSEARCH, IPA, MEDLINE, MRCK\*, MSDS-OHS, NIOSHTIC, PDLCOM\*, PIRA, PROMT, RTECS\*, TOXCENTER, TULSA, ULIDAT, USAN, USPAT2, USPATFULL, VETU, VTB  
(\*File contains numerically searchable property data)  
Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*  
(\*Enter CHEMLIST File for up-to-date regulatory information)  
DT.CA Caplus document type: Book; Conference; Dissertation; Journal; Patent; Preprint; Report  
RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses); NORL (No role in record)  
RLD.P Roles for non-specific derivatives from patents: BIOL (Biological study); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)  
RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses); NORL (No role in record)  
RLD.NP Roles for non-specific derivatives from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); OCCU

Search done by Noble Jarrell

(Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT  
(Reactant or reagent); USES (Uses)  
CRN (7664-38-2)



● A1

6359 REFERENCES IN FILE CA (1907 TO DATE)  
108 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
6374 REFERENCES IN FILE CAPLUS (1907 TO DATE)  
4 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

L5 ANSWER 4 OF 4 REGISTRY COPYRIGHT 2004 ACS on STN  
RN 151-21-3 REGISTRY  
CN Sulfuric acid monododecyl ester sodium salt (8CI, 9CI) (CA INDEX NAME)

OTHER NAMES:

CN Adeka Hope LS 35  
CN Adeka Hope LS 90  
CN Akyposal NLS  
CN Akyposal SDS  
CN Alscoap LN 40A  
CN Alscoap LN 90  
CN Alscoap MP 90N  
CN Alscoap SP 40  
CN Aquarex Me  
CN Avirol 101  
CN Avirol SL 2010  
CN Berol 452  
CN Bio-Soft SDBS 60  
CN Calfoam SLS 30  
CN Carsonol SLS-S  
CN Conco Sulfate WAS  
CN Cycloryl 21LS  
CN Cycloryl 580  
CN Dehydag Sulfate GL  
CN Dermacide  
CN Dodecyl sodium sulfate  
CN Dodecyl sulfate sodium salt  
CN Dreft  
CN Duponol C  
CN Duponol ME  
CN Duponol QC  
CN Duponol WA  
CN Duponol WA Dry  
CN Duponol WAQ  
CN Duponol WAQE  
CN Duponol WAQM  
CN Emal 10  
CN Emal 10 Needle  
CN Emal 10 Powder  
CN Emal 2F  
CN Emal 2F Needle  
CN Emal 2F30  
CN Emal O  
CN Emal OS  
CN Empicol 0303  
CN Empicol 0303VA  
CN Empicol BSD 70  
CN Empicol LPZ  
CN Empicol LS 30  
CN Empicol LX 28  
CN Empicol LX 28R  
CN Empicol LX 42  
CN Empicol LXSV 938U  
CN Empicol LXV  
CN Empicol LY 28S

ADDITIONAL NAMES NOT AVAILABLE IN THIS FORMAT - Use FCN, FIDE, or ALL for

Search done by Noble Jarrell



DISPLAY  
 DR 12738-53-3, 12765-21-8, 8012-56-4, 1334-67-4, 1335-72-4, 172826-72-1,  
 121481-64-9, 58640-35-0, 57176-54-2, 64441-33-4, 129203-37-8, 51222-39-0,  
 61711-39-5, 111726-87-5, 74433-77-5, 145269-44-9, 152155-52-7,  
 156108-01-9, 191490-40-1, 237743-45-2, 303179-49-9  
 MF C12 H26 O4 S . Na  
 CI COM  
 LC STN Files: ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, AQUIRE, BEILSTEIN\*,  
 BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CANCERLIT, CAOLD, CAPLUS, CASREACT,  
 CBNB, CEN, CHEMCATS, CHEMINFORMRX, CHEMLIST, CIN, CSCHM, CSNB, DDFU,  
 DETHERM\*, DIOGENES, DRUGU, EMBASE, ENCOMPLIT, ENCOMPLIT2, ENCOMPPAT,  
 ENCOMPPAT2, GMELIN\*, HSDB\*, IFICDB, IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*,  
 MSDS-OHS, NIOSHTIC, PDLCOM\*, PHAR, PIRA, PROMT, RTECS\*, TOXCENTER,  
 TULSA, ULIDAT, USAN, USPAT2, USPATFULL, VETU, VTB  
 (\*File contains numerically searchable property data)  
 Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*  
 (\*\*Enter CHEMLIST File for up-to-date regulatory information)  
 DT.CA Caplus document type: Book; Conference; Dissertation; Journal; Patent;  
 Preprint; Report  
 RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study);  
 MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC  
 (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses);  
 NORL (No role in record)  
 RLD.P Roles for non-specific derivatives from patents: ANST (Analytical  
 study); BIOL (Biological study); PREP (Preparation); PROC (Process); PRP  
 (Properties); USES (Uses)  
 RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological  
 study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU  
 (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT  
 (Reactant or reagent); USES (Uses); NORL (No role in record)  
 RLD.NP Roles for non-specific derivatives from non-patents: ANST (Analytical  
 study); BIOL (Biological study); FORM (Formation, nonpreparative); OCCU  
 (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT  
 (Reactant or reagent); USES (Uses)  
 CRN (151-41-7)

HO<sub>3</sub>SO- (CH<sub>2</sub>)<sub>11</sub>-Me

● Na

\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

30993 REFERENCES IN FILE CA (1907 TO DATE)  
 324 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 31054 REFERENCES IN FILE CAPLUS (1907 TO DATE)  
 32 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

=> b wpiX  
 FILE 'WPIX' ENTERED AT 11:27:38 ON 16 DEC 2004  
 COPYRIGHT (C) 2004 THE THOMSON CORPORATION

FILE LAST UPDATED: 13 DEC 2004 <20041213/UP>  
 MOST RECENT DERWENT UPDATE: 200480 <200480/DW>  
 DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,  
 PLEASE VISIT:  
[http://www.stn-international.de/training\\_center/patents/stn\\_guide.pdf](http://www.stn-international.de/training_center/patents/stn_guide.pdf) <<<

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE  
<http://thomsonderwent.com/coverage/latestupdates/> <<<

>>> FOR INFORMATION ON ALL DERWENT WORLD PATENTS INDEX USER  
 GUIDES, PLEASE VISIT:  
<http://thomsonderwent.com/support/userguides/> <<<

>>> NEW! FAST-ALERTING ACCESS TO NEWLY-PUBLISHED PATENT  
 DOCUMENTATION NOW AVAILABLE IN DERWENT WORLD PATENTS INDEX  
 FIRST VIEW - FILE WPIFV.  
 FOR FURTHER DETAILS: <http://www.thomsonderwent.com/dwpifv> <<<

Search done by Noble Jarrell

>>> NEW DISPLAY FORMAT HITSTR ADDED ALLOWING DISPLAY OF  
HIT STRUCTURES WITHIN THE BIBLIOGRAPHIC DOCUMENT <<<

>>> SMILES and ISOSMILES strings are no longer available as  
Derwent Chemistry Resource display fields <<<

=> d all 18 tot

L8 ANSWER 1 OF 1 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN  
AN 2001-168447 [17] WPIX  
DNC C2001-050284  
TI Novel multivalent immunogenic composition for conferring protection  
against infection caused by *Hameophilus influenzae* and *Moraxella*  
*catarrhalis* comprises four antigens derived from each of the two  
microorganisms.  
DC B04 D16  
IN KLEIN, M H; LOOSMORE, S M; SASAKI, K; YANG, Y  
PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD  
CYC 95  
PI WO 2001005424 A2 20010125 (200117)\* EN 58 A61K039-00  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TZ UG ZW  
W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM  
DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC  
LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE  
SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW  
AU 2000059586 A 20010205 (200128) A61K039-00  
EP 1200122 A2 20020502 (200236) EN A61K039-116  
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI  
US 6391313 B1 20020521 (200239) A61K039-116 <--  
AU 767096 B 20031030 (200382) A61K039-00  
NZ 516819 A 20031219 (200404) A61K039-00  
ADT WO 2001005424 A2 WO 2000-CA811 20000711; AU 2000059586 A AU  
2000-59586 20000711; EP 1200122 A2 EP 2000-945494 20000711, WO  
2000-CA811 20000711; US 6391313 B1 US 1999-353617 19990715; AU 767096  
B AU 2000-59586 20000711; NZ 516819 A NZ 2000-516819 20000711, WO  
2000-CA811 20000711  
FDT AU 2000059586 A Based on WO 2001005424; EP 1200122 A2 Based on WO  
2001005424; AU 767096 B Previous Publ. AU 2000059586, Based on WO  
2001005424; NZ 516819 A Based on WO 2001005424  
PRAI US 1999-353617 19990715  
IC ICM A61K039-00; A61K039-116  
ICS A61P031-04  
AB WO 200105424 A UPAB: 20010328  
NOVELTY - A multivalent immunogenic composition (I) for conferring  
protection in a host against disease caused by both *Hameophilus influenzae*  
(HI) and *Moraxella catarrhalis* (MC) comprising four different antigens, of  
which at least one antigen is from HI and one antigen is from MC, is new.  
Additionally three of the antigens of (I) are adhesins, and one is from  
MC.  
ACTIVITY - Auditory; antibacterial.  
MECHANISM OF ACTION - Vaccine.  
Groups of five BALB/C mice were immunized subcutaneously on days 1, 29  
and 43 with one of the mouse H91A Hin47 + rHMW + rHia + r200 kDa vaccines.  
Blood samples were taken on days 0, 14, 28, 42 and 56. Groups of five  
Hartley outbred guinea pigs were immunized intramuscularly on days 1, 29  
and 43 with the vaccine as described above. Blood samples were taken on  
days 0, 14, 28, 42 and 56. Anti-H91A Hin47, anti-rHMW, anti-rHia and  
anti-r200 kDa IgG antibody titers were determined by antigen specific  
enzyme linked immunosorbant assays (ELISAs). The results of the  
immunogenicity studies showed that the final bleed sera obtained from mice  
immunized with 0.3 mu g, or 3.0 mu g each of H91A Hin47 + rHMW + rHia with  
0, 0.3, 1.0, 3.0 or 10.0 mu g of added r200 kDa, all had high antibody  
titers to H91A Hin47 component. The final bleed sera obtained from the  
mice immunized with 3.0 mu g each of H91A Hin47 + rHMW + rHia with 0, 0.3,  
1.0, 3.0 or 10.0 mu g of added r200 kDa, all had high titer antibodies to  
the rHMW apparent enhancing or inhibiting effect on the anti-rHMW response  
with the addition of the r200 kDa component. Mice immunized with 0.3 mu g  
each of H91A Hin 47 + HMW + rHia with 0, 0.3, 1.0, 3.0 or 10.0 mu g of  
added r200 kDa, all had high titer antibodies to the rHia component. There  
was no apparent enhancing or inhibiting effect on the anti-rHia response  
with the addition of the r200 kDa component. The final bleed sera obtained  
from guinea pigs immunized with 25 mu g or 50 mu g each of H91A Hin47 +  
rHMW + rHia with 0, 25, 50 or 100 mu g of added r200 kDa, all had high

Search done by Noble Jarrell

*Applicant*

titer antibodies to the H91A Hin47 component. Also final bleed sera obtained from guinea pigs immunized with 25 mu g or 50 mu g each of H91A Hin47 + rHMW + rHia with 0, 25, 50 or 100 mu g of added r200 kDa, all had titer antibodies to the rHMW component. There was no apparent enhancing or inhibiting effect on the anti-rHMW response upon the addition of the r200 kDa antigen.

USE - (I) is useful for immunizing a host against infection caused by both HI and MC including otitis media (claimed).

ADVANTAGE - The multivalent vaccine can confer protection against encapsulated and unencapsulated HI and MC diseased in a safe and efficient manner.

Dwg.0/14

FS CPI

FA AB; DCN

MC CPI: B04-B04C1; B14-A01; B14-A01A; B14-N02; B14-S11B; D05-C02; D05-H07;  
D05-H12F

=> b home

FILE 'HOME' ENTERED AT 11:27:45 ON 16 DEC 2004

=>

=&gt; d his

(FILE 'HOME' ENTERED AT 11:24:56 ON 16 DEC 2004)

FILE 'HCAPLUS' ENTERED AT 11:25:27 ON 16 DEC 2004

E WO2000-CA811/AP,PRN  
 L1 1 WO2000-CA811/AP,PRN  
 L2 1 US6391313/PN  
 L3 1 L1-2

FILE 'REGISTRY' ENTERED AT 11:26:35 ON 16 DEC 2004

FILE 'HCAPLUS' ENTERED AT 11:26:37 ON 16 DEC 2004  
 L4 TRA L3 1- RN : 4 TERMS

FILE 'REGISTRY' ENTERED AT 11:26:37 ON 16 DEC 2004  
 L5 4 SEA L4

FILE 'WPIX' ENTERED AT 11:26:42 ON 16 DEC 2004

E WO2000-CA811/AP,PRN  
 L6 1 WO2000-CA811/AP,PRN  
 L7 1 US6391313/PN  
 L8 1 L6-7

FILE 'HCAPLUS' ENTERED AT 11:46:19 ON 16 DEC 2004

E IMMUNOSTIMULANTS/CT  
 E E3+ALL  
 L9 16542 IMMUNOSTIMULANTS+NT/CT  
 E IMMUNE ADJUVANTS/CT  
 E E3+ALL  
 L10 9606 IMMUNE ADJUVANTS/CT OR L9 (L) ADJUV?  
 E VACCINES/CT  
 E E3+ALL  
 E CONTRACEPTIVES/CT  
 E E3+ALL  
 L11 564 CONTRACEPTIVES/CT (L) ?IMMUN?/BI  
 L12 40848 VACCINES/CT  
 E IMMUNITY/CT  
 E E3+ALL  
 L13 63934 IMMUNITY+NT/CT  
 E IMMUNOTHERAPY/CT  
 E E3+ALL  
 L14 12699 IMMUNOTHERAPY+NT/CT  
 E THERAPEUTICS/CT  
 E E3+ALL  
 E E2  
 E E3+ALL  
 L15 29488 THERAPY+OLD,NT/CT (L) ?IMMUN?/BI  
 E RADIOTHERAPY/CT  
 E E3+ALL  
 E HAEMOPHILUS INFLUENZAE/CT  
 E E3+ALL  
 E E4+ALL  
 L16 7294 HAEMOPHILUS+OLD,NT/CT  
 E MORAXELLA CATARRHALIS/CT  
 E E3+ALL  
 E E4  
 E E3+ALL  
 L17 1861 MORAXELLA+OLD,NT/CT  
 E ANTIGENS/CT  
 E E3+ALL  
 L18 QUE ANTIGENS+NT/CT  
 L19 80 L16 AND L17 AND L18  
 E ADHESIN/CT  
 E E5+ALL  
 E AGGLUTININS AND LECTINS/CT  
 E E3+ALL  
 L20 2116 "AGGLUTININS AND LECTINS"+OLD,NT/CT (L) ?ADHESIN?/BI  
 L21 2 L19 AND L20  
 E LOOSMORE S/AU  
 L22 83 E3-7  
 E YANG Y/AU  
 L23 786 E3,E26  
 E YANG YAN/AU  
 L24 321 E3,E31  
 E YANG, YANPING/AU

Search done by Noble Jarrell

*Invento  
Search*

E KLEIN ,/AU  
 E KLEIN M/AU  
 L25 513 E3,E12  
 E KLEIN MICHAEL/AU  
 L26 140 E3,E12  
 E SASAKI K/AU  
 L27 702 E3-6  
 E SASAKI KEN/AU  
 L28 889 E3-10  
 L29 182 (AVENTIS (1A) PASTEUR)/CS,PA  
 L30 3 L19 AND L22-29  
 L31 77 L19 NOT L30  
 L32 57 L31 AND L9-15  
 L33 QUE PY<=1999 OR AY<=1999 OR PRY<=1999 OR PD<19990715 OR PRD<199  
 L34 26 L32 AND L33  
 SEL AN 4 15 23 24 L34  
 L35 22 L34 NOT E1-8  
 L36 1 L20 AND L35  
 L37 21 L35 NOT L36  
 E OTITIS MEDIA/CT  
 E E3+ALL  
 E EAR, DISEASE/CT  
 E E3=ALL  
 E EAR, DISEASE/CT  
 E E3+ALL  
 L38 2966 "EAR, DISEASE"+NT/CT  
 E EAR/CT  
 E E3=ALL  
 E EAR/CT  
 E E3+ALL  
 L39 1839 EAR+OLD,NT/CT (L) DISEAS?  
 L40 9750 EAR+OLD,NT/CT  
 L41 818 L38-40 (L) OTITIS(1A) MEDIA  
 L42 3 L35 AND L41  
 L43 4 L36 OR L42  
 L44 18 L35 NOT L43

FILE 'WPIX' ENTERED AT 12:43:52 ON 16 DEC 2004

L45 65479 A61K039/IPC OR (B04-B04C? OR C04-B04C? OR B04-G01 OR C04-G01)/M  
 L46 28924 (B12-A01 OR C12-A01 OR B14-A01A OR C14-A01A)/MC OR MORAXELLA/BI  
 L47 28459 (B02-V02 OR C02-V02 OR B14-S11? OR C14-S11? OR D05-H07)/MC OR (  
 L48 989 L45 AND L46 AND L47  
 L49 205 (AVENTIS (1A) PASTEUR)/CS,PA  
 E LOOSMORE S/AU  
 L50 39 E3-4  
 E YANG Y/AU  
 L51 1938 E3,E19  
 E KELIN M/AU  
 E KLEIN M/AU  
 L52 333 E3,E13  
 E SASAKI K/AU  
 L53 1213 E3-7  
 L54 29 L48 AND L49-53  
 L55 960 L48 NOT L54  
 L56 4 L55 AND ADHESIN?/BIX  
 L57 8186 (B14-N02 OR C14-N02 OR B12-L04 OR C12-L04)/MC  
 L58 40 L55 AND L57  
 L59 0 L58 AND ADHESIN?/BIX  
 L60 18 L58 NOT (PY>1999 OR PRY>1999 OR AY>1999)  
 SEL AN 1-3 10 12  
 L61 5 E1-5 AND L60

=> b hcap

FILE 'HCAPLUS' ENTERED AT 13:09:10 ON 16 DEC 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing

Search done by Noble Jarrell

of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 Dec 2004 VOL 141 ISS 25  
FILE LAST UPDATED: 15 Dec 2004 (20041215/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d all 144 tot

L44 ANSWER 1 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
AN 2002:505235 HCAPLUS  
DN 137:62165  
ED Entered STN: 05 Jul 2002  
TI Producing antibodies with attenuated bacteria with altered DNA adenine methylase activity  
IN Mahan, Michael J.; Heithoff, Douglas M.; Low, David A.; Sinsheimer, Robert L.  
PA USA  
SO U.S. Pat. Appl. Publ., 44 pp., Cont.-in-part of U. S. Ser. No. 612,116.  
CODEN: USXXCO  
DT Patent  
LA English  
IC ICM A61K039-02  
ICS C12N001-21  
NCL 424200100  
CC 15-2 (Immunochemistry)  
Section cross-reference(s): 3, 14  
FAN.CNT 7

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI US 2002086032	A1	20020704	US 2001-927896	20010809 <--
PRAI US 1999-183043P	P	19990202	<--	
US 1999-198250P	P	19990505	<--	
US 2000-495614	A2	20000201		
US 2000-612116	A2	20000707		

#### CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
US 2002086032	ICM	A61K039-02
	ICS	C12N001-21
	NCL	424200100
US 2002086032	ECLA	A61K039/002; A61K039/02; A61K039/106; A61K039/112 <--
AB	The present invention is directed towards methods of inducing antibodies using an attenuated strain of pathogenic bacteria (e.g., Haemophilus, Escherichia coli, and/or Salmonella) having non-reverting genetic mutations relative to the wild-type organism which alter activity of DNA adenine methylase (Dam). The invention further includes compns. comprised of the attenuated bacteria and methods using these compns. to elicit an immune response and immunize a subject with highly specific antibodies. The invention also provides methods producing antibodies to heterologous antigens which the attenuated bacteria are engineered to produce.	
ST	antibody bacteria DNA adenine methylase; vaccine Salmonella vector DNA adenine methylase	
IT	<b>Vaccines</b> (AIDS; attenuated bacteria with altered DNA adenine methylase activity for expression of heterologous antigens of HIV)	
IT	Antibodies and Immunoglobulins RL: BSU (Biological study, unclassified); BIOL (Biological study) (IgG; attenuated bacteria with altered DNA adenine methylase activity for induction of)	
IT	Animal virus Arbovirus Ascaris lumbricoides Aspergillus fumigatus Astrovirus Bacillus anthracis Blastomyces dermatitidis Bordetella pertussis Borrelia burgdorferi Campylobacter Candida Chlamydia pneumoniae Chlamydia trachomatis	

*Clostridium tetani*  
*Coccidioides*  
*Cryptococcus neoformans*  
*Cytomegalovirus*  
*Dengue virus*  
*Entamoeba histolytica*  
*Giardia lamblia*  
*Helicobacter pylori*  
*Hepatitis A virus*  
*Hepatitis B virus*  
*Hepatitis C virus*  
*Hepatitis E virus*  
*Hepatitis GB virus C/G*  
*Hepatitis delta virus*  
*Histoplasma capsulatum*  
*Human coxsackievirus*  
*Human echovirus*  
*Human herpesvirus 2*  
*Human herpesvirus 3*  
*Human herpesvirus 4*  
*Human papillomavirus*  
*Human parainfluenza virus*  
*Human poliovirus*  
*Influenza virus*  
*Japanese encephalitis virus*  
*Leptospira*  
*Measles virus*  
*Moraxella catarrhalis*  
*Mycobacterium leprae*  
*Mycobacterium tuberculosis*  
*Mycoplasma pneumoniae*  
*Neisseria gonorrhoeae*  
*Neisseria meningitidis*  
*Norwalk virus*  
*Paracoccidioides brasiliensis*  
*Paramyxovirus*  
*Parasite*  
*Pinworm*  
*Plasmodium* (malarial genus)  
*Pseudomonas aeruginosa*  
*Rabies virus*  
*Respiratory syncytial virus*  
*Rhinovirus*  
*Rotavirus*  
*Rubella virus*  
*Schistosoma*  
*Staphylococcus aureus*  
*Staphylococcus saprophyticus*  
*Streptococcus group A*  
*Streptococcus group B*  
*Taenia*  
*Toxoplasma gondii*  
*Treponema pallidum*  
*Trichomonas vaginalis*  
*Variola virus*

(attenuated bacteria with altered DNA adenine methylase activity for expression of heterologous antigens of)

- IT Mycosis
  - (attenuated bacteria with altered DNA adenine methylase activity for expression of heterologous antigens of fungi associated with)
- IT Sexually transmitted diseases
  - (attenuated bacteria with altered DNA adenine methylase activity for expression of heterologous antigens of microorganisms associated with)
- IT Gene, microbial
  - RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
  - (dam; of attenuated bacteria with altered DNA adenine methylase activity)
- IT Antigens
  - RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
  - (heterologous; expression in attenuated bacteria with altered DNA adenine methylase activity)
- IT *Escherichia*
  - Eubacteria*
    - Haemophilus*
    - Salmonella*
    - Vibrio*

Yersinia  
(immunostimulation by attenuated bacteria with altered DNA adenine methylase activity)

IT **Vaccines**  
(of attenuated bacteria with altered DNA adenine methylase activity)

IT **Antigens**  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(tumor-associated; expression in attenuated bacteria with altered DNA adenine methylase activity)

IT **Bos taurus**  
**Gallus domesticus**  
**Human**  
(vaccination with attenuated bacteria expressing altered DNA adenine methylase activity and heterologous antigens)

IT **Food poisoning**  
(vaccination with attenuated bacteria expressing altered DNA adenine methylase activity and heterologous antigens in relation to)

IT **Anti-AIDS agents**  
(vaccines; attenuated bacteria with altered DNA adenine methylase activity for expression of heterologous antigens of HIV)

IT 69553-52-2, Dam methylase  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(immunostimulation by attenuated bacteria with altered DNA adenine methylase activity)

L44 ANSWER 2 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
AN 2002:488067 HCAPLUS  
DN 137:62150  
ED Entered STN: 28 Jun 2002  
TI Bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial or antibacterial agents  
IN Mahan, Michael J.; Heithoff, Douglas M.; Low, David A.; Sinsheimer, Robert L.  
PA USA  
SO U.S. Pat. Appl. Publ., 44 pp., Cont.-in-part of U.S. Ser. No. 612,116.  
CODEN: USXXCO  
DT Patent  
LA English  
IC ICM A61K039-02  
ICS C12N001-21; C12N015-74  
NCL 424200100  
CC 15-2 (Immunochemistry)  
Section cross-reference(s): 1, 3, 10, 17  
FAN.CNT 7

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002081317	A1	20020627	US 2001-927788	20010809 <--
	ZA 2001005305	A	20020627	ZA 2001-5305	20010627 <--
PRAI	US 1999-183043P	P	19990202	<--	
	US 1999-241951	A	19990202	<--	
	US 1999-198250P	P	19990505	<--	
	US 1999-305603	A	19990505	<--	
	US 2000-495614	A2	20000201		
	US 2000-612116	A2	20000707		

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
US 2002081317	ICM	A61K039-02
	ICS	C12N001-21; C12N015-74
	NCL	424200100
US 2002081317	ECLA	A61K039/002; A61K039/02; A61K039/106; A61K039/112 <--

AB Immunogenic compns. are disclosed which are comprised of bacteria which are pathogenic in their native state but which are rendered non-pathogenic in a manner which alters the native level or activity of DNA adenine methylase (dam). The genome is also artificially engineered to express a heterologous antigen such as an immunogenic antigen of a virus, protozoa, parasite or fungi. The microorganism with mutated dam is also useful for identifying or developing antimicrobial or antibacterial agents.

ST DNA adenine methylase mutation microorganism vaccine; antimicrobial antibacterial vaccine antigen DNA adenine methylase mutation

IT Hepatitis  
(A; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Hepatitis  
(B; bacteria with mutated DNA adenine methylase for use as vaccine and



screening or development of antimicrobial and antibacterial agents)

IT Hepatitis  
(C; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Hepatitis  
(D; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Hepatitis  
(E; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Antibodies and Immunoglobulins  
RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(IgG; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Haemophilus influenzae  
(NT; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Toxins  
RL: ADV (Adverse effect, including toxicity); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(Shiga; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Drug screening  
(antibacterial; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Fungi  
Parasite  
Protozoa  
Virus  
(antigen; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Arbovirus  
Ascaris lumbricoides  
Aspergillus fumigatus  
Astrovirus  
Bacillus anthracis  
Blastomyces dermatitidis  
Bordetella pertussis  
Borrelia burgdorferi  
Bos taurus  
Campylobacter  
Candida  
Chlamydia pneumoniae  
Chlamydia trachomatis  
Coccidioides immitis  
Cryptococcus neoformans  
Cytomegalovirus  
Dengue virus  
Drug delivery systems  
Entamoeba histolytica  
Eubacteria  
Food poisoning  
Gallus domesticus  
Genetic vectors  
Giardia lamblia  
Helicobacter pylori  
Hepatitis GB virus C/G  
Hepatitis virus  
Herpesviridae  
Histoplasma capsulatum  
Human  
Human coxsackievirus  
Human echovirus  
Human herpesvirus  
Human herpesvirus 1  
Human herpesvirus 2  
Human herpesvirus 3  
Human herpesvirus 4  
Human immunodeficiency virus  
Human papillomavirus  
Human parainfluenza virus  
Human poliovirus  
Influenza virus

Japanese encephalitis virus  
 Leptospira  
 Mammalia  
 Measles virus  
 Molecular cloning  
   *Moraxella catarrhalis*  
 Mutagenesis  
 Mycobacterium leprae  
 Mycobacterium tuberculosis  
 Mycoplasma pneumoniae  
 Mycosis  
 Neisseria gonorrhoeae  
 Neisseria meningitidis  
 Norwalk virus  
 Paracoccidioides brasiliensis  
 Paramyxovirus  
 Pathogen  
 Pathogenic bacteria  
 Pinworm  
 Plasmodium (malarial genus)  
 Pseudomonas aeruginosa  
 Rabies virus  
 Respiratory syncytial virus  
 Rhinovirus  
 Rodentia  
 Rotavirus  
 Rubella virus  
 Salmonella  
 Salmonella enteritidis  
 Salmonella typhi  
 Salmonella typhimurium  
 Schistosoma  
 Sexually transmitted diseases  
 Shigella  
 Staphylococcus saprophyticus  
 Streptococcus group A  
 Streptococcus group B  
 Taenia  
 Tetanus  
 Toxoplasma gondii  
 Treponema pallidum  
 Trichomonas vaginalis  
 Typhoid fever  
   Vaccines  
 Vibrio  
 Vibrio cholerae  
 Yersinia  
 Yersinia pseudotuberculosis  
   (bacteria with mutated DNA adenine methylase for use as vaccine and  
   screening or development of antimicrobial and antibacterial agents)  
 IT Antibodies and Immunoglobulins  
   Antigens  
   RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified);  
   THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES  
   (Uses)  
   (bacteria with mutated DNA adenine methylase for use as vaccine and  
   screening or development of antimicrobial and antibacterial agents)  
 IT Gene, microbial  
   RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL  
   (Biological study); USES (Uses)  
   (bacteria with mutated DNA adenine methylase for use as vaccine and  
   screening or development of antimicrobial and antibacterial agents)  
 IT Infection  
   (bacterial; bacteria with mutated DNA adenine methylase for use as  
   vaccine and screening or development of antimicrobial and antibacterial  
   agents)  
 IT Antibacterial agents  
   Antimicrobial agents  
   (development; bacteria with mutated DNA adenine methylase for use as  
   vaccine and screening or development of antimicrobial and antibacterial  
   agents)  
 IT Immunity  
   (disorder, antigen; bacteria with mutated DNA adenine methylase for use  
   as vaccine and screening or development of antimicrobial and  
   antibacterial agents)  
 IT Escherichia coli

(enterotoxigenic; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Respiratory tract, disease  
Urinary tract, disease  
(infection; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Intestinal bacteria  
(pathogenic, infection; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT **Antigens**  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(tumor-associated; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT **Haemophilus influenzae**  
(type b; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT **Escherichia coli**  
(uropathogenic; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Infection  
(variola; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Infection  
(vector-born; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT Infection  
(viral; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT 69553-52-2, Dam methylase  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(mutation; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial and antibacterial agents)

IT 439624-92-7 439624-93-8 439624-94-9 439624-95-0  
RL: PRP (Properties)  
(unclaimed sequence; bacteria with mutated DNA adenine methylase for use as vaccine and screening or development of antimicrobial or antibacterial agents)

L44 ANSWER 3 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 2002:107146 HCAPLUS

DN 136:166052

ED Entered STN: 10 Feb 2002

TI Vaccine composition

IN Berthet, Francois-Xavier Jacques; Dalemans, Wilfried; Denoel, Philippe; Dequesne, Guy; Feron, Christiane; Garcon, Nathalie; Lobet, Yves; Poolman, Jan; Thiry, Georges; Thonnard, Joelle; Voet, Pierre

PA Smithkline Beecham Biologicals SA, Belg.

SO PCT Int. Appl., 125 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K039-00

CC 15-2 (Immunochemistry)

Section cross-reference(s): 3

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002009746	A2	20020207	WO 2001-EP8857	20010731
	WO 2002009746	A3	20020613		
	WO 2002009746	C1	20021114		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				

Search done by Noble Jarrell

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,  
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,  
 BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

EP 1208214 A2 20020529 EP 2000-956369 20000731 <--  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL

CA 2425037 AA 20020207 CA 2001-2425037 20010731  
 AU 2001085856 A5 20020213 AU 2001-85856 20010731  
 EP 1307224 A2 20030507 EP 2001-965152 20010731  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR

US 2004126389 A1 20040701 US 2003-343561 20030915  
 PRAI EP 2000-956369 A 20000731  
 GB 2001-3170 A 20010208  
 GB 1999-18319 A 19990803 <--  
 WO 2000-EP7424 W 20000731  
 WO 2001-EP8857 W 20010731

## CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 2002009746	ICM	A61K039-00
US 2004126389	ECLA	A61K039/02; A61K039/39

AB The present invention relates to the field of vaccine formulation, particularly the field of novel adjuvant compns. comprising outer membrane vesicles (or blebs), and advantageous methods of detoxifying these compns., and advantageous methods of use of such adjuvants. The novel adjuvant for Gram-neg. bacterial vaccine is a capsular polysaccharide or detoxified lipid A portion of LPS derived from engineered Neisseria meningitidis serogroup A, B, Y or W; Hemophilus influenzae; Streptococcus pneumoniae; or Moraxella catarrhalis. These engineered bacteria have reduced or switched off expression of one or more gene selected from htrB, msbB, .pxK, pmrA, pmrB, pmrE, pmrF, galE, siaA, siaB, siaC, siaD, ctrA, ctrB, ctrC and ctrD. Vaccines comprising the adjuvant and pathogen-derived antigen is especially useful for protecting elderly patients against the pathogen.

ST vaccine adjuvant outer membrane vesicle bleb; Gram neg bacteria bleb prepn adjuvant; Neisseria meningitidis bleb detoxified lipid A; Streptococcus pneumoniae Hemophilus influenzae capsular polysaccharide

IT Gene, microbial  
 RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
 (D15; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Proteins  
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (D; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Proteins  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (OMP (outer membrane protein); outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Immunostimulants  
 (adjuvants; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Organelle  
 (bleb; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Glycolipids  
 Lipopolysaccharides  
 Polysaccharides, biological studies  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (capsular; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Drug delivery systems  
 (carriers; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
 RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
 (cps; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(ctrA; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(ctrB; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(ctrC; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(ctrD; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Aging, animal  
(elderly; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(galE; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Proteins  
RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation)  
(green fluorescent; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Neisseria meningitidis  
(group A; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Neisseria meningitidis  
(group B; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Neisseria meningitidis  
(group W; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Neisseria meningitidis  
(group Y; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(hsf; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(htrB; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(lpxK; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(msbB; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(nspA; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial  
RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
(omp85; outer membrane vesicles or detoxified lipid A as adjuvant for Gram-neg. bacterial vaccines)

IT DNA sequences  
Detergents

Gram-negative bacteria  
*Haemophilus influenzae*  
*Moraxella catarrhalis*  
*Neisseria meningitidis*  
*Streptococcus pneumoniae*

**Vaccines**

(outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Promoter (genetic element)

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT **Antigens**

Polysaccharides, biological studies

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL  
 (Biological study); USES (Uses)  
 (outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Cell wall

(outer membrane, vesicles; outer membrane vesicles or detoxified lipid  
 A as adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (pilQ; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (pIdA; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (pmrA; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (pmrB; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (pmrE; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (pmrF; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Infection

(pneumococcal; outer membrane vesicles or detoxified lipid A as  
 adjuvant for Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (porA; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (porB; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (siaA; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (siaB; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial

RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL

(Biological study); PROC (Process)  
 (siaC; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial  
 RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (siaD; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT Gene, microbial  
 RL: BSU (Biological study, unclassified); REM (Removal or disposal); BIOL  
 (Biological study); PROC (Process)  
 (tbpA; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT *Haemophilus influenzae*  
 (type b; outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT 1404-24-6, Polymyxin A  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL  
 (Biological study); USES (Uses)  
 (chimeric; outer membrane vesicles or detoxified lipid A as adjuvant  
 for Gram-neg. bacterial vaccines)

IT 397430-36-3, DNA (Synthetic plasmid vector CMK(+))  
 RL: BSU (Biological study, unclassified); BUU (Biological use,  
 unclassified); PRP (Properties); BIOL (Biological study); USES (Uses)  
 (nucleotide sequence; outer membrane vesicles or detoxified lipid A as  
 adjuvant for Gram-neg. bacterial vaccines)

IT 397430-37-4 397430-38-5 397430-39-6 397430-40-9 397430-41-0  
 397430-42-1  
 RL: BSU (Biological study, unclassified); PRP (Properties); REM (Removal  
 or disposal); BIOL (Biological study); PROC (Process)  
 (nucleotide sequence; outer membrane vesicles or detoxified lipid A as  
 adjuvant for Gram-neg. bacterial vaccines)

IT 83-44-3  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
 (Uses)  
 (outer membrane vesicles or detoxified lipid A as adjuvant for  
 Gram-neg. bacterial vaccines)

IT 397431-26-4, 4: PN: WO0209746 SEQID: 4 unclaimed DNA 397431-27-5, 5: PN:  
 WO0209746 SEQID: 5 unclaimed DNA 397431-28-6, 6: PN: WO0209746 SEQID: 6  
 unclaimed DNA 397431-29-7, 8: PN: WO0209746 SEQID: 8 unclaimed DNA  
 397431-30-0, 9: PN: WO0209746 SEQID: 9 unclaimed DNA 397431-31-1  
 397431-32-2 397431-33-3 397431-34-4 397431-35-5 397431-36-6  
 397431-37-7 397431-38-8 397431-39-9 397431-40-2 397431-41-3  
 397431-42-4 397431-43-5 397431-44-6 397431-45-7 397431-46-8  
 397431-47-9 397431-48-0 397431-49-1 397431-50-4 397431-51-5  
 397431-52-6 397431-53-7 397431-54-8 397431-55-9 397431-56-0  
 397431-57-1 397431-58-2 397431-59-3 397431-60-6 397431-61-7  
 397431-62-8 397431-63-9 397431-64-0 397431-65-1 397431-66-2  
 397431-67-3 397431-68-4 397431-69-5 397431-70-8 397431-71-9  
 397431-72-0 397431-73-1 397431-74-2 397431-75-3 397431-76-4  
 397431-77-5 397431-78-6 397431-79-7 397431-80-0 397431-81-1  
 397431-82-2 397431-83-3 397431-84-4 397431-85-5 397431-86-6  
 397431-87-7 397431-88-8 397431-89-9 397431-90-2 397431-91-3  
 397431-92-4 397431-93-5 397431-94-6 397431-95-7 397431-96-8  
 397431-97-9 397431-98-0 397431-99-1 397432-00-7 397432-01-8  
 397432-02-9 397432-03-0 397432-04-1 397432-05-2 397432-06-3  
 397432-07-4 397432-08-5 397432-09-6 397432-10-9 397432-11-0  
 397432-12-1 397432-13-2 397432-14-3 397432-15-4 397432-16-5  
 397432-17-6 397432-18-7 397432-19-8 397432-20-1 397432-21-2  
 397432-22-3 397432-23-4 397432-24-5 397432-25-6 397432-26-7  
 397432-27-8 397432-28-9 397432-29-0 397432-30-3 397432-31-4  
 397432-32-5 397432-33-6 397432-34-7 397432-35-8 397432-36-9  
 397432-37-0 397432-38-1 397432-39-2 397432-40-5 397432-41-6  
 397432-42-7 397432-43-8 397432-44-9 397432-45-0 397432-46-1  
 397432-47-2 397432-48-3 397432-49-4 397432-50-7 397432-51-8  
 397432-52-9 397432-53-0 397432-54-1 397432-55-2 397432-56-3  
 397432-57-4 397432-58-5 397432-59-6 397432-60-9 397432-61-0  
 397432-62-1 397432-63-2 397432-64-3 397432-65-4 397432-66-5  
 397432-67-6 397432-68-7 397432-69-8 397432-70-1 397432-71-2  
 397432-72-3 397432-73-4 397432-74-5  
 RL: PRP (Properties)  
 (unclaimed nucleotide sequence; vaccine composition)

L44 ANSWER 4 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 2000:688113 HCAPLUS  
 DN 133:265640

Search done by Noble Jarrell

ED Entered STN: 29 Sep 2000  
 TI Bacterial polysaccharide antigen vaccine  
 IN Capiiau, Carine; Deschamps, Marguerite; Desmons, Pierre Michel; Laferriere, Craig Antony Joseph; Poolman, Jan; Prieels, Jean-paul  
 PA Smithkline Beecham Biologicals SA, Belg.  
 SO PCT Int. Appl., 79 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM A61K039-385  
 ICS A61K039-39; A61K039-02; A61K039-005; A61K039-116; A61P031-04  
 CC 15-2 (Immunochemistry)  
 Section cross-reference(s): 3  
 FAN.CNT 5

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000056360	A2	20000928	WO 2000-EP2468	20000317 <--
WO 2000056360	A3	20010125		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2366314	AA	20000928	CA 2000-2366314	20000317 <--
NZ 513841	A	20010928	NZ 2000-513841	20000317 <--
EP 1163000	A2	20011219	EP 2000-912626	20000317 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
TR 200102739	T2	20011221	TR 2001-200102739	20000317 <--
BR 2000009163	A	20011226	BR 2000-9163	20000317 <--
TR 200102735	T2	20020422	TR 2001-200102735	20000317 <--
TR 200102736	T2	20020422	TR 2001-200102736	20000317 <--
AU 750913	B2	20020801	AU 2000-34307	20000317 <--
AU 2000034307	A5	20001009		
JP 2002540075	T2	20021126	JP 2000-606264	20000317 <--
NZ 513840	A	20040227	NZ 2000-513840	20000317 <--
NZ 513842	A	20040528	NZ 2001-513842	20010317 <--
NO 2001004325	A	20011114	NO 2001-4325	20010905 <--
ZA 2001007638	A	20020611	ZA 2001-7638	20010917 <--
ZA 2001007637	A	20020621	ZA 2001-7637	20010917 <--
ZA 2001007640	A	20020911	ZA 2001-7640	20010917 <--
PRAI GB 1999-6437	A	19990319	<--	
GB 1999-9077	A	19990420	<--	
GB 1999-9466	A	19990423	<--	
GB 1999-16677	A	19990715	<--	
WO 2000-EP2468	W	20000317		

## CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 2000056360	ICM	A61K039-385
	ICS	A61K039-39; A61K039-02; A61K039-005; A61K039-116; A61P031-04
AB	The present invention relates to the field of bacterial polysaccharide antigen vaccines. In particular, the present invention relates to bacterial polysaccharides conjugated to protein D from H. influenzae.	
ST	bacteria polysaccharide antigen vaccine protein D	
IT	Neisseria meningitidis	
	(C; bacterial polysaccharide antigen vaccine)	
IT	Proteins, specific or class	
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)	
	(Cbpa or choline-binding protein A; bacterial polysaccharide antigen vaccine)	
IT	Proteins, specific or class	
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)	
	(D; bacterial polysaccharide antigen vaccine)	
IT	Proteins, specific or class	
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)	
	(PsaA; bacterial polysaccharide antigen vaccine)	
IT	Proteins, specific or class	
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)	
	(PspA (pneumococcal surface protein A); bacterial polysaccharide antigen vaccine)	



IT Proteins, specific or class  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(PspC; bacterial polysaccharide antigen vaccine)

IT Immunity  
(Th1 adjuvant; bacterial polysaccharide antigen vaccine)

IT Polysaccharides, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(Vi; bacterial polysaccharide antigen vaccine)

IT Neisseria meningitidis  
(Y; bacterial polysaccharide antigen vaccine)

IT Immunostimulants  
(adjuvants, Th1; bacterial polysaccharide antigen vaccine)

IT Haemophilus influenzae  
Immunostimulants  
Pathogen  
Salmonella typhi  
Susceptibility (genetic)  
Trypanosoma cruzi  
Vaccines  
(bacterial polysaccharide antigen vaccine)

IT Lipopolysaccharides  
Saponins  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(bacterial polysaccharide antigen vaccine)

IT Infection  
(bacterial; bacterial polysaccharide antigen vaccine)

IT Polysaccharides, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(capsular; bacterial polysaccharide antigen vaccine)

IT Antigens  
Polysaccharides, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(conjugates; bacterial polysaccharide antigen vaccine)

IT Glycolipoproteins  
Glycolipoproteins  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(glycan-containing, phospho-; bacterial polysaccharide antigen vaccine)

IT Neisseria meningitidis  
(group B polysaccharide; bacterial polysaccharide antigen vaccine)

IT Oligosaccharides, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(lipopeptidophospho; bacterial polysaccharide antigen vaccine)

IT Moraxella catarrhalis  
Shigella sonnei  
(lipopolysaccharide; bacterial polysaccharide antigen vaccine)

IT Mucopolysaccharides, biological studies  
Mucopolysaccharides, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(lipoproteoglycans, phospho-; bacterial polysaccharide antigen vaccine)

IT Proteins, specific or class  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(outer surface proteins; bacterial polysaccharide antigen vaccine)

IT Lipopeptides  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(phosphoglycan; bacterial polysaccharide antigen vaccine)

IT Hemolysins  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(pneumolysins; bacterial polysaccharide antigen vaccine)

IT Bacteria (Eubacteria)  
Cryptococcus neoformans  
Mycobacterium  
Neisseria meningitidis  
Staphylococcus aureus  
Streptococcus agalactiae  
Streptococcus pneumoniae  
(polysaccharide; bacterial polysaccharide antigen vaccine)

IT Proteins, specific or class  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(secreted; bacterial polysaccharide antigen vaccine)

IT Oligonucleotides  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(stimulatory CpG-containing; bacterial polysaccharide antigen vaccine)

IT Toxoids  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(tetanus; bacterial polysaccharide antigen vaccine)

IT 7784-30-7, Aluminum phosphate 9001-50-7, Glyceraldehyde-3-phosphate

dehydrogenase 128478-31-9

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(bacterial polysaccharide antigen vaccine)

L44 ANSWER 5 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 2000:68366 HCAPLUS  
 DN 132:127726  
 ED Entered STN: 28 Jan 2000  
 TI Adjuvant and vaccine compositions containing monophosphoryl lipid A  
 IN Laposta, Vincent James; Eldridge, John Hayward  
 PA American Cyanamid Company, USA  
 SO PCT Int. Appl., 35 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM A61K039-00  
 CC 63-6 (Pharmaceuticals)  
 Section cross-reference(s): 15

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000003744	A2	20000127	WO 1999-US15942	19990713 <--
WO 2000003744	A3	20000427		
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2333578	AA	20000127	CA 1999-2333578	19990713 <--
AU 9951018	A1	20000207	AU 1999-51018	19990713 <--
AU 763770	B2	20030731		
BR 9912067	A	20010410	BR 1999-12067	19990713 <--
EP 1096954	A2	20010509	EP 1999-935566	19990713 <--
EP 1096954	B1	20041006		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
US 6306404	B1	20011023	US 1999-352526	19990713 <--
JP 2002520373	T2	20020709	JP 2000-559878	19990713 <--
US 2002025330	A1	20020228	US 2001-943028	20010830 <--
US 6635261	B2	20031021		
PRAI US 1998-115392	A	19980714	<--	
US 1998-155270P	P	19980714	<--	
US 1999-352526	A3	19990713	<--	
WO 1999-US15942	W	19990713	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 2000003744	ICM	A61K039-00
US 2002025330	ECLA	A61K039/39 <--
AB	The invention pertains to adjuvant and vaccine compns. of monophosphoryl lipid A, sugar, and optionally an amine-based surfactant, which when frozen and thawed or lyophilized and reconstituted reform a colloidal suspension having a light transmission of greater than or equal to 88 % as measured spectrophotometrically.	
ST	vaccine adjuvant formulation monophosphoryl lipid A	
IT	Chlamydia	
	Colloids	
	Haemophilus influenzae	
	Helicobacter pylori	
	Human herpesvirus	
	Human immunodeficiency virus	
	Human papillomavirus	
	Human parainfluenza virus	
	Influenza virus	
	Measles virus	
	Moraxella catarrhalis	
	Neisseria gonorrhoeae	
	Neisseria meningitidis	
	Norwalk virus	
	Optical absorption	
	Respiratory syncytial virus	
	Rotavirus	
	Salmonella typhi	

Search done by Noble Jarrell

Solvents  
 Spectrophotometry  
 Streptococcus group A  
 Streptococcus group B  
 Streptococcus pneumoniae  
 Turbidimetry  
 Vaccines  
 (adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT Allergens  
   Antigens  
     RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
     (adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT Carbohydrates, biological studies  
     RL: PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
     (adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT Immunostimulants  
     (adjuvants; adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT Surfactants  
     (amine-based; adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT Bacteria (Eubacteria)  
   Neoplasm  
   Parasite  
   Virus  
     (antigens of; adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT Polysaccharides, biological studies  
     RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
     (capsular; adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT Drug delivery systems  
     (carriers; adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT Physiological saline solutions  
     (diluent; adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT Drug delivery systems  
     (freeze-dried; adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT Lipopolysaccharides  
     RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
     (of Salmonella minnesota R595; adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT 7784-30-7, Aluminum phosphate 220048-47-5  
     RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
     (adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT 50-99-7, Dextrose, biological studies 57-48-7, D-Fructose, biological studies 57-50-1, biological studies 59-23-4, Galactose, biological studies 63-42-3, Lactose 69-79-4, Maltose 99-20-7, Trehalose 499-40-1, Isomaltose 3458-28-4, Mannose  
     RL: PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
     (adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT 7732-18-5, Water, uses  
     RL: NUU (Other use, unclassified); USES (Uses)  
     (diluent; adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 IT 102-71-6, biological studies 121-44-8, biological studies  
     RL: MOA (Modifier or additive use); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
     (surfactant; adjuvant and vaccine compns. containing monophosphoryl lipid A)  
 L44 ANSWER 6 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 1999:626070 HCAPLUS  
 DN 131:262583  
 ED Entered STN: 01 Oct 1999  
 TI Haemophilus influenzae B-DTPa combination vaccine  
 IN Artois, Claude; De Heyder, Koen; Desmons, Pierre; Garcon, Nathalie;

Mainil, Roland  
 PA Smithkline Beecham Biologicals SA, Belg.  
 SO PCT Int. Appl., 36 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM A61K039-295  
 ICS A61K039-39; A61K039-05; A61K039-08; A61K039-09; A61K039-095;  
 A61K039-10; A61K039-102; A61K039-13; A61K039-29  
 CC 63-3 (Pharmaceuticals)

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9948525	A1	19990930	WO 1999-EP1959	19990322 <--
	W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	CA 2325436	AA	19990930	CA 1999-2325436	19990322 <--
	AU 9934172	A1	19991018	AU 1999-34172	19990322 <--
	AU 735619	B2	20010712		
	BR 9909037	A	20001205	BR 1999-9037	19990322 <--
	TR 200002737	T2	20001221	TR 2000-200002737	19990322 <--
	EP 1066053	A1	20010110	EP 1999-915692	19990322 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI				
	JP 2002507581	T2	20020312	JP 2000-537572	19990322 <--
	NZ 506604	A	20030228	NZ 1999-506604	19990322 <--
	ZA 2000004956	A	20020108	ZA 2000-4956	20000918 <--
	NO 2000004758	A	20001108	NO 2000-4758	20000922 <--
	US 2003022304	A1	20030130	US 2002-217572	20020813 <--
PRAI	GB 1998-6456	A	19980325	<--	
	WO 1999-EP1959	W	19990322	<--	
	US 2000-647032	B1	20001031		

## CLASS

	PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
	WO 9948525	ICM	A61K039-295
		ICS	A61K039-39; A61K039-05; A61K039-08; A61K039-09; A61K039-095; A61K039-10; A61K039-102; A61K039-13; A61K039-29
	US 2003022304	ECLA	A61K039/05; A61K039/08; A61K039/09; A61K039/095; A61K039/10; A61K039/102; A61K039/13; A61K039/29; A61K039/39 <--
AB	This invention relates to a general method by which either extemporaneously prepared or liquid Haemophilus influenzae B (Hib)/DTPa combination vaccines can be made in order to avoid Hib interference while being able to maintain the maximum, stable adsorption of each antigen onto the aluminum-based adjuvant on which it is most immunogenic. In so doing, pertussis antigens in combination vaccines of the present invention are stably retained in their most potent form. Examples are given for the vaccines using Al hydroxide or Al phosphate as adjuvants.		
ST	vaccine Haemophilus diphtheria tetanus pertussis		
IT	<b>Vaccines</b>		
	(Haemophilus influenzae B-DTPa combination vaccine)		
IT	<b>Antigens</b>		
	Polysaccharides, biological studies		
	RL: BOC (Biological occurrence); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); USES (Uses)		
	(Haemophilus influenzae B-DTPa combination vaccine)		
IT	<b>Immunostimulants</b>		
	(adjuvants; Haemophilus influenzae B-DTPa combination vaccine)		
IT	Hepatitis A virus		
	Human poliovirus		
	(antigens; Haemophilus influenzae B-DTPa combination vaccine)		
IT	Streptococcus pneumoniae		
	(capsular polysaccharide and proteins; Haemophilus influenzae B-DTPa combination vaccine)		
IT	<b>Toxoids</b>		
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)		

(diphtheria; Haemophilus influenzae B-DTPa combination vaccine)

IT *Neisseria meningitidis*  
(group A, capsular polysaccharide; Haemophilus influenzae B-DTPa combination vaccine)

IT *Neisseria meningitidis*  
(group C, capsular polysaccharide; Haemophilus influenzae B-DTPa combination vaccine)

IT **Antigens**  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(hepatitis B surface; Haemophilus influenzae B-DTPa combination vaccine)

IT *Moraxella catarrhalis*  
(outer membrane proteins; Haemophilus influenzae B-DTPa combination vaccine)

IT Toxoids  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(pertussis; Haemophilus influenzae B-DTPa combination vaccine)

IT Toxoids  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(tetanus; Haemophilus influenzae B-DTPa combination vaccine)

IT *Haemophilus influenzae*  
(type b; Haemophilus influenzae B-DTPa combination vaccine)

IT 7784-30-7, Aluminum phosphate 21645-51-2, Aluminum hydroxide, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(Haemophilus influenzae B-DTPa combination vaccine)

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE  
(1) Corbel, M; BIOLOGICALS 1994, V22(4), P353 HCAPLUS  
(2) Corbel, M; BIOLOGICALS 1997, V25/3, P351  
(3) Ellis, R; VACCINE 1999, V17(13-14), P1635 MEDLINE  
(4) Slaoui Moncef Mohamed; WO 9746255 A 1997 HCAPLUS  
(5) Smithkline Beecham Biolog; WO 9324148 A 1993 HCAPLUS  
(6) Smithkline Beecham Biolog; WO 9700697 A 1997 HCAPLUS

L44 ANSWER 7 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1999:597423 HCAPLUS

DN 131:213104

ED Entered STN: 22 Sep 1999

TI Antigenic conjugates of conserved lipopolysaccharides of gram negative bacteria

IN Arumugham, Rasappa G.; Fortuna-Nevin, Maria; Apicella, Michael A.; Gibson, Bradford W.

PA American Cyanamid Company, USA

SO Eur. Pat. Appl., 18 pp.

CODEN: EPXXDW

DT Patent

LA English

IC ICM A61K039-385

ICI A61K039-02, A61K039-095

CC 15-2 (Immunochemistry)

Section cross-reference(s): 14, 63

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 941738	A1	19990915	EP 1999-301747	19990309 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
CA 2264970	AA	19990910	CA 1999-2264970	19990308 <--
AU 9919540	A1	19990923	AU 1999-19540	19990309 <--
AU 766184	B2	20031009		
JP 11322793	A2	19991124	JP 1999-61354	19990309 <--
BR 9902008	A	20000509	BR 1999-2008	19990309 <--
PRAI US 1998-37529	A	19980310	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
EP 941738	ICM	A61K039-385
	ICI	A61K039-02, A61K039-095

AB Antigenic conjugates are provided which comprise a carrier protein covalently bonded to the conserved portion of a lipopolysaccharide of a gram neg. bacteria, wherein said conserved portion of the lipopolysaccharide comprises the inner core and lipid A portions of said lipopolysaccharide, said conjugate eliciting a cross reactive immune response against heterologous strains of said gram neg. bacteria. The carrier protein is selected from CRM197, tetanus toxin, diphtheria toxin,

pseudomonas exotoxin A, cholera toxin, group A streptococcal toxin, pneumolysin of Streptococcus pneumoniae, filamentous hemagglutinin (FHA), FHA of Bordetella pertussis, pili or pilins of Neisseria gonorrhoeae or meningitidis, outer membrane proteins of Neisseria meningitidis, C5A peptidase of Streptococcus and surface protein of Moraxella catarrhalis.

ST gram neg bacteria lipopolysaccharide carrier protein; vaccine lipopolysaccharide carrier conjugate immune adjuvant

IT Hemagglutinins  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (FHA (filamentous hemagglutinin); conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT Proteins, specific or class  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (OMP (outer membrane protein), carrier; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT Proteins, specific or class  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (SU (surface), carrier; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT Immunostimulants  
 (adjuvants; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT Toxins  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (bacterial; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT Streptococcus  
 (carrier; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT Toxins  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (cholera; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT Bordetella  
 Bordetella pertussis  
 Chlamydia  
 Escherichia coli  
 Gram-negative bacteria  
 Haemophilus  
 Haemophilus ducreyi  
 Haemophilus influenzae  
 Helicobacter pylori  
 Klebsiella  
 Moraxella catarrhalis  
 Neisseria  
 Neisseria gonorrhoeae  
 Neisseria meningitidis  
 Pilus  
 Proteus mirabilis  
 Pseudomonas  
 Pseudomonas aeruginosa  
 Salmonella  
 Salmonella minnesota  
 Salmonella typhimurium  
 Shigella  
 Streptococcus pneumoniae  
 Vaccines  
 Vibrio cholerae  
 (conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT Lipid A

**O antigen****Pilins**

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Lipopolysaccharides**

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Carriers**

(conjugates; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Antigens**

Proteins, specific or class

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(conjugates; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Toxins**

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(diphtheria, carrier; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Toxins**

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(exotoxin A, carrier; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Drug delivery systems**

(injections, i.m.; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Drug delivery systems**

(injections, i.v.; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Drug delivery systems**

(injections, s.c.; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Drug delivery systems**

(intradermal; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Drug delivery systems**

(nasal, intra-; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Drug delivery systems**

(ophthalmic; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Drug delivery systems**

(oral; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Hemolysins**

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(pneumolysins, carrier; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Drug delivery systems**

(solns., i.p.; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

**IT Toxins**

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(tetanus, carrier; conjugates of conserved lipopolysaccharides of gram

neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT Streptococcus group A  
(toxin; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT Drug delivery systems  
(vaginal; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT 100179-39-3, C5A Peptidase  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(carrier; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

IT 51-85-4, Cystamine 1071-93-8, Adipic acid dihydrazide 1892-57-5, EDAC 6539-14-6, Traut's reagent 42014-51-7, Bromoacetic acid N-hydroxysuccinimide ester 57757-57-0 64202-52-4 64987-85-5, SMCC 68181-17-9, SPDP 72252-96-1, SIAB 76931-93-6, SATA 79886-55-8, Succinimidyl 4-(p-maleimidophenyl)butyrate 150205-95-1 150244-18-1 158913-22-5  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(linker; conjugates of conserved lipopolysaccharides of gram neg. bacteria and carrier proteins for eliciting cross reactive immune response against heterologous strains of gram neg. bacteria)

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE  
(1) Baker, P; Infection and Immunity 1994, V62(6), P2257 HCAPLUS  
(2) Rune, A; Microbial Pathogenesis 1997, V23(3), P139  
(3) Stanislavsky, E; FEMS Microbiology Reviews 1997, V21(3), P243 HCAPLUS

L44 ANSWER 8 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
AN 1999:571730 HCAPLUS  
DN 131:213099  
ED Entered STN: 09 Sep 1999  
TI Vaccine for Moraxella catarrhalis  
IN Murphy, Timothy F.  
PA The Research Foundation of State University of New York, USA  
SO U.S., 20 pp., Cont.-in-part of U.S. 5,607,846.

CODEN: USXXAM  
DT Patent  
LA English  
IC ICM A61K039-02  
ICS C07K014-00

NCL 424251100  
CC 15-2 (Immunochemistry)  
Section cross-reference(s): 3

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 5948412	A	19990907	US 1997-810655	19970303 <--
	US 5607846	A	19970304	US 1994-245758	19940517 <--
	CA 2189971	AA	19951123	CA 1995-2189971	19950420 <--
	CA 2189971	C	20030729		
	ES 2202361	T3	20040401	ES 1995-917165	19950420 <--
PRAI	US 1994-245758	A2	19940517	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
US 5948412	ICM	A61K039-02
	ICS	C07K014-00
	NCL	424251100
US 5948412	ECLA	C07K014/21B
US 5607846	ECLA	C07K014/21B

AB Comps. comprising outer membrane protein E, and peptides and oligopeptides thereof, of Moraxella catarrhalis are described. Addnl., nucleotide sequences encoding the protein, peptide, or oligopeptide are disclosed, as well as recombinant vectors containing these sequences. Protein, peptide, or oligopeptide can be produced from host cell systems containing these recombinant vectors. Peptides and oligopeptides can also be chemical synthesized. Disclosed are the uses of the protein, peptides and oligopeptides as antigens in antigenic formulations for vaccine applications or for generating antisera of diagnostic or therapeutic use; and as antigens in diagnostic immunoassays. The nucleotide sequences are useful for constructing vectors for use as vaccines for insertions into



attenuated bacteria in constructing a recombinant bacterial vaccine and for inserting into a viral vector in constructing a recombinant viral vaccine. Also described is the use of nucleotide sequences related to the gene encoding E as primers and/or probes in mol. diagnostic assays for the detection of *M. catarrhalis*.

- ST Moraxella catarrhalis outer membrane protein E; vaccine antiserum outer membrane protein E; gene protein E epitope probe primer
- IT Primers (nucleic acid)  
 Primers (nucleic acid)  
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (DNA; vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)
- IT Proteins, specific or class  
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (OMP (outer membrane protein), E; vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)
- IT Polysaccharides, biological studies  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (capsular; vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)
- IT Drug delivery systems  
 (carriers; vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)
- IT Diagnosis  
 (immunodiagnosis; vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)
- IT DNA  
 DNA  
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (primer; vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)
- IT Animal cell line  
 Antiserums  
 Bacteria (Eubacteria)  
 DNA sequences  
 Epitopes  
 Filamentous fungi  
 Haemophilus influenzae  
 Insect (Insecta)  
 Mammal (Mammalia)  
 Molecular cloning  
 Moraxella catarrhalis  
 Neisseria meningitidis  
 Protein sequences  
 Pseudomonas aeruginosa  
 Staphylococcus aureus  
 Streptococcus pneumoniae  
 Vaccines  
 Virus vectors  
 Yeast  
 (vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)
- IT Gene, microbial  
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)
- IT Antigens  
 Lipopolysaccharides  
 Polysaccharides, biological studies  
 RNA  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)
- IT Probes (nucleic acid)  
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)

IT 159869-80-4 174065-50-0 242799-79-7  
 RL: PRP (Properties)  
 (amino acid sequence; vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)

IT 174066-42-3  
 RL: PRP (Properties)  
 (nucleotide sequence; vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)

IT 173432-99-0 173433-06-2 173433-09-5 173433-10-8 173433-11-9  
 173433-12-0 173433-13-1  
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (vaccine for Moraxella catarrhalis comprising outer membrane protein E epitope and primers and probes for genetic diagnosis)

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE  
 (1) Bartos; J Infect Dis 1988, V158, P761 HCAPLUS  
 (2) Bhushan; Abstracts Gen Meet Am Soc Microbiol 1991, V97, P30  
 (3) Maciver; J Infect Dis 1993, V168, P469 MEDLINE

L44 ANSWER 9 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 1999:487519 HCAPLUS  
 DN 131:120851  
 ED Entered STN: 06 Aug 1999  
 TI Nonrecombinant subunit vaccine  
 IN Gerlach, Gerald-F.; Goethe, Ralph  
 PA Germany  
 SO Ger. Offen., 22 pp.  
 CODEN: GWXXBX  
 DT Patent  
 LA German  
 IC ICM A61K039-102  
 ICS A61K039-095  
 CC 63-4 (Pharmaceuticals)  
 Section cross-reference(s): 15

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19753176	A1	19990729	DE 1997-19753176	19971120 <--
DE 19753176	C2	20000427		
PRAI DE 1997-19753176		19971120 <--		

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
DE 19753176	ICM	A61K039-102
	ICS	A61K039-095

AB The title bacterial vaccines are obtained by (1) cultivation of (preferably gram-neg.) pathogenic bacteria, preferably under mineral or nutrient deficiency stress or heat stress, and (2) enrichment of protective antigens from the bacteria by use of detergents, especially steroidal detergents such as cholic acid. This procedure exts. various protective antigens (especially lipoproteins) from the outer membrane without lysing the bacteria and thus without causing release of extraneous proteins. The subunit vaccine can be used as a marker vaccine for differentiation of vaccinated from infected subjects by ELISA. Thus, Actinobacillus pleuropneumoniae 811/051 (serotype 9) was cultivated in PPLO medium + Iso Vitale X at 37.degree. under Fe deficiency conditions (100 .mu.M 2,2'-dipyridyl), centrifuged, and resuspended in distilled water, and transferrin-binding protein A was extracted from the outer membrane with 0.075% Na deoxycholate. This extract and a similar extract from serotype 2 were combined 1:2, diluted 1:10, and mixed with HCHO 0.05 and Emulsigen Plus 20% for use as a vaccine in swine.

ST bacteria vaccine outer membrane protein; Actinobacillus vaccine detergent extn; pleuropneumonia vaccine extn deoxycholate

IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (OMP (outer membrane protein); nonrecombinant subunit vaccine)

IT Detergents  
 (anionic; nonrecombinant subunit vaccine)

IT Mineral elements, biological studies

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (bacteria deficiency in; nonrecombinant subunit vaccine)

IT Nutrition, microbial  
 (deficiency; nonrecombinant subunit vaccine)

IT Immunoassay  
 (enzyme-linked immunosorbent assay, for bacterial outer membrane proteins; nonrecombinant subunit vaccine)

IT Swine  
 (immunization of, against pleuropneumonia; nonrecombinant subunit vaccine)

IT Diagnosis  
 (immunodiagnosis, of pleuropneumonia; nonrecombinant subunit vaccine)

IT Detergents  
 (nonionic; nonrecombinant subunit vaccine)

IT Actinobacillus equuli  
 Actinobacillus pleuropneumoniae  
 Chelating agents  
 Detergents  
 Escherichia coli  
 Gram-negative bacteria  
 Haemophilus actinomycetemcomitans  
 Haemophilus agni  
 Haemophilus influenzae  
 Haemophilus paragallinarum  
 Haemophilus parasuis  
 Haemophilus somnus  
 Mannheimia haemolytica  
 Moraxella bovis  
 Moraxella catarrhalis  
 Moraxella lacunata  
 Neisseria gonorrhoeae  
 Neisseria meningitidis  
 Neisseriaceae  
 Pasteurella avium  
 Pasteurella multocida  
 Pasteurellaceae  
 Stress, microbial  
 Vaccines  
 (nonrecombinant subunit vaccine)

IT Antigens  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (nonrecombinant subunit vaccine)

IT Bile acids  
 Bile salts  
 Quaternary ammonium compounds, biological studies  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (nonrecombinant subunit vaccine)

IT Antibodies  
 RL: ANT (Analyte); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (to Actinobacillus pleuropneumoniae; nonrecombinant subunit vaccine)

IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (transferrin-binding; nonrecombinant subunit vaccine)

IT Nutrition, animal  
 (undernutrition; nonrecombinant subunit vaccine)

IT Detergents  
 (zwitterionic; nonrecombinant subunit vaccine)

IT 7439-89-6, Iron, biological studies  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (chelating agents for; nonrecombinant subunit vaccine)

IT 60-00-4, EDTA, biological studies 67-43-6D, salts 70-51-9, Deferrioxamine 139-13-9 366-18-7, 2,2'-Dipyridyl 12111-24-9  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (nonrecombinant subunit vaccine)

IT 57-09-0, Cetyltrimethylammonium bromide 81-24-3, Taurocholic acid 81-25-4, Cholic acid 83-44-3, Deoxycholic acid 98-11-3D, Benzenesulfonic acid, alkyl derivs., biological studies 151-21-3, SDS, biological studies 302-95-4, Sodium deoxycholate 360-65-6,

Glycodeoxycholic acid 361-09-1, Sodium cholate 475-31-0, Glycocholic acid 516-50-7, Taurodeoxycholic acid 2044-56-6, Lithium dodecyl sulfate 7631-98-3, Sodium N-laurylsarcosinate 9002-92-0, Brij 35 9002-93-1, Triton X-100 9004-95-9, Brij 58 9005-64-5, Tween 20 9005-65-6, Tween 80 9036-19-5, Nonidet P40 9043-30-5, Genapol X-080 14933-09-6, N-Tetradecyl-N,N-dimethyl-3-ammonio-1-propanesulfonate 25339-99-5, Sucrose monolaurate 29836-26-8 59122-55-3 68894-53-1, Tergitol 69227-93-6 75621-03-3, CHAPS 82473-24-3, CHAPSO 85261-20-7, MEGA 10 85316-98-9, MEGA 8 86295-19-4, N-Dodecyl-N,N-dimethyl-3-ammonio-1-propanesulfonate 86303-23-3, Deoxy BigCHAP 106392-12-5, Synperonic PE/F 68 232601-34-2, Tween 48  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (nonrecombinant subunit vaccine)

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Anon; DD 242716 A3 HCAPLUS
- (2) Anon; EP 37931 A2 HCAPLUS
- (3) Anon; US 4845036 HCAPLUS
- (4) Wpids; AU 9061356 A HCAPLUS
- (5) Wpids; AU 9534351 A

L44 ANSWER 10 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1998:800024 HCAPLUS

DN 130:51336

ED Entered STN: 22 Dec 1998

TI Laft mutants of pathogenic gram-negative bacteria

IN Apicella, Michael A.; Gibson, Bradford W.; Nichols, Wade A.

PA University of Iowa Research Foundation, USA; University of California

SO PCT Int. Appl., 31 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K039-02

ICS A01N063-00; C12N001-00; C12N001-20

CC 15-2 (Immunochemistry)

Section cross-reference(s): 3, 10

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9853851	A1	19981203	WO 1998-US10881	19980528 <--
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9877010	A1	19981230	AU 1998-77010	19980528 <--
PRAI US 1997-47791P	P	19970528 <--		
WO 1998-US10881	W	19980528 <--		

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
------------	-------	------------------------------------

WO 9853851	ICM	A61K039-02	
	ICS	A01N063-00; C12N001-00; C12N001-20	
WO 9853851	ECLA	C12N009/10C1	<--

AB A method is provided for identifying, isolating, and producing lipooligosaccharide (LOS) mutants of gram-neg. bacterial pathogens. The method comprises mutating the laft gene of a gram-neg. bacterial pathogen so that there is a lack of a functional Lipid A fatty acid transferase protein. The resulting LOS mutants lack one or more secondary acyl chains as compared to the LOS contained in the wild type gram-neg. bacterial pathogen. The LOS isolated from the laft mutants displays substantially reduced toxicity as compared to that of the wild type strain. Also, the present invention provides methods for using a vaccine formulation containing the laft mutants, the endotoxin isolated therefrom, or the endotoxin isolated therefrom which is then conjugated to a carrier protein, to immunize an individual against infections caused by gram-neg. bacterial pathogens by administering a prophylactically effective amount of the vaccine formulation.

ST lipid A fatty acid transferase gene; lipopolysaccharide endotoxin vaccine gram neg bacteria

IT Immunostimulants

(adjuvants; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic

- gram-neg. bacteria)
- IT Microorganism
  - (antigen; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)
- IT Infection
  - (bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)
- IT Drug delivery systems
  - (carriers; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)
- IT Toxins
  - RL: PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
  - (endotoxins; gram-neg. bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections)
- IT Bordetella pertussis
- Escherichia coli
- Gram-negative bacteria
- Haemophilus ducreyi
- Haemophilus influenzae
- Human adenovirus
- Human parainfluenza virus
- Influenza virus
- Legionella pneumophila
- Moraxella catarrhalis
- Mycoplasma pneumoniae
- Neisseria gonorrhoeae
- Pneumocystis carinii
- Pseudomonas aeruginosa
- Respiratory syncytial virus
- Streptococcus group A
- Streptococcus pneumoniae
- Vaccines
  - (gram-neg. bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections)
- IT Antigens
  - RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
  - (gram-neg. bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections)
- IT Lipopolysaccharides
  - RL: PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
  - (gram-neg. bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections)
- IT Drug delivery systems
  - (injections, i.m.; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)
- IT Drug delivery systems
  - (injections, i.v.; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)
- IT Drug delivery systems
  - (injections, s.c.; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)
- IT Drug delivery systems
  - (intradermal; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)
- IT Gene, microbial
  - RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
  - (laft (lipid A fatty acid transferase); gram-neg. bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections)
- IT Gene, microbial
  - RL: BSU (Biological study, unclassified); BIOL (Biological study)
  - (msbB; gram-neg. bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections)
- IT Drug delivery systems
  - (mucosal; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)
- IT Drug delivery systems

(nasal, intra-; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)

IT Gene  
RL: BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(open reading frame; gram-neg. bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections)

IT Drug delivery systems  
(ophthalmic; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)

IT Drug delivery systems  
(oral; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)

IT Drug delivery systems  
(solns., i.p.; bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections by pathogenic gram-neg. bacteria)

IT 115926-32-4  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(gram-neg. bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections)

IT 217181-27-6  
RL: PRP (Properties)  
(nucleotide sequence; gram-neg. bacterium having mutated lipid A fatty acid transferase gene as vaccine for preventing infections)

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE  
(1) Clementz; J Biol Chem 1997, V272(16), P10353 HCAPLUS  
(2) Jones; Infect Immun 1997, V65(11), P4778 HCAPLUS  
(3) Lee; J Biol Chem 1995, V270(45), P27151 HCAPLUS  
(4) Somerville; J Clin Invest 1996, V97(2), P359 HCAPLUS  
(5) Sprouse; US 5641492 A 1997 HCAPLUS  
(6) Sunshine; J Bacteriol 1997, V179(17), P5521 HCAPLUS

L44 ANSWER 11 OF 18 'HCAPLUS COPYRIGHT 2004 ACS on STN  
AN 1993:17456 HCAPLUS

DN 118:17456

ED Entered STN: 24 Jan 1993

TI Use of the purA gene as a selectable marker in stabilization and integration of plasmid or bacteriophage cloning vectors

IN Brey, Robert Newton, III; Fulginiti, James Peter; Anilionis, Algis

PA American Cyanamid Co., USA

SO Eur. Pat. Appl., 29 pp.

CODEN: EPXXDW

DT Patent

LA English

IC ICM C12N015-74

ICS A61K039-112

ICI C12N015-74, C12R001-42

CC 3-2 (Biochemical Genetics)

Section cross-reference(s): 10, 15

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 512260	A2	19921111	EP 1992-105887	19920406 <--
	EP 512260	A3	19930728		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, PT, SE				
	AT 202800	E	20010715	AT 1992-105887	19920406 <--
	ES 2160573	T3	20011116	ES 1992-105887	19920406 <--
	PT 512260	T	20011228	PT 1992-105887	19920406 <--
	JP 05192161	A2	19930803	JP 1992-134375	19920428 <--
	JP 3320095	B2	20020903		
	NO 9201729	A	19921104	NO 1992-1729	19920430 <--
	CA 2067862	AA	19921104	CA 1992-2067862	19920501 <--
	CA 2067862	C	20031230		
	AU 9215959	A1	19921105	AU 1992-15959	19920501 <--
	AU 654347	B2	19941103		
	US 5919663	A	19990706	US 1995-380297	19950130 <--
	US 5961983	A	19991005	US 1995-448907	19950524 <--
	GR 3036487	T3	20011130	GR 2001-401346	20010831 <--
PRAI	US 1991-695706	A	19910503	<--	
	US 1994-204903	B1	19940302	<--	
	US 1995-380297	A3	19950130	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
EP 512260	ICM	C12N015-74
	ICS	A61K039-112
	ICI	C12N015-74, C12R001-42
US 5961983	ECLA	A61K039/015
AB	Host bacteria carrying deletions in the purA gene (for adenylosuccinate synthetase) are used as hosts for cloning vectors carrying the purA gene as a selectable marker. The vector is stabilized by selection and the purA gene also acts as a site for integration of the plasmid. The use of these vectors does not involve the use of antibiotic resistance markers and is therefore particularly suitable for hosts used in live vaccines. A pUC8-based plasmid carrying the Escherichia coli purA gene and the gene for the nontoxic subunit of the E. coli heat-labile enterotoxin was constructed and introduced into Salmonella dublin, S. typhimurium or Salmonella vaccine strains carrying deletions in the purA gene and transformants selected on minimal medium. This plasmid was maintained in cultures grown on a minimal medium without loss for 80 generations but lost rapidly in the absence of selection (1% retention in 40 generations). When the purA gene was used in integrating vectors the prototrophic phenotype was 100% stable for at least 80 generations in the presence or absence of selection.	
ST	purA gene selectable marker cloning vector; Salmonella adenine auxotrophy selectable marker	
IT	Bordetella pertussis Chlamydia trachomatis Clostridium tetani Corynebacterium diphtheriae Escherichia coli Haemophilus influenzae Klebsiella pneumoniae Moraxella catarrhalis Neisseria gonorrhoeae Neisseria meningitidis Parasite Pseudomonas aeruginosa Staphylococcus aureus Streptococcus pneumoniae Streptococcus pyogenes Vibrio cholerae (antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)	
IT	Aromatic hydrocarbons, biological studies RL: BPN (Biosynthetic preparation); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation) (biosynthesis of, microorganism deficient in as host for expression vectors, complementing genes on plasmids as selectable markers for)	
IT	Plasmid and Episome Virus, bacterial (cloning vector, purA gene as selectable marker for, adenine auxotrophic host for)	
IT	Antigens RL: BIOL (Biological study) (genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)	
IT	Campylobacter Escherichia Salmonella Salmonella dublin Salmonella typhimurium Shigella Vibrio Yersinia (in live vaccines, heterologous antigen genes in, stabilization or integration of, purA gene as selectable marker in)	
IT	Vaccines (live, bacteria carrying cloned antigen genes for, purA gene as a selectable marker for cloning vectors in)	
IT	Plasmid and Episome (pX3005, gene for heat-labile enterotoxin subunit of Escherichia coli on, integration and expression in Salmonella of, purA gene as selectable marker in)	
IT	Plasmid and Episome (pX3006, gene for circumsporozoite antigen of Plasmodium berghei on, integration and expression in Salmonella of, purA gene as selectable marker in)	

- IT Plasmid and Episome  
(pX3007, chimeric gene for fusion protein of circumsporozoite antigen of Plasmodium berghei and Escherichia coli enterotoxin on, integration and expression in Salmonella of, purA gene as selectable marker in)
- IT Plasmid and Episome  
(pX3009, gene for circumsporozoite antigen of Plasmodium berghei on, integration and expression in Salmonella of, purA gene as selectable marker in)
- IT Plasmid and Episome  
(pX3010, gene for heat-labile enterotoxin subunit of Escherichia coli on, integration and expression in Salmonella of, purA gene as selectable marker in)
- IT **Antigens**  
RL: BIOL (Biological study)  
(CS (circumsporozoite), gene for, expression in enterobacteria of, in live vaccines, purA gene for stabilization or integration of antigen genes in, fusion proteins with heat-labile enterotoxin subunit in relation to)
- IT Virus, animal  
(Epstein-Barr, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(adeno-, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(corona-, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(cytomegalo-, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT **Toxins**  
RL: BIOL (Biological study)  
(entero-, LT, gene for, expression in enterobacteria of, in live vaccines, purA gene for stabilization or integration of enterotoxin genes in, fusion proteins with circumsporozoite antigens in relation to)
- IT Virus, animal  
(hepatitis A, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(hepatitis B, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(hepatitis C, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(hepatitis, non-A, non-B, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(herpes simplex 1, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(herpes simplex 2, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(human T-cell leukemia type I, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(human T-cell leukemia type II, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(human immunodeficiency 1, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(human immunodeficiency 2, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(influenza, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(measles, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)
- IT Virus, animal  
(papilloma, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)



IT Virus, animal  
(parainfluenza, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)

IT Fungi  
(pathogenic, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)

IT Virus, animal  
(polio-, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)

IT Virus, animal  
(respiratory syncytial, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)

IT Virus, animal  
(rota-, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)

IT Virus, animal  
(rubella, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)

IT Virus, animal  
(varicella-zoster, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)

IT Virus, animal  
(yellow fever, antigens of, genes for, vaccine bacteria carrying, cloning vectors using purA gene as selectable marker in)

IT Gene, microbial  
RL: BIOL (Biological study)  
(purA, as selectable marker for cloning vectors, deletion of host copy of gene in)

IT 73-24-5, Adenine, biological studies  
RL: BIOL (Biological study)  
(auxotrophy for, as selectable marker for cloning vectors, deletion of host copy of gene in)

IT 120-73-0, Purine  
RL: BIOL (Biological study)  
(biosynthesis of, microorganism deficient in as host for expression vectors, complementing genes on plasmids as selectable markers for)

IT 9023-57-8, Adenylosuccinate synthetase  
RL: BIOL (Biological study)  
(gene for, as selectable marker for cloning vectors, deletion of host copy of gene in)

L44 ANSWER 12 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1991:499235 HCAPLUS

DN 115:99235

ED Entered STN: 06 Sep 1991

TI A method for isolating and purifying transferrin and lactoferrin receptor proteins from bacteria and the preparation of vaccines containing the same

IN Schryvers, Anthony Bernard

PA University Technologies International Inc., Can.

SO PCT Int. Appl., 34 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K039-095

ICS A61K039-102; A61K039-02

CC 63-3 (Pharmaceuticals)

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9012591	A1	19901101	WO 1990-CA131	19900426 <--
	W: AU, CA, JP, KR, SU				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, IT, LU, NL, SE				
	US 5292869	A	19940308	US 1990-507481	19900409 <--
	AU 9055261	A1	19901116	AU 1990-55261	19900426 <--
	AU 649950	B2	19940609		
	JP 04506794	T2	19921126	JP 1990-506296	19900426 <--
	JP 3335622	B2	20021021		
	EP 528787	A1	19930303	EP 1990-906093	19900426 <--
	EP 528787	B1	19981202		
	R: AT, BE, CH, DE, DK, ES, FR, GB, IT, LI, LU, NL, SE				
	ES 2127184	T3	19990416	ES 1990-906093	19900426 <--
	CA 2051808	C	19991214	CA 1990-2051808	19900426 <--
	JP 2002316942	A2	20021031	JP 2002-54731	19900426 <--
	US 6060058	A	20000509	US 1995-483881	19950607 <--
PRAI	US 1989-344356	A	19890427	<--	
	US 1990-507481	A	19900409	<--	

JP 1990-506296	A3	19900426	<--
WO 1990-CA131	A	19900426	<--
US 1991-639365	A3	19910110	<--
US 1992-851005	B1	19920312	<--
US 1994-207719	B1	19940309	<--

## CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
------------	-------	------------------------------------

WO 9012591	ICM	A61K039-095
	ICS	A61K039-102; A61K039-02

AB A method of isolating and purifying transferrin and lactoferrin receptor proteins from bacterial pathogens by affinity chromatog. is described. The proteins are used for preparing vaccine antigens. The vaccine antigens are effective in preventing diseases caused by bacterial pathogens containing lactoferrin and transferrin receptor proteins. The human-lactoferrin binding protein from *Neisseria meningitidis* is identified and characterized. It is incorporated into vaccine preps.

ST vaccine bacteria lactoferrin receptor; transferrin receptor vaccine bacteria

IT **Vaccines**  
(against bacteria, lactoferrin and transferrin receptors as)

IT **Receptors**  
RL: BIOL (Biological study)  
(for lactoferrin and transferrin, from bacteria, vaccines containing)

IT Actinobacillus suis  
Haemophilus avium  
Haemophilus gallinarum  
Haemophilus influenzae  
Haemophilus paragallinarum  
Haemophilus pleuropneumoniae  
Haemophilus somnus  
Haemophilus suis  
Moraxella catarrhalis  
Neisseria gonorrhoeae  
Neisseria lactamicus  
Neisseria meningitidis  
Pasteurella haemolytica  
Pasteurella multocida  
(lactoferrin and transferrin receptors from, vaccines containing)

IT Lactoferrins  
Transferrins  
RL: BIOL (Biological study)  
(receptors for, from bacteria, vaccines containing)

IT **Antigens**  
RL: BIOL (Biological study)  
(vaccines, against bacteria, lactoferrin and transferrin receptors as)

IT Proteins, specific or class  
RL: BIOL (Biological study)  
(lactoferrin-binding, vaccines containing)

IT Proteins, specific or class  
RL: BIOL (Biological study)  
(transferrin-binding, vaccines containing)

L44 ANSWER 13 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1991:171296 HCAPLUS

DN 114:171296

ED Entered STN: 03 May 1991

TI Cytokine and hormone carriers for conjugate vaccines

IN Pillai, Subramonia; Eby, Ronald

PA Praxis Biologics, Inc., USA

SO PCT Int. Appl., 50 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K047-48

ICS A61K039-385

CC 63-3 (Pharmaceuticals)

Section cross-reference(s): 2, 15

FAN.CNT 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9101146	A1	19910207	WO 1990-US3983	19900716 <--
W: AU, CA, FI, JP, KR, NO				
RW: AT, BE, CH, DE, DK, ES, FR, GB, IT, LU, NL, SE				
CA 2063271	AA	19910115	CA 1990-2063271	19900716 <--

AU 9060550	A1	19910222	AU 1990-60550	19900716 <--
AU 651949	B2	19940811		
EP 482068	A1	19920429	EP 1990-911070	19900716 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, IT, LI, LU, NL, SE				
JP 04506662	T2	19921119	JP 1990-510469	19900716 <--
NO 9200160	A	19920305	NO 1992-160	19920113 <--
PRAI US 1989-380566	A	19890714	<--	
WO 1990-US3983	A	19900716	<--	

## CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
------------	-------	------------------------------------

WO 9101146	ICM	A61K047-48
	ICS	A61K039-385

AB Immunogenic conjugates are disclosed comprising a carbohydrate-containing antigen or other antigen bound to or genetically fused with a cytokine, lymphokine, hormone, or growth factor having immunomodulating activity, wherein the cytokine, lymphokine, hormone, or growth factor is capable of modifying immunogenicity of the carbohydrate-containing antigen. The cytokine or lymphokine can be an interleukin or an interferon. The immunogenic conjugate can be used in vaccine and covaccine formulations.

ST vaccine cytokine hormone carrier

IT Animal growth regulators

Hormones

Interferons

RL: PREP (Preparation)

(antigen bound to, for vaccine preparation)

IT Lymphokines and Cytokines

RL: PREP (Preparation)

(antigen conjugates, for vaccine preparation)

IT Carbohydrates and Sugars, biological studies

Oligosaccharides

Polysaccharides, biological studies

RL: PREP (Preparation)

(antigens containing, conjugates with cytokines and hormones, for vaccine preparation)

IT Fungi

Parasite

Virus, animal

(antigens of, conjugates with cytokines and hormones, for vaccine preparation)

IT Bordetella pertussis

Clostridium tetani

Corynebacterium diphtheriae

Escherichia coli

Haemophilus influenzae

Klebsiella pneumoniae

Moraxella catarrhalis

Neisseria gonorrhoeae

Neisseria meningitidis

Pseudomonas aeruginosa

Staphylococcus aureus

Streptococcus pneumoniae

Streptococcus pyogenes

Vibrio cholerae

(capsular polymers of, conjugates with cytokines and hormones, for vaccine preparation)

IT Antigen

RL: BIOL (Biological study)

(carbohydrate and hormone conjugates, in vaccine formulations)

IT Vaccines

(immunogenic conjugates with cytokines and hormones in formulations of)

IT Lipopolysaccharides

RL: PREP (Preparation)

(of bacteria, conjugates with cytokines and hormones, in vaccine preparation)

IT Peptidoglycans

RL: PREP (Preparation)

(of bacterial cell wall, conjugates with cytokines and hormones, in vaccine preparation)

IT Lymphokines and Cytokines

RL: PREP (Preparation)

(interleukin 1.alpha., antigen bound to, for vaccine preparation)

IT Lymphokines and Cytokines

RL: PREP (Preparation)

(interleukin 1.beta., antigen bound to, for vaccine preparation)

IT Lymphokines and Cytokines

RL: PREP (Preparation)  
(interleukin 2, antigen bound to, for vaccine preparation)

IT Lymphokines and Cytokines  
RL: PREP (Preparation)  
(tumor necrosis factor, antigen conjugates, for vaccine preparation)

IT 9002-62-4D, Prolactin, antigen conjugates 9002-72-6D, Somatotropin,  
antigen conjugates 9004-10-8D, Insulin, antigen conjugates  
62229-50-9D, Epidermal growth factor, antigen conjugates 62683-29-8D,  
Granulocyte colony-stimulating factor, antigen conjugates 82197-76-0D,  
Polyribosylribitolphosphate, antigen conjugates 83869-56-1D, Granulocyte  
macrophage colony stimulating factor, antigen conjugates  
RL: BIOL (Biological study)  
(vaccines from)

L44 ANSWER 14 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
AN 1990:558671 HCAPLUS  
DN 113:158671  
ED Entered STN: 27 Oct 1990  
TI T-cell epitope as carrier molecule for conjugate vaccines  
IN Bixler, Garvin; Pillai, Subramonia; Insel, Richard  
PA Praxis Biologics, Inc., USA  
SO PCT Int. Appl., 103 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
IC ICM A61K039-385  
ICS C07K015-04; A61K039-155  
CC 63-3 (Pharmaceuticals)  
Section cross-reference(s): 15  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 8906974	A2	19890810	WO 1989-US388	19890131 <--
	WO 8906974	A3	19890824		
	W: AU, DK, FI, JP, NO				
	RW: AT, BE, CH, DE, FR, GB, IT, LU, NL, SE				
	AU 8930654	A1	19890825	AU 1989-30654	19890131 <--
	AU 634153	B2	19930218		
	EP 399001	A1	19901128	EP 1989-908669	19890131 <--
	EP 399001	B1	19940727		
	R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE				
	JP 03502691	T2	19910620	JP 1989-502396	19890131 <--
	JP 2921574	B2	19990719		
	NO 9002909	A	19900827	NO 1990-2909	19900629 <--
	NO 179164	B	19960513		
	NO 179164	C	19960821		
	DK 9001829	A	19900731	DK 1990-1829	19900731 <--
	DK 174416	B1	20030217		
	US 5785973	A	19980728	US 1995-481923	19950607 <--
PRAI	US 1988-150688	A	19880201	<--	
	US 1989-304783	B1	19890131	<--	
	WO 1989-US388	A	19890131	<--	
	US 1992-828711	B1	19920131	<--	
	US 1993-164989	B1	19931209	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 8906974	ICM	A61K039-385
	ICS	C07K015-04; A61K039-155
US 5785973	ECLA	A61K039/385; A61K047/48R; C07K014/195

AB Conjugates between T-cell epitopes (recognition sites) for bacterial products such as tetanus toxin and medically useful substances such as antigens, haptens, or antigenic determinants are prepared. These conjugates elicit antibody responses and are useful in vaccine preps. The use of the T-cell epitope, as opposed to a larger peptide containing the epitope, provides an economic advantage in that it may be readily prepared as well as a safety advantage in avoidance of use of the whole protein. The conjugates also stimulate antibodies against tumor-specific or tumor-associated antigens and are useful in the immunization of infants whose immune system is not fully developed. The DeLisi and Berzofsky algorithm (1985) for potential amphipathic regions was applied to diphtheria toxin cross-reactive material (CRM) and 6 regions were identified. Peptides corresponding to these CRM regions were synthesized (synthesis given) and those stimulating T-cells were conjugated to phosphoribosylribitol phosphate (PRP, capsular polymer of Haemophilus influenzae b). The conjugates were capable of stimulating antibodies to PRP.

ST T lymphocyte epitope conjugate vaccine; safety T lymphocyte epitope conjugate vaccine

IT Allergy inhibitors  
Neoplasm inhibitors  
Vaccines  
(antigen conjugates with T-cell epitopes of bacterial products as)

IT Microorganism  
Neoplasm, composition  
Parasite  
Virus  
(antigens of, conjugates with T-cell epitopes of bacterial products, as vaccines)

IT *Haemophilus influenzae*  
*Neisseria meningitidis*  
(capsular antigen and outer membrane protein of, conjugates with T-cell epitopes of bacterial products, as vaccines)

IT *Salmonella typhi*  
*Streptococcus pneumoniae*  
(capsular antigens of, conjugates with T-cell epitopes of bacterial products, as vaccines)

IT Allergens  
Antigens  
Carbohydrates and Sugars, biological studies  
RL: BIOL (Biological study)  
(conjugates with T-cell epitopes of bacterial products, as vaccines)

IT *Escherichia coli*  
*Moraxella catarrhalis*  
*Neisseria gonorrhoeae*  
*Streptococcus pyogenes*  
(outer membrane proteins of, conjugates with T-cell epitopes of bacterial products, as vaccines)

IT Lymphocyte  
(B-, epitopes reactive with, conjugates with T-cell epitopes, as vaccines)

IT Glycoproteins, specific or class  
RL: BIOL (Biological study)  
(F (fusion), conjugates, with T-cell epitopes of bacterial products, as vaccine)

IT Proteins, specific or class  
RL: BIOL (Biological study)  
(OMP (outer membrane protein), conjugates, with T-cell epitopes of bacterial products, as vaccines)

IT Lymphocyte  
(T-, bacterial epitope reactive with, conjugates with antigens, as vaccines)

IT Disease  
(autoimmune, treatment of, antigen conjugates with T-cell epitopes of bacterial products for)

IT Lipopolysaccharides  
RL: BIOL (Biological study)  
(conjugates, of gram-neg. bacteria, with T-cell epitopes of bacterial products, as vaccines)

IT Toxins  
Toxoids  
RL: BIOL (Biological study)  
(diphtheria, T-cell epitopes of, antigen conjugates, as vaccines)

IT Bacteria  
(gram-neg., lipopolysaccharides of, conjugates with T-cell epitopes of bacterial products, as vaccines)

IT Toxins  
Toxoids  
RL: BIOL (Biological study)  
(pertussis, T-cell epitopes of, antigen conjugates, as vaccines)

IT Virus, animal  
(respiratory syncytial, vaccine for, F protein conjugates with T-cell epitopes of bacterial products as)

IT Toxins  
Toxoids  
RL: BIOL (Biological study)  
(tetanus, T-cell epitopes of, antigen conjugates, as vaccines)

IT 129774-60-3, Toxin (corynephage .beta. strain ATCC 53281 reduced)  
RL: BIOL (Biological study)  
(T-cell epitopes of, in vaccine preparation)

IT 128516-95-0D, antigen conjugates 128786-78-7D, antigen conjugates  
129813-87-2D, antigen conjugates 129813-88-3D, antigen conjugates  
129836-17-5D, antigen conjugates 129836-18-6D, antigen conjugates

129851-39-4D, antigen conjugates  
 RL: BIOL (Biological study)  
 (as vaccines)

IT 128516-95-0 128786-78-7 129813-87-2 129813-88-3 129836-17-5  
 129836-18-6 129851-39-4  
 RL: BIOL (Biological study)  
 (in vaccine preparation)

IT 129813-89-4P 129813-90-7P 129813-91-8P 129813-92-9P 129813-93-0P  
 129813-94-1P 129813-95-2P 129813-96-3P 129813-97-4P 129813-98-5P  
 129813-99-6P 129814-00-2P 129814-01-3P 129836-19-7P  
 RL: PRP (Properties); PREP (Preparation)  
 (preparation and amino acid sequence of, of diphtheria toxin cross-reactive  
 material, in vaccine preparation)

IT 82197-76-0DP, Polyribosylribitol phosphate, conjugates with synthetic  
 peptides of diphtheria toxin cross-reactive material 129813-89-4DP,  
 conjugates with polyribosylribitol phosphate 129813-90-7DP, conjugates  
 with polyribosylribitol phosphate 129813-91-8DP, conjugates with  
 polyribosylribitol phosphate 129813-92-9DP, conjugates with  
 polyribosylribitol phosphate 129813-93-0DP, conjugates with  
 polyribosylribitol phosphate 129813-94-1DP, conjugates with  
 polyribosylribitol phosphate 129813-95-2DP, conjugates with  
 polyribosylribitol phosphate 129813-96-3DP, conjugates with  
 polyribosylribitol phosphate 129813-97-4DP, conjugates with  
 polyribosylribitol phosphate 129813-98-5DP, conjugates with  
 polyribosylribitol phosphate 129813-99-6DP, conjugates with  
 polyribosylribitol phosphate 129814-00-2DP, conjugates with  
 polyribosylribitol phosphate 129814-01-3DP, conjugates with  
 polyribosylribitol phosphate 129836-19-7DP, conjugates with  
 polyribosylribitol phosphate  
 RL: PREP (Preparation)  
 (preparation of, in preparation of vaccines for Haemophilus influenzae)

L44 ANSWER 15 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 1990:520772 HCAPLUS  
 DN 113:120772  
 ED Entered STN: 29 Sep 1990  
 TI Milk antibody production with shaped polymer microparticles for  
 controlled-release of antigens  
 IN Beck, Lee R.  
 PA Stolle Research and Development Corp., USA  
 SO U.S., 9 pp. Cont. of U.S. Ser. No. 576,001, abandoned.  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 IC ICM A61K039-00  
 NCL 424088000  
 CC 63-3 (Pharmaceuticals)  
 Section cross-reference(s): 15, 17  
 FAN.CNT 7

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4919929	A	19900424	US 1986-910297	19860917 <--
	AU 605159	B2	19910110	AU 1987-76086	19870724 <--
	AU 8776086	A1	19890127		
	US 4956349	A	19900911	US 1988-177223	19880404 <--
	US 5242691	A	19930907	US 1990-580382	19900911 <--
	CA 2072658	AA	19910507	CA 1990-2072658	19901030 <--
	CA 2072658	C	20010220		
	WO 9106321	A1	19910516	WO 1990-US6215	19901030 <--
	W: AU, CA, JP, NO				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE				
	AU 9066149	A1	19910531	AU 1990-66149	19901030 <--
	AU 644820	B2	19931223		
	JP 05501801	T2	19930408	JP 1990-515068	19901030 <--
	JP 2912451	B2	19990628		
	EP 593440	A1	19940427	EP 1990-915892	19901030 <--
	EP 593440	B1	19960821		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
	AT 141513	E	19960915	AT 1990-915892	19901030 <--
	NO 9201748	A	19920701	NO 1992-1748	19920504 <--
	US 5352462	A	19941004	US 1992-966741	19921027 <--
PRAI	US 1984-576001	B1	19840201	<--	
	US 1982-384625	B2	19820603	<--	
	US 1983-576001	B1	19830201	<--	
	US 1983-546162	A3	19831027	<--	
	US 1986-910297	A	19860917	<--	

US 1987-1848	A2	19870109	<--
US 1988-177223	A2	19880404	<--
US 1989-431639	A	19891106	<--
US 1990-580382	A2	19900911	<--

## CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
US 4919929	ICM	A61K039-00
	NCL	424088000
US 5352462	ECLA	A23C009/20; A24B015/30; A61K009/16H6D4; A61K035/20; A61K035/54; A61K035/74; A61K039/395A <--
AB		Milk having elevated IgG antibody levels is produced by (1) i.m. or s.c. implantation within a bovidae of a hyperimmunization response-eliciting amount of an antigenic substance incorporated within shaped microparticles of a biocompatible matrix material which causes controlled-release of the antigen, thereby prolonging antigenic activity within the bovidae; and (2) recovering milk having an elevated level of antibody. The immunized state may be attained more rapidly by simultaneously administering the shaped matrix and the antigenic substance in liquid form. A polyvalent antigen sample (S-100) containing .apprx.26 kinds of heat-killed bacteria was microencapsulated in biodegradable lactide-glycolide copolymer microparticles (diameter < 250 .mu.m). The microparticles were suspended in vehicle (Tween 20 and CMC) and injected i.m. one time into cows. Increased antibody titer was shown in the milk.
ST		controlled release antigen vaccine; cattle milk antibody prodn
IT		<b>Vaccines</b> (antigens in controlled-release microparticles as, for elevated milk antibody production)
IT		Actinobacillus equuli Actinobacillus lignieresii Actinobacillus seminis Bacillus cereus Brucella melitensis Campylobacter fetus Campylobacter fetus intestinalis Cell Chlamydia psittaci Clostridium tetani Corynebacterium pyogenes Corynebacterium renale Enterobacter aerogenes Escherichia coli Fusobacterium necrophorum Gardnerella vaginalis Haemophilus ducreyi Haemophilus influenzae Klebsiella pneumoniae, oxides Leptospira interrogans pomona Listeria monocytogenes Moraxella bovis Mycobacterium tuberculosis Mycoplasma bovigenitalium Mycoplasma hominis Neisseria gonorrhoeae Pasteurella haemolytica Pasteurella multocida Propionibacterium acnes Proteus vulgaris Pseudomonas aeruginosa Pseudomonas maltophilia Rhodococcus equi Salmonella abortusovis Salmonella dublin Salmonella enteritidis Salmonella heidelberg Salmonella paratyphi Salmonella typhimurium Shigella dysenteriae Staphylococcus aureus Staphylococcus epidermidis Streptococcus agalactiae Streptococcus bovis Streptococcus dysgalactiae Streptococcus equisimilis Streptococcus mitis

Streptococcus mutans  
 Streptococcus pneumoniae  
 Streptococcus pyogenes  
 Streptococcus salivarius  
 Streptococcus sanguis  
 Streptococcus uberis  
 Treponema pallidum  
 Virus, animal  
 Yersinia enterocolitica  
 (antigens of, polymer microparticles containing, for controlled-release vaccines and elevated milk antibody production)

IT Milk  
 (elevated antibody production in, controlled-release microencapsulated antigen vaccines for)

IT Disease  
 (lymphopathia venereum, antigens of, polymer microparticles containing, for controlled-release vaccines and elevated milk antibody production)

IT Antibodies  
 RL: PRP (Properties)  
 (milk containing elevated levels of, production of, controlled-release microencapsulated antigen vaccine for)

IT Antigens  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (polymer microparticles containing, for controlled-release vaccines and elevated milk antibody production)

IT Cattle  
 (vaccination of, with controlled-release microencapsulated antigen vaccines)

IT Bovidae  
 (vaccination of, with controlled-release microencapsulated antigen vaccines, for production of milk containing elevated levels of IgG antibodies)

IT Immunoglobulins  
 RL: BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)  
 (G, milk containing elevated levels of, production of, controlled-release microencapsulated antigen vaccine for)

IT Venereal disease  
 (granuloma inguinale, antigens of, polymer microparticles containing, for controlled-release vaccines and elevated milk antibody production)

IT Streptococcus  
 (group B, antigens of, polymer microparticles containing, for controlled-release vaccines and elevated milk antibody production)

IT Polyethers, biological studies  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (ortho esters, antigen microencapsulation in, for controlled-release vaccines and elevated milk antibody production)

IT 24980-41-4 25248-42-4, Poly[oxy(1-oxo-1,6-hexanediyl)] 25266-42-6  
 26009-03-0, Poly[oxy(1-oxo-1,2-ethanediyl)] 26023-30-3,  
 Poly[oxy(1-methyl-2-oxo-1,2-ethanediyl)] 26100-51-6 26124-68-5  
 31621-87-1 34346-01-5  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (antigen microencapsulation in, for controlled-release vaccines and elevated milk antibody production)

L44 ANSWER 16 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 1990:42564 HCAPLUS  
 Correction of: 1988:461449  
 DN 112:42564  
 Correction of: 109:61449  
 ED Entered STN: 04 Feb 1990  
 TI Complexes of vitamin B12 and biologically active agents for oral drug delivery  
 IN Russell-Jones, Gregory John; De Aizpurua, Henry James; Howe, Peter Allan; Burge, Geoffery Lewis  
 PA Biotechnology Australia Pty. Ltd., Australia  
 SO Eur. Pat. Appl., 19 pp.  
 CODEN: EPXXDW  
 DT Patent  
 LA English  
 IC ICM A61K047-00  
 ICS A61K037-02; A61K037-24  
 CC 63-5 (Pharmaceuticals)  
 Section cross-reference(s): 1



## FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 220030	A2	19870429	EP 1986-307849	19861010 <--
	EP 220030	A3	19870930		
	EP 220030	B1	19910619		
	R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE				
	ZA 8607703	A	19870624	ZA 1986-7703	19861009 <--
	IN 165029	A	19890805	IN 1986-CA738	19861009 <--
	IL 80264	A1	19911121	IL 1986-80264	19861009 <--
	CA 1330791	A1	19940719	CA 1986-520162	19861009 <--
	WO 8702251	A1	19870423	WO 1986-AU299	19861010 <--
	W: AU, DK, FI, JP, KR, NO, SU, US				
	AU 8665289	A1	19870505	AU 1986-65289	19861010 <--
	AU 587658	B2	19890824		
	CN 86107590	A	19870520	CN 1986-107590	19861010 <--
	CN 1045391	B	19991006		
	JP 63501015	T2	19880414	JP 1986-505525	19861010 <--
	JP 08000779	B4	19960110		
	AT 64534	E	19910715	AT 1986-307849	19861010 <--
	ES 2051690	T3	19940701	ES 1986-307849	19861010 <--
	DK 8702925	A	19870609	DK 1987-2925	19870609 <--
	DK 167099	B1	19930830		
	US 5428023	A	19950627	US 1993-61343	19930517 <--
	US 5589463	A	19961231	US 1995-479635	19950607 <--
	US 5807832	A	19980915	US 1995-483811	19950607 <--
PRAI	AU 1985-2838	A	19851010	<--	
	EP 1986-307849	A	19861010	<--	
	WO 1986-AU299	A	19861010	<--	
	US 1987-84821	B1	19870609	<--	
	AU 1988-2838	A	19881010	<--	
	US 1990-600137	B1	19901019	<--	
	US 1991-759697	B1	19910909	<--	
	US 1993-61343	A3	19930517	<--	

## CLASS

	PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
	EP 220030	ICM	A61K047-00
		ICS	A61K037-02; A61K037-24
	US 5589463	ECLA	A61K039/00; A61K039/35; A61K039/385; A61K047/48H4 <--
	US 5807832	ECLA	A61K047/48H4 <--
AB	Vitamin B12 is covalently bonded to biol. active substances such as hormones, proteins, antigens, haptens, and antibiotics. The B12 in the complex can still react with intrinsic factor in the intestine, so the natural uptake mechanism for B12 is utilized to deliver various otherwise nonabsorbable compds. to the circulation. Vitamin B12 was hydrolyzed to give the monocarboxylic acid, which was coupled to N-hydroxysuccinamide, and treated with Lys-6-leutiniz hormone releasing hormone (I) to give B12-I. This complex induced ovulation in mice following oral administration, whereas LHRH, orally, did not induce ovulation. A B12-neomycin complex was as effective orally in mice against Salmonella typhimurium as i.m. neomycin or i.m. B12-neomycin, whereas neomycin orally was ineffective.		
ST	oral absorption drug vitamin B12 complex; intestine absorption drug vitamin B12 complex; vaccine vitamin B12 complex oral absorption; hormone vitamin B12 complex oral absorption; antibiotic vitamin B12 complex oral absorption		
IT	Blood-brain barrier		
	Intestine, metabolism		
	(absorption by, of vitamin B12-biol. active agent complexes)		
IT	Pollen		
	(antigen of, complexes with vitamin B12, for oral administration to stimulate immune response)		
IT	Kapot		
	(antigens of, complexes with vitamin B12, for oral administration to stimulate immune response)		
IT	Wheat		
	(chaff, antigens of, complexes with vitamin B12, for oral administration to stimulate immune response)		
IT	Antibiotics		
	(complexes with vitamin B12, for oral administration)		
IT	Allergens		
	Haptens		
	Hormones		
	RL: BIOL (Biological study)		
	(complexes with vitamin B12, for oral delivery)		

IT Albumins, compounds  
Interferons  
RL: BIOL (Biological study)  
(complexes with vitamin B12, for oral drug delivery)

IT Canidae  
Dermatophagoides pteronyssinus  
Felidae  
Swine  
(hair of, antigen of, complexes with vitamin B12, for oral administration to stimulate immune response)

IT Placenta  
(passage of substances across, vitamin B12-active agent complexes for)

IT Cholera  
Coccidiosis  
Diphtheria  
Haemophilus influenzae  
Influenza  
Klebsiella pneumoniae  
Measles  
Moraxella catarrhalis  
Mycobacterium BCG  
Plague  
Rubella  
Salmonella typhi  
Streptococcus  
Streptococcus pneumoniae  
Tetanus  
Tuberculosis  
Variola  
Yellow fever  
(protein derived from or immunogens against, complexes with vitamin B12 for oral vaccination)

IT Ovulation  
(stimulation of, by orally administered vitamin B12-LHRH complex)

IT Allergy inhibitors  
(vitamin B12 complexes with hapten or antigen for)

IT Immunomodulators  
(vitamin B12-biol. active agent complexes as)

IT Antigens  
RL: BIOL (Biological study)  
(B12, complexes with vitamin, for oral delivery)

IT Proteins, specific or class  
RL: BIOL (Biological study)  
(complexes, with vitamin B12, for oral delivery)

IT Polysaccharides, compounds  
RL: BIOL (Biological study)  
(complexes, with vitamin B12, for oral drug delivery)

IT Embryo  
(fetus, drug delivery in, vitamin B12-active agent complexes for)

IT Vaccines  
(oral, vitamin B12 complexes with antigens as)

IT 6539-14-6 37434-06-3 38285-78-8 57683-72-4 57757-57-0 62069-75-4  
68181-17-9 74676-97-4 74676-98-5 81069-02-5 98897-08-6  
102568-45-6 111105-75-0 111105-76-1  
RL: BIOL (Biological study)  
(coupling agent, for preparation of vitamin B12-biol. active agent complexes)

IT 6066-82-6P  
RL: PREP (Preparation)  
(crosslinking agent for preparation of vitamin B12-biol. active agent complexes)

IT 68-19-9DP, Vitamin B12, complexes with bovine serum albumin  
RL: PREP (Preparation)  
(preparation and stimulation of immune response by, in oral administration)

IT 88326-63-0DP, Zincobinamide, complexes with biol. active agents  
111070-88-3DP, complexes with biol. active agents  
RL: PREP (Preparation)  
(preparation of, for oral drug administration)

IT 51-17-2DP, 1H-Benzimidazole, derivs., complexes with biol. active agents  
57-42-1DP, vitamin B12 complexes 58-14-0DP, vitamin B12 complexes  
58-22-0DP, vitamin B12 complexes 59-47-2DP, vitamin B12 complexes  
61-32-5DP, vitamin B12 complexes 61-33-6DP, vitamin B12 complexes  
68-19-9DP, Vitamin B12, complexes with biol. active agents 1404-04-2DP,  
Neomycin, vitamin B12 complexes 1867-66-9DP, vitamin B12 complexes  
4697-36-3DP, vitamin B12 complexes 7361-61-7DP, vitamin B12 complexes  
9002-61-3DP, vitamin B12 complexes 9002-70-4DP, vitamin B12 complexes

9004-10-8DP, Insulin, vitamin B12 complexes 9004-66-4DP, Iron dextran, vitamin B12 complexes 9034-40-6DP, Luteinizing hormone-releasing factor, vitamin B12 complexes 13408-75-8DP, complexes with biol. active agents 13422-51-0DP, complexes with biol. active agents 13422-52-1DP, complexes with biol. active agents 13422-55-4DP, complexes with biol. active agents 13870-90-1DP, complexes with biol. active agents 14978-39-3DP, complexes with biol. active agents 15041-07-3DP, complexes with biol. active agents 15671-27-9DP, complexes with biol. active agents 18559-94-9DP, vitamin B12 complexes 20623-13-6DP, complexes with biol. active agents 23208-66-4DP, complexes with biol. active agents 23388-02-5DP, complexes with biol. active agents 52671-12-2DP, vitamin B12 complexes 57285-09-3DP, Inhibin, vitamin B12 complexes 112076-75-2DP, complexes with biol. active agents  
 RL: PREP (Preparation)  
 (preparation of, for oral drug delivery)

L44 ANSWER 17 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1981:145332 HCAPLUS

DN 94:145332

ED Entered STN: 12 May 1984

TI Antigens and vaccines containing them

IN Hours, Michel; Pourquier, Andre

PA Fr.

SO Fr. Demande, 11 pp.

CODEN: FRXXBL

DT Patent

LA French

IC A61K039-02; C12K005-00

CC 63-3 (Pharmaceuticals)

Section cross-reference(s): 15

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 2446111	A1	19800808	FR 1979-1301	19790112 <--
	FR 2446111	B1	19820702		
PRAI	FR 1979-1301	A	19790112	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
------------	-------	------------------------------------

FR 2446111	IC	A61K039-02IC C12K005-00
------------	----	-------------------------

AB Antigenic complexes were isolated from organisms (*Streptococcus pyogenes*, *S. aureus*, *Diplococcus pneumoniae*, *Neisseria catarrhalis*, *N. elongata*, *Escherichia coli*, *Klebsiella pneumoniae*, *Hemophilus influenzae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*) by an improved process in which the cell membrane was isolated and then solubilized by the simultaneous action of Na deoxycholate [302-95-4] (0.5-5 mg/mg proteins), lysozyme [9001-63-2] (1 mg/mL), and EDTA Na [64-02-8] (25 mM) at pH .apprx.7 for 12-24 h at 20.degree.. The extract was centrifuged and the supernatant liquid could be used directly as a source of antigens without removal of the solubilizers. The solubilizers protected the antigenic complexes against contaminant proteases. The preparation of vaccines from these antigens is discussed.

ST antigen bacteria vaccine

IT *Branhamella catarrhalis*

*Escherichia coli*

*Haemophilus influenzae*

*Klebsiella pneumoniae*

*Neisseria elongata*

*Proteus mirabilis*

*Pseudomonas aeruginosa*

*Staphylococcus aureus*

*Streptococcus pneumoniae*

*Streptococcus pyogenes*

(antigen separation from cell membranes of, for vaccine manufacture)

IT Vaccines

(manufacture of, antigen separation for)

IT Antigens

RL: PROC (Process)

(of bacteria cell membranes, separation of, for vaccines)

IT 302-95-4

RL: BIOL (Biological study)

(bacteria cell membrane solubilization by EDTA and lysozyme and, in antigen preparation)

IT 9001-63-2

RL: BIOL (Biological study)

(bacteria cell membrane solubilization by deoxycholate and EDTA and, in

antigen preparation)  
 IT 64-02-8  
 RL: BIOL (Biological study)  
 (bacteria cell membrane solubilization by deoxycholate and lysozyme  
 and, in antigen preparation)

L44 ANSWER 18 OF 18 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 1973:470125 HCAPLUS  
 DN 79:70125  
 ED Entered STN: 12 May 1984  
 TI Autogenous vaccine. Preparation technique and efficiency factors  
 AU Cury, Rolando  
 CS Fac. Med. Vet. Zootec., Univ. Sao Paulo, Sao Paulo, Brazil  
 SO Revista de Saude Publica (1972), 6(4), 371-83  
 CODEN: RSPUB9; ISSN: 0034-8910  
 DT Journal  
 LA Portuguese  
 CC 63-3 (Pharmaceuticals)  
 AB Seeding procedures were used to obtain antigenous vaccines. Bacteria were  
 preselected. Media were tested for sterility before seeding by heating  
 them at 37.degree. for 24 hr. Simple media (pH 7.4), agar (pH 7.4),  
 semisolid agar (pH 6.8-7.0), thioglycolate-dextrose, and agar-triptose  
 were used. Cultures of Streptococcus, Staphylococcus, Enterobacteriaceae,  
 Haemophilus, Bordetella, Pasteurella, Moraxella, Clostridium, Pseudomonas,  
 Neisseria, and Brucella were prepared Adequate culture conditions are given  
 in each case. I2, HCHO, and the heat were used to inactivate the bacteria  
 to prevent damage of existing antigenous agents. Instructions to apply  
 vaccines prepared are given.

ST bacteria vaccine  
 IT Bacteria  
 Bordetella  
 Brucella  
 Clostridium  
 Enterobacteriaceae  
 Haemophilus  
 Moraxella  
 Neisseria  
 Pasteurella  
 Pseudomonas  
 Staphylococcus  
 Streptococcus  
 (antigens of, vaccines of)

IT Vaccines  
 (of bacteria antigens)

IT Antigens  
 RL: BIOL (Biological study)  
 (of bacteria, vaccines of)

=> d all 136 tot

L36 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 2000:741941 HCAPLUS  
 DN 133:320987  
 ED Entered STN: 20 Oct 2000  
 TI Conserved adhesin motif and methods of use thereof  
 IN Lupas, Andrei Nicolae  
 PA Smithkline Beecham Corporation, USA; Smithkline Beecham PLC  
 SO PCT Int. Appl., 85 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM A61K038-00  
 ICS A61K039-00; A61K039-395; C07H021-04; C07K001-00; C07K016-00;  
 C12N005-00; C12N007-00; C12N015-09; C12P021-08; G01N033-53  
 CC 15-2 (Immunochemistry)  
 Section cross-reference(s): 1, 3, 63  
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000061165	A1	20001019	WO 2000-US9866	20000413 <--
W: JP, US				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL,				
PT, SE				
PRAI US 1999-129073P	P	19990413 <--		

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 2000061165	ICM ICS	A61K038-00 A61K039-00; A61K039-395; C07H021-04; C07K001-00; C07K016-00; C12N005-00; C12N007-00; C12N015-09; C12P021-08; G01N033-53
AB		Isolated polypeptides which are conserved in eubacterial extracellular domains are identified in five pathogens of the beta and gamma branches of proteobacteria. These polypeptides, alone or as fusion proteins with a second protein, are useful in the generation of antibodies or other antagonists. The peptides, fusion proteins, and antibodies are useful as vaccine components or therapeutic agents against bacterial infection or as diagnostic reagents. These polypeptides are also useful in screening methods for other agonists and antagonists which may be used in diagnosis, therapy, and as vaccines.
ST		proteobacteria infection adhesin motif antibody vaccine
IT		Chaperonins RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (DnaK, fusion protein; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Proteins, specific or class RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (GST, fusion protein; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Heat-shock proteins RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (HSP 70, fusion protein; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Proteins, specific or class RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (NS1 (nonstructural, 1), fusion protein; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Proteins, specific or class RL: BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (UspA1; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Proteins, specific or class RL: BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (UspA2; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Proteins, specific or class RL: BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (YadA outer membrane adhesin; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Receptors RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process) (adhesin; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Immunostimulants (adjuvants; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Antibodies RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (anti-idiotypic; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Infection (bacterial; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
IT		Drug delivery systems

- (carriers; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT **Antigens**  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (composition; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT Actinobacillus  
 Bacteria (Eubacteria)  
 Drug screening  
 Escherichia coli  
**Haemophilus**  
 Influenza  
 Labels  
**Moraxella**  
 Mycobacterium  
 Neisseria  
 Pathogen  
 Protein motifs  
 Protein sequences  
 Proteobacteria  
 Simulation and Modeling, physicochemical  
**Vaccines**  
 Yersinia  
 Yersinia pestis  
 (conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT Fusion proteins (chimeric proteins)  
 RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT **Adhesins**  
 RL: BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT Antibodies  
 Nucleic acids  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT Toxoids  
 RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (diphtheria, fusion protein; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT Immunoglobulins  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (fragments; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT .alpha.-Factor (microbial)  
 RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (fusion protein; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT Diagnosis  
 (immunodiagnosis; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT Animal cell  
 (mammalian; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)
- IT Antibodies  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (monoclonal; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)

infection)

IT Animal virus  
(recombinant; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)

IT Genetic element  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(regulatory sequence; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)

IT Toxoids  
RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(tetanus, fusion protein; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)

IT 302579-96-0 302579-98-2  
RL: PRP (Properties)  
(Unclaimed; conserved adhesin motif and methods of use thereof)

IT 301857-38-5 302323-37-1 302323-38-2 302323-39-3 302323-42-8  
302323-48-4 302323-49-5 302323-50-8 302323-51-9 302323-52-0  
302323-53-1 302323-54-2 302323-55-3 302323-56-4 302323-57-5  
302352-24-5 302798-58-9 302798-59-0  
RL: BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)

IT 60267-61-0P, Ubiquitin  
RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(fusion protein; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)

IT 9030-53-9, Galactokinase 9031-11-2, .beta.-Galactosidase  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(fusion protein; conserved adhesin motif and fusion proteins for use as vaccine or diagnostic agent and therapeutic agent against bacterial infection)

IT 302684-47-5 302684-48-6 302684-49-7 302684-50-0 302684-51-1  
302684-52-2 302684-53-3 302684-54-4 302684-55-5 302684-56-6  
302684-57-7 302684-58-8 302684-59-9 302684-60-2 302684-61-3  
302684-62-4 302684-63-5 302684-64-6 302684-65-7 302684-66-8  
302684-67-9 302684-68-0 302684-69-1 302684-70-4 302684-71-5  
302684-72-6 302684-73-7 302684-74-8 302684-75-9 302684-76-0  
302684-77-1 302684-78-2 302684-79-3 302684-80-6 302684-81-7  
302684-82-8 302684-83-9 302684-84-0 302684-85-1 302684-86-2  
302684-87-3 302684-88-4 302684-89-5 302684-90-8 302905-45-9  
302905-47-1 302905-52-8 302905-61-9 302905-74-4 302906-00-9  
302906-01-0 302906-02-1  
RL: PRP (Properties)  
(unclaimed protein sequence; conserved adhesin motif and methods of use thereof)

IT 302323-40-6 302323-41-7 302323-43-9 302323-44-0 302323-45-1  
302323-46-2 302323-47-3 302323-58-6 302323-59-7 302323-60-0  
302323-61-1 302323-62-2 302323-63-3 302323-64-4 302323-65-5  
302323-66-6 302325-08-2 302325-09-3 302325-10-6 302325-11-7  
302579-61-9 302579-63-1 302579-65-3 302579-67-5 302579-69-7  
302579-71-1 302579-74-4 302579-81-3 302579-86-8 302579-88-0  
302579-91-5 302579-99-3 302580-00-3 302580-01-4 302580-02-5  
302580-04-7 302580-05-8 302580-06-9 302580-07-0 302580-08-1  
302580-09-2 302580-10-5 302580-11-6 302580-12-7 302580-13-8  
302580-16-1 302580-19-4 302580-20-7 302580-22-9 302580-24-1  
302580-25-2 302580-26-3 302580-27-4 302580-28-5 302580-29-6  
302580-30-9 302580-31-0 302580-32-1 302580-33-2 302580-34-3  
302580-35-4 302580-36-5 302580-38-7 302580-39-8 302580-40-1  
302580-41-2 302580-42-3 302684-46-4 302798-57-8  
RL: PRP (Properties)  
(unclaimed sequence; conserved adhesin motif and methods of use thereof)

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Skurnik; Molecular Microbiology 1989, V3, P517 HCAPLUS
- (2) The Board Of Regents The University Of Texas System; WO 9828333 A2 1998 HCAPLUS

Search done by Noble Jarrell

=> d all 142 tot

L42 ANSWER 1 OF 3 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 1999:495189 HCAPLUS  
 DN 131:129044  
 ED Entered STN: 10 Aug 1999  
 TI Vaccine composition comprising milled lyophilizate of antigenic whole cells  
 IN Hafner, Roderick Peter  
 PA Raby Limited, UK  
 SO PCT Int. Appl., 33 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM A61K039-00  
 ICS A61K039-02  
 CC 15-2 (Immunochemistry)  
 Section cross-reference(s): 63  
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9938529	A1	19990805	WO 1999-GB287	19990128 <--
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9922897	A1	19990816	AU 1999-22897	19990128 <--
PRAI GB 1998-1870	A	19980128		<--
WO 1999-GB287	W	19990128		<--

# CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 9938529	ICM	A61K039-00
	ICS	A61K039-02
WO 9938529	ECLA	A61K039/00; A61K039/102; A61K039/02 <--
AB	A vaccine composition for the prevention of bacterial or fungal infections of mucosal surfaces comprises a lyophilizate of antigenic whole cells milled to a particle size of from about 20 to 350<mm. The vaccine may contain killed organisms such as Haemophilus influenzae or Pseudomonas aeruginosa and is useful, for example, for preventing the colonization by these organisms of patients suffering from chronic lung diseases or, in previously colonized patients, for preventing the occurrence of acute infection of the respiratory tract.	
ST	vaccine lyophilized bacteria fungus yeast antigen	
IT	Immunostimulants (adjuvants; vaccine composition comprising milled lyophilizate of antigenic whole cells)	
IT	Infection (bacterial, secondary; vaccine composition comprising milled lyophilizate of antigenic whole cells)	
IT	Drug delivery systems (capsules; vaccine composition comprising milled lyophilizate of antigenic whole cells)	
IT	Eye, disease (conjunctivitis, bacterial; vaccine composition comprising milled lyophilizate of antigenic whole cells)	
IT	Ear (disease, eustachian tube infection; vaccine composition comprising milled lyophilizate of antigenic whole cells)	
IT	Respiratory tract (disease, exacerbation; vaccine composition comprising milled lyophilizate of antigenic whole cells)	
IT	Mammary gland Urogenital tract (disease, infection; vaccine composition comprising milled lyophilizate of antigenic whole cells)	
IT	Nose (diseases, infection; vaccine composition comprising milled lyophilizate of antigenic whole cells)	



IT Escherichia coli  
(enteropathogenic; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Escherichia coli  
(enterotoxigenic; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Digestive tract  
(gastroenteritis; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Digestive tract  
Eye, disease  
Mouth  
Respiratory tract  
Urogenital tract  
Vagina  
(infection; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Ear  
(middle, infection; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Pharynx  
(nasopharynx, infection; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Ear  
(otitis, otitis media; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Pharynx  
(pharyngitis; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Drug delivery systems  
(powders; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Drug delivery systems  
(tablets; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Tonsil  
(tonsillitis; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Digestive tract  
(ulcer; vaccine composition comprising milled lyophilizate of antigenic whole cells)

IT Bacteria (Eubacteria)  
Burkholderia cepacia  
Candida  
Candida albicans  
Candida glabrata  
Candida krusei  
Chlamydia trachomatis  
Cholera  
Common cold  
Corynebacterium parvum  
Diarrhea  
Diphtheria  
Fermentation  
Fungi  
Granulicatella adiacens  
Haemophilus influenzae  
Helicobacter pylori  
Klebsiella pneumoniae  
Klebsiella pneumoniae ozaenae  
Lactococcus lactis  
Meningitis  
Microorganism  
Moraxella catarrhalis  
Mycobacterium BCG  
Mycobacterium tuberculosis  
Mycosis  
Neisseria gonorrhoeae  
Neisseria meningitidis  
Pertussis  
Pneumonia  
Pseudomonas  
Pseudomonas aeruginosa  
Salmonella typhi  
Sexually transmitted diseases  
Staphylococcus aureus

Streptococcus  
 Streptococcus mutans  
 Streptococcus pneumoniae  
 Streptococcus pyogenes  
 Tuberculosis  
 Typhoid fever  
 Vaccines  
 Vibrio cholerae  
 Virus  
 Yeast

(vaccine composition comprising milled lyophilizate of antigenic whole cells)

## IT Antigen

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(vaccine composition comprising milled lyophilizate of antigenic whole cells)

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Angus, R; DEVELOPMENTS IN BIOLOGICAL STANDARDIZATION 1984, V56, P659  
 MEDLINE

(2) Esquisabel, A; JOURNAL OF MICROENCAPSULATION 1997, V14(5), P627 HCAPLUS

L42 ANSWER 2 OF 3 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1999:464181 HCAPLUS

DN 131:86860

ED Entered STN: 29 Jul 1999

TI Lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals

IN Gu, Xin-Xing; Robbins, John B.

PA The Government of the United States of America, Department of Health and Human, USA

SO PCT Int. Appl., 60 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K039-02

ICS A61K039-385; C08B037-00

CC 15-2 (Immunochemistry)

Section cross-reference(s): 63

FAN.CNT 3

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 9936086	A1	19990722	WO 1999-US590	19990112 <--
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2315746	AA	19990722	CA 1999-2315746	19990112 <--
AU 9922212	A1	19990802	AU 1999-22212	19990112 <--
BR 9906902	A	20001017	BR 1999-6902	19990112 <--
EP 1047447	A1	20001102	EP 1999-902170	19990112 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2002509115	T2	20020326	JP 2000-539859	19990112 <--
US 6685949	B1	20040203	US 2000-610034	20000705 <--
US 2004126381	A1	20040701	US 2003-688115	20031017 <--
US 2004115214	A1	20040617	US 2003-729027	20031205 <--
PRAI US 1998-71483P	P	19980113	<--	
US 1996-16020P	P	19960423	<--	
US 1997-842409	A3	19970423	<--	
WO 1999-US590	W	19990112	<--	
US 2000-610034	A2	20000705		
US 2001-789017	A2	20010220		
US 2001-288695P	P	20010503		
WO 2001-US32331	A1	20011016		

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 9936086	ICM	A61K039-02
	ICS	A61K039-385; C08B037-00

Search done by Noble Jarrell

WO 9936086 ECLA A61K039/02; A61K039/385; C12P019/04 <--  
 US 2004126381 ECLA A61K039/02; A61K039/102; A61K039/385; C12P019/04 <--  
 US 2004115214 ECLA A61K039/02; A61K039/385; C12P019/04 <--

AB A conjugate vaccine for *Moraxella catarrhalis* comprising isolated lipooligosaccharide from which esterified fatty acids have been removed, to produce a detoxified lipooligosaccharide (dLOS), or from which lipid A has been removed, to produce a detoxified oligosaccharide (OS), which is linked to an immunogenic carrier. The immunogenic carrier is selected from the group consisting of UspA or CD derived from *M. catarrhalis*, tetanus toxoid, HMP derived from *Haemophilus influenza*, diphtheria toxoid, detoxified *P. aeruginosa* toxin A, cholera toxin, pertussis toxin, hepatitis B surface or core antigen, rotavirus VP 7 protein, CRM, CRM197, CRM3201 and respiratory syncytial virus F and G protein. The vaccine is useful for preventing otitis media and respiratory infections caused by *M. catarrhalis* in mammals, including humans.

ST *Moraxella catarrhalis* lipooligosaccharide vaccine conjugate; fatty acid lipid A removal vaccine

IT Proteins, specific or class  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (CD; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Proteins, specific or class  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (F; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Proteins, specific or class  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (UspA; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Glycoproteins, specific or class  
 Proteins, specific or class  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (VP7; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Alums  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (adjuvant; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Immunostimulants  
 (adjuvants; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Antigens  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (carrier; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Chemistry  
 (chemical compds., linker; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Toxins  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (cholera; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Proteins, specific or class  
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (conjugates; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Glycolipids  
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (detoxified and conjugated; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Toxins  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (diphtheria, CRM and CRM197 and CRM3201; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Toxoids  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (diphtheria; lipooligosaccharide-based vaccine for prevention of *Moraxella* (Branhamella) catarrhalis infections in mammals)

IT Toxins  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (exotoxins; lipooligosaccharide-based vaccine for prevention of

- Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Antigens**  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (hepatitis B core; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Antigens**  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (hepatitis B surface; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Proteins, specific or class**  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (high-mol.-weight; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Carriers**  
 (immunogenic; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Respiratory tract**  
 (infection; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Drug delivery systems**  
 (injections, i.m.; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Drug delivery systems**  
 (injections, s.c.; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Proteins, specific or class**  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (linker; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Clostridium perfringens**  
**Haemophilus influenzae**  
**Moraxella catarrhalis**  
 Pseudomonas aeruginosa  
 Respiratory syncytial virus  
 Rotavirus  
**Vaccines**  
 (lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Fatty acids, processes**  
**Lipid A**  
 RL: REM (Removal or disposal); PROC (Process)  
 (lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **G proteins (guanine nucleotide-binding proteins)**  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Lipid A**  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (monophosphates, adjuvant; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Drug delivery systems**  
 (mucosal; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Drug delivery systems**  
 (nasal, intra-; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Ear**  
 (otitis, otitis media;  
 lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Drug delivery systems**  
 (parenterals; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Toxins**  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (pertussis; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Toxoids**  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (tetanus; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)
- IT **Toxins**  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (toxin A; lipooligosaccharide-based vaccine for prevention of Moraxella

(Branhamella) catarrhalis infections in mammals)

IT Acids, biological studies  
Bases, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(treatment; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)

IT 99-20-7D, Trehalose, dimycolate derivative  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(adjuvant; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)

IT 60-32-2, .epsilon.-Aminohexanoic acid 1071-93-8, Adipic acid dihydrazide 24954-67-4, p-Nitrophenylethyl amine 32449-92-6, D-Glucuronolactone  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(linker; lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)

IT 302-01-2, Hydrazine, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(lipooligosaccharide-based vaccine for prevention of Moraxella (Branhamella) catarrhalis infections in mammals)

RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE  
(1) Edebrink, P; CARBOHYDR RES 1995, V266(2), P237 HCAPLUS  
(2) Gibson, B; WO 9853851 A 1998 HCAPLUS  
(3) Gu, X; INFECTION AND IMMUNITY 1993, V61(5), P1873 HCAPLUS  
(4) Gu, X; INFECTION AND IMMUNITY 1996, V64(10), P4047 HCAPLUS  
(5) Gu, X; INFECTION AND IMMUNITY 1998, V66(5), P1891 HCAPLUS  
(6) Kelly, J; ANALYTICAL BIOCHEMISTRY 1996, V233(1), P15 HCAPLUS  
(7) Murphy, T; MICROBIOLOGICAL REVIEWS 1996, V60(2), P267 HCAPLUS

L42 ANSWER 3 OF 3 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1997:128048 HCAPLUS

DN 126:211022

ED Entered STN: 26 Feb 1997

TI Vaccines for nontypeable Haemophilus influenzae

IN Green, Bruce A.; Zlotnick, Gary W.

PA Praxis Biologics, Inc., USA

SO U.S., 24 pp., Cont.-in-part of U.S. Ser. No. 320, 971, abandoned.

CODEN: USXXAM

DT Patent

LA English

IC ICM A61K039-102

ICS A61K039-385

NCL 424256100

CC 15-2 (Immunochemistry)

FAN.CNT 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5601831	A	19970211	US 1990-491466	19900309 <--
CA 2047681	AA	19900910	CA 1990-2047681	19900309 <--
CA 2047681	C	20000201		
EP 606921	A1	19940720	EP 1994-100492	19900309 <--
EP 606921	B1	20000802		
R: AT, BE, CH, DE, DK, ES, FR, GB, IT, LI, LU, NL, SE				
ES 2063965	T3	19950116	ES 1990-905112	19900309 <--
AT 195076	E	20000815	AT 1994-100492	19900309 <--
US 5780601	A	19980714	US 1995-447653	19950523 <--
US 5955580	A	19990921	US 1995-449406	19950523 <--
US 6420134	B1	20020716	US 1995-448097	19950523 <--
PRAI US 1989-320971	B2	19890309	<--	
EP 1990-905112	A3	19900309	<--	
US 1990-491466	A3	19900309	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
US 5601831	ICM	A61K039-102
	ICS	A61K039-385
	NCL	424256100
US 5601831	ECLA	C07K014/285; C07K016/12A30; C07K019/00 <--
EP 606921	ECLA	C07K014/285 <--
US 5780601	ECLA	C07K014/285; C07K016/12A30; C07K019/00 <--
US 5955580	ECLA	C07K014/285; C07K016/12A30; C07K019/00 <--
US 6420134	ECLA	C07K014/285; C07K016/12A30; C07K019/00 <--
AB		Protein "e" of H. influenzae, a lipoprotein of approx. 28,000 daltons, has been purified and sequenced. Protein "e" and peptides or proteins having

a shared epitope, can be used to vaccinate against non-typable (and typable) *H. influenzae* and to prevent otitis media caused by *H. influenzae*. For this purpose, protein "e" or derivs. thereof can be produced in native, synthetic or recombinant forms and can be administered alone or in conjunction with other antigens of *H. influenzae*. Protein "e" can also be used in multivalent vaccines designed for *H. influenzae* and one or more other infectious organisms. Protein "e" was isolated from *Haemophilus* cell envelopes and characterized, polyclonal anti-protein "e" antiserum and monoclonal anti-protein "e" antibodies were prepared, protein "e" gene was isolated and nucleotide sequence was determined and mol. cloning of the gene was performed, bactericidal activity of vaccine comprising protein "e" subunit was studied, and synergy of anti-protein "e" with other antibodies were demonstrated.

- ST vaccine *Haemophilus influenzae* protein e antibody
- IT Proteins, specific or class  
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (E; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Proteins, specific or class  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (OMP (outer membrane protein); protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Immunostimulants  
 (adjuvants; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Antibodies  
 RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (bactericidal; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Peptides, biological studies  
 Proteins, general, biological studies  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (carrier; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Toxins  
 Toxoids  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (diphtheria; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Lipoproteins  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (e; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Toxins  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (exotoxin A, *Pseudomonas*; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT *Pseudomonas*  
 (exotoxin A; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Gene, microbial  
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); PREP (Preparation)  
 (for *Haemophilus influenzae* protein "e"; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT *Escherichia coli*  
 (heat labile toxin; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Toxins  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (heat-labile, *Escherichia coli*; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Antibodies  
 RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (monoclonal; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)
- IT Ear  
 (otitis media; protein "e" and gene of *Haemophilus*)

influenza and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT Rotavirus  
(particles; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT Antiserums  
Bacterium (genus)  
DNA sequences  
Fungi  
    *Haemophilus influenzae*  
Microorganism  
    *Moraxella catarrhalis*  
Parasite  
Protein sequences  
Respiratory syncytial virus  
Staphylococcus aureus  
Streptococcus pneumoniae  
Streptococcus pyogenes  
Vaccines  
Virus  
(protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT Opsonins  
RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT Antigens  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT Oligosaccharides, biological studies  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT Polysaccharides, biological studies  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT Fusion proteins (chimeric proteins)  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT Toxins  
Toxoids  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(tetanus; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT 145110-32-3, Lipoprotein e (*Haemophilus influenzae* clone pPX513 gene hel precursor protein moiety reduced)  
RL: PRP (Properties)  
(amino acid sequence; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT 135622-17-2, DNA (*Haemophilus influenzae* type b strain Eagan clone pPX513 lipoprotein e gene)  
RL: PRP (Properties)  
(nucleotide sequence; protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

IT 82197-76-0, Polyribosylribitolphosphate  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(protein "e" and gene of *Haemophilus influenzae* and antibodies and vaccines for nontypeable *Haemophilus influenzae*)

=> d all 130 tot

L30 ANSWER 1 OF 3 HCAPLUS COPYRIGHT 2004 ACS on STN  
AN 2002:237317 HCAPLUS  
DN 136:261813  
ED Entered STN: 28 Mar 2002  
TI Transferrin receptor-encoding genes from *Haemophilus influenzae* strains and their uses for diagnostics and medical treatment  
IN Loosmore, Sheena M.; Harkness, Robin E.; Schryvers, Anthony B.;

Search done by Noble Jarrell

Chong, Pele; Gray-Owen, Scott; Yang, Yan-ping; Murdin, Andrew  
D.; Klein, Michel H.

PA Aventis Pasteur Limited, Can.

SO U.S., 280 pp., Cont.-in-part of Ser. No. US 1995-483577, filed on 7 Jun  
1995, now

CODEN: USXXAM

DT Patent

LA English

IC ICM A61K038-21

ICS A61K038-16; A01N063-00; C12P019-34

NCL 424256100

CC 15-2 (Immunochemistry)

Section cross-reference(s): 1, 3, 6

FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6361779	B1	20020326	US 1996-649518	19960517
	US 5922562	A	19990713	US 1994-337483	19941108
	US 6015688	A	20000118	US 1995-483577	19950607
	CA 2223503	AA	19961219	CA 1996-2223503	19960607
	WO 9640929	A2	19961219	WO 1996-CA399	19960607
	WO 9640929	A3	19970306		
	W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI				
	RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN				
	AU 9661177	A1	19961230	AU 1996-61177	19960607
	AU 716506	B2	20000224		
	EP 833920	A2	19980408	EP 1996-918543	19960607
	EP 833920	B1	20040818		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
	JP 11506335	T2	19990608	JP 1997-500057	19960607
	JP 3516688	B2	20040405		
	BR 9608482	A	20010731	BR 1996-8482	19960607
	AT 274059	E	20040915	AT 1996-918543	19960607
	US 2003088086	A1	20030508	US 2002-43344	20020114
PRAI	US 1993-148968	B2	19931108		
	US 1993-175116	B2	19931229		
	US 1994-337483	A2	19941108		
	US 1995-483577	A2	19950607		
	US 1996-649518	A	19960517		
	WO 1996-CA399	W	19960607		

#### CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
US 6361779	ICM	A61K038-21
	ICS	A61K038-16; A01N063-00; C12P019-34
	NCL	424256100
US 6361779	ECLA	C07K014/285
US 5922562	ECLA	C07K014/285
US 6015688	ECLA	C07K014/285
WO 9640929	ECLA	C07K014/285
US 2003088086	ECLA	C07K014/285

AB Purified and isolated genes are provided which encodes transferrin receptor proteins Tbp1 and/or Tbp2 of Haemophilus influenzae type b strains DL63, Eagan, Minna, PAK12085, and SB33 and the non-typeable strains SB12, SB29, SB30, and SB32. The nucleic acid sequence may be used to produce peptides free of contaminants derived from bacteria normally containing the Tbp1 or Tbp2 proteins for purposes of diagnostics and medical treatment. Furthermore, the nucleic acid mol. may be used in the diagnosis of infection. Also provided are recombinant Tbp1 or Tbp2 and methods for purification of the same. Live vectors expressing epitopes of transferrin receptor protein for vaccination are provided. Thus, poliovirus vectors incorporating the H. influenzae strain DL63 Tbp2 are neutralized by guinea-pig antisera raised against peptide LEGGFYGP, indicating that the viruses express this sequence in an antigenically recognizable form. Since H. influenzae Tbp2 is produced in low amts by Escherichia coli, the Eagan strain Tbp2 gene was truncated from its 3'-end using an Erase-a-base kit to produce a number of truncated analogs of Tbp2. The yield of Eagan rTbp2 is significantly increased by truncation of the C-terminal region of the protein. The infant rat model of bacteremia confirms the protective ability of anti-(truncated analogs of transferrin receptor protein Tbp2) antibodies even after removal of up to half of the



Tbp2 sequence.

ST transferrin receptor gene sequence Haemophilus; antigenicity transferrin receptor Haemophilus; vaccination transferrin receptor Haemophilus

IT Plasmid vectors  
(JD-1468-29 and JD-1424-2-8, for expression in Escherichia coli; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Gene, microbial  
Transferrin receptors  
RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); DGN (Diagnostic use); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(Tbp1 and Tbp2; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Moraxella catarrhalis  
(antiserum cross-reactivity with; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Immunoassay  
(enzyme, development and cross-reactivity of; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Diagnosis  
(mol.; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Escherichia coli  
(plasmid vectors JD-1468-29 and JD-1424-2-8 for expression in; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Viral vectors  
(poliovirus type 1; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT DNA sequences  
Epitopes  
Haemophilus influenzae  
Molecular cloning  
Protein sequences  
Vaccines  
(transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Promoter (genetic element)  
RL: BUU (Biological use, unclassified); PRP (Properties); BIOL (Biological study); USES (Uses)  
(transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Human poliovirus 1  
(vector; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT 405178-16-7P 405178-17-8P 405178-18-9P 405178-19-0P 405178-20-3P  
405178-21-4P 405178-23-6P 405178-24-7P 405178-26-9P 405178-28-1P  
405178-30-5P 405178-32-7P  
RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); DGN (Diagnostic use); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(amino acid sequence; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT 405178-36-1P 405178-37-2P 405178-38-3P 405178-39-4P 405178-40-7P  
405178-41-8P 405178-42-9P 405178-43-0P 405178-44-1P 405178-45-2P  
405178-46-3P 405178-47-4P 405178-48-5P  
RL: BPN (Biosynthetic preparation); PAC (Pharmacological activity); PRP (Properties); BIOL (Biological study); PREP (Preparation)  
(amino acid sequence; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT 167769-62-2 167769-63-3  
RL: PAC (Pharmacological activity); PRP (Properties); BIOL (Biological study)  
(antigenic peptide epitope; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT 405178-12-3P 405178-13-4P 405178-14-5P 405178-15-6P 405178-22-5P  
405178-25-8P 405178-27-0P 405178-29-2P 405178-31-6P 405178-33-8P

405178-34-9P 405178-35-0P

RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified);  
DGN (Diagnostic use); PRP (Properties); THU (Therapeutic use); BIOL  
(Biological study); PREP (Preparation); USES (Uses)

(nucleotide sequence; transferrin receptor-encoding genes from  
Haemophilus influenzae strains and their uses for diagnostics and  
medical treatment)

IT	405180-53-2	405180-54-3	405180-55-4	405180-56-5	405180-57-6
	405180-58-7	405180-59-8	405180-60-1	405180-68-9	405180-69-0
	405180-70-3	405180-71-4	405180-72-5	405180-73-6	405180-74-7
	405180-75-8	405180-76-9	405180-77-0	405180-78-1	

RL: PRP (Properties)

(unclaimed nucleotide sequence; transferrin receptor-encoding genes  
from Haemophilus influenzae strains and their uses for diagnostics and  
medical treatment)

IT	405180-06-5	405180-07-6	405180-08-7	405180-09-8	405180-10-1
	405180-11-2	405180-12-3	405180-13-4	405180-14-5	405180-15-6
	405180-16-7	405180-17-8	405180-18-9	405180-19-0	405180-20-3
	405180-21-4	405180-22-5	405180-23-6	405180-24-7	405180-25-8
	405180-26-9	405180-27-0	405180-28-1	405180-29-2	405180-30-5
	405180-31-6	405180-32-7	405180-33-8	405180-34-9	405180-35-0
	405180-36-1	405180-37-2	405180-38-3	405180-39-4	405180-40-7
	405180-41-8	405180-42-9	405180-43-0	405180-44-1	405180-45-2
	405180-46-3	405180-47-4	405180-48-5	405180-49-6	405180-50-9
	405180-51-0	405180-52-1	405180-61-2	405180-62-3	405180-63-4
	405180-64-5	405180-65-6	405180-66-7	405180-67-8	

RL: PRP (Properties)

(unclaimed protein sequence; transferrin receptor-encoding genes from  
Haemophilus influenzae strains and their uses for diagnostics and  
medical treatment)

IT	161228-75-7	229032-30-8	229032-31-9	229032-32-0	229032-33-1
	229032-34-2	229032-35-3	229032-36-4	229032-37-5	229032-38-6
	229032-39-7	229032-40-0	229032-41-1	229032-42-2	229032-43-3
	229032-44-4	229032-45-5	229032-46-6	229032-47-7	229032-48-8
	229157-61-3	229157-62-4	229157-63-5	229157-64-6	229157-65-7
	404572-60-7	404572-61-8	404572-62-9	404572-63-0	404572-64-1
	404572-65-2	404572-66-3	404572-67-4	404572-68-5	404572-69-6
	404572-70-9	404572-72-1	404572-74-3	404572-76-5	404572-78-7
	404572-80-1	404572-82-3	404572-84-5		

RL: PRP (Properties)

(unclaimed sequence; transferrin receptor-encoding genes from  
Haemophilus influenzae strains and their uses for diagnostics and  
medical treatment)

RE.CNT 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Anon; WO 9412641 1994 HCAPLUS
- (2) Anon; WO 9513370 1995 HCAPLUS
- (3) Barber; US 5194254 A 1993 HCAPLUS
- (4) Frasch; US 4601903 A 1986
- (5) Ghrayeb; Embo J 1984, V3, P2437 HCAPLUS
- (6) Gordon; US 4496538 A 1985 HCAPLUS
- (7) Gray-Owen; Infect Immun 1995, V4, P1201
- (8) Griffiths; Fems Microbiol Lett 1993, V109(1), P85 HCAPLUS
- (9) Holland; Infect Immun 1992, V60, P2986 HCAPLUS
- (10) Hopp, T; Journal of Immunological Methods 1986, V88, P1 HCAPLUS
- (11) Itakura; Science 1977, V198, P1056 HCAPLUS
- (12) Jarosik; Infection and Immunity 1994, V62, P2470 HCAPLUS
- (13) Legrain; Gene 1993, V130, P73 HCAPLUS
- (14) Lockhoff; Chem Ind Ed Engl 1991, V30, P1611
- (15) Lockoff; US 4855283 A 1989 HCAPLUS
- (16) Mickelsen; Infect Immun 1981, V33, P555 HCAPLUS
- (17) Milich; US 4599230 A 1986 HCAPLUS
- (18) Milich; US 4599231 A 1986 HCAPLUS
- (19) Moloney; US 4258029 A 1981 HCAPLUS
- (20) Morton; Infection and Immunity 1993, V61, P4033 HCAPLUS
- (21) Mordin; J Gen Viral 1992, V73, P607 HCAPLUS
- (22) Mordin; Microbial Pathogenesis 1991, V10, P27 HCAPLUS
- (23) Nixon-George; J Immunol 1990, V14, P4798
- (24) Ogunnariwo; Avian Dis 1992, V36, P655 MEDLINE
- (25) O'Hagan; Clin Pharmacokinet 1992, V22, P1 MEDLINE
- (26) Panezutti; Infection and Immunity 1993, V61, P1867 HCAPLUS
- (27) Roosi-Campos; Vaccine 1992, V10, P512
- (28) Sambrook; Molecular Cloning, A Lab Manual V3, P16.2
- (29) Schryvers; US 5141743 A 1992 HCAPLUS
- (30) Schryvers; J Infect Dis 1992, V165(suppl 1), PS103
- (31) Schryvers; Molec Microbiol 1988, V2, P467 HCAPLUS

Search done by Noble Jarrell

- (32) Schyrrvers; Can J Microbiol 1989, V35, P409  
 (33) Schyrrvers; Med Microbiol 1989, V29, P121  
 (34) Short; Nucl Acids Res 1988, V16, P7583 HCAPLUS  
 (35) Studier; US 4952496 A 1990 HCAPLUS  
 (36) Ulmer; Curr Opinion Invest Drugs 1993, V2(9), P983  
 (37) Vanderwerf; Proc Natl Acad Sci 1986, V83, P2330 HCAPLUS  
 (38) Vyas; US 4596792 A 1986 HCAPLUS  
 (39) Wilton; FEMS Microbiology Letters 1993, V107, P59 HCAPLUS

L30 ANSWER 2 OF 3 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 2002:213747 HCAPLUS  
 DN 136:242991  
 ED Entered STN: 21 Mar 2002  
 TI Transferrin receptor-encoding genes from Haemophilus influenzae strains  
 and their uses for diagnostics and medical treatment  
 IN Loosmore, Sheena M.; Harkness, Robin E.; Schryvers, Anthony B.;  
 Chong, Pele; Gray-Owen, Scott; Yang, Yan-Ping; Murdin, Andrew  
 D.; Klein, Michel H.  
 PA Aventis Pasteur Limited, Can.  
 SO U.S., 264 pp., Cont.-in-part of U.S. Ser. No. 175,116, abandoned.  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 IC ICM C12N001-21  
 ICS C12N005-10; C12N015-03; C12N015-63  
 NCL 435252300  
 CC 3-3 (Biochemical Genetics)  
 Section cross-reference(s): 1, 6, 15

## FAN.CNT 5

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6358727	B1	20020319	US 1996-637654	19960805
WO 9513370	A1	19950518	WO 1994-CA616	19941107
W: AU, BR, CA, CN, FI, JP, KR, NO, NZ, RU, UA, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
PRAI US 1993-148968	B2	19931108		
US 1993-175116	B2	19931229		
WO 1994-CA616	W	19941107		

## CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
US 6358727	ICM	C12N001-21
	ICS	C12N005-10; C12N015-03; C12N015-63
	NCL	435252300
US 6358727	ECLA	C07K014/285
WO 9513370	ECLA	C07K014/285

- AB Purified and isolated genes are provided which encodes transferrin receptor proteins Tbp1 and/or Tbp2 of Haemophilus influenzae type b strains DL63, Eagan, MinnA, PAK12085, and SB33 and the non-typeable strains SB12, SB29, SB30, and SB32. The nucleic acid sequence may be used to produce peptides free of contaminants derived from bacteria normally containing the Tbp1 or Tbp2 proteins for purposes of diagnostics and medical treatment. Furthermore, the nucleic acid mol. may be used in the diagnosis of infection. Also provided are recombinant Tbp1 or Tbp2 and methods for purification of the same. Live vectors expressing epitopes of transferrin receptor protein for vaccination are provided. Thus, poliovirus vectors incorporating the H. influenzae strain DL63 Tbp2 are neutralized by guinea-pig antisera raised against peptide LEGGFYGP, indicating that the viruses express this sequence in an antigenically recognizable form.
- ST transferrin receptor gene sequence Haemophilus; antigenicity transferrin receptor Haemophilus; vaccination transferrin receptor Haemophilus
- IT Plasmid vectors  
 (JD-1468-29 and JD-1424-2-8, for expression in Escherichia coli; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)
- IT Gene, microbial  
 Transferrin receptors  
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); DGN (Diagnostic use); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (Tbp1 and Tbp2; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)
- IT Moraxella catarrhalis  
 (antiserum cross-reactivity with; transferrin receptor-encoding genes

from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Immunoassay  
(enzyme, development and cross-reactivity of; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Diagnosis  
(mol.; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Escherichia coli  
(plasmid vectors JD-1468-29 and JD-1424-2-8 for expression in; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Viral vectors  
(poliovirus type 1; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT DNA sequences  
Epitopes  
Haemophilus influenzae  
Molecular cloning  
Protein sequences  
Vaccines  
(transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Promoter (genetic element)  
RL: BUU (Biological use, unclassified); PRP (Properties); BIOL (Biological study); USES (Uses)  
(transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT Human poliovirus 1  
(vector; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT 404796-06-1P 404796-07-2P 404796-08-3P 404796-09-4P 404796-10-7P  
404796-11-8P 404796-12-9P 404796-13-0P 404796-14-1P 404796-15-2P  
404796-16-3P 404796-17-4P  
RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); DGN (Diagnostic use); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(amino acid sequence; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT 167769-62-2 167769-63-3  
RL: PAC (Pharmacological activity); PRP (Properties); BIOL (Biological study)  
(antigenic peptide epitope; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT 404795-94-4P 404795-95-5P 404795-96-6P 404795-97-7P 404795-98-8P  
404795-99-9P 404796-00-5P 404796-01-6P 404796-02-7P 404796-03-8P  
404796-04-9P 404796-05-0P  
RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); DGN (Diagnostic use); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(nucleotide sequence; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT 404818-86-6 404818-87-7 404818-88-8 404818-89-9 404818-90-2  
404818-91-3 404818-92-4 404818-93-5 404819-01-8 404819-02-9  
404819-03-0 404819-04-1 404819-05-2 404819-06-3 404819-07-4  
404819-08-5 404819-09-6 404819-10-9 404819-11-0  
RL: PRP (Properties)  
(unclaimed nucleotide sequence; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT 404818-39-9 404818-40-2 404818-41-3 404818-42-4 404818-43-5  
404818-44-6 404818-45-7 404818-46-8 404818-47-9 404818-48-0  
404818-49-1 404818-50-4 404818-51-5 404818-52-6 404818-53-7  
404818-54-8 404818-55-9 404818-56-0 404818-57-1 404818-58-2  
404818-59-3 404818-60-6 404818-61-7 404818-62-8 404818-63-9  
404818-64-0 404818-65-1 404818-66-2 404818-67-3 404818-68-4  
404818-69-5 404818-70-8 404818-71-9 404818-72-0 404818-73-1  
404818-74-2 404818-75-3 404818-76-4 404818-77-5 404818-78-6  
404818-79-7 404818-80-0 404818-81-1 404818-82-2 404818-83-3  
404818-84-4 404818-85-5 404818-86-6 404818-87-7 404818-88-8

404818-97-9 404818-98-0 404818-99-1 404819-00-7

RL: PRP (Properties)

(unclaimed protein sequence; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

IT	161228-75-7	229032-30-8	229032-31-9	229032-32-0	229032-33-1
	229032-34-2	229032-35-3	229032-36-4	229032-37-5	229032-38-6
	229032-39-7	229032-40-0	229032-41-1	229032-42-2	229032-43-3
	229032-44-4	229032-45-5	229032-46-6	229032-47-7	229032-48-8
	229157-61-3	229157-62-4	229157-63-5	229157-64-6	229157-65-7
	404572-60-7	404572-61-8	404572-62-9	404572-63-0	404572-64-1
	404572-65-2	404572-66-3	404572-67-4	404572-68-5	404572-69-6
	404572-70-9	404572-72-1	404572-74-3	404572-76-5	404572-78-7
	404572-80-1	404572-82-3	404572-84-5		

RL: PRP (Properties)

(unclaimed sequence; transferrin receptor-encoding genes from Haemophilus influenzae strains and their uses for diagnostics and medical treatment)

RE.CNT 60 THERE ARE 60 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Anon; WO 9306861 1993 HCAPLUS
- (2) Anon; WO 9308283 1993 HCAPLUS
- (3) Barcak; Methods Enzymol 1991, V204, P321 HCAPLUS
- (4) Berber; US 5194254 A 1993 HCAPLUS
- (5) Berkowitz; J Pediatr 1987, V110, P509 MEDLINE
- (6) Black; Pediatr Infect Dis J 1991, V10, P97 MEDLINE
- (7) Bluestone, N; Engl J Med 1982, V306, P1399
- (8) Cheng; Nature 1978, V375, P615
- (9) Chou; Annual Reviews of Biochemistry 1978, V47, P251 HCAPLUS
- (10) Claesson; J Pediatr 1989, V114, P97 MEDLINE
- (11) Cornelissen; J of Bacteriology 1992, V174, P5788 HCAPLUS
- (12) Danve; Vaccine 1993, V11, P1214 HCAPLUS
- (13) Derea; Nature 1989, V342, P651
- (14) Freach; US 4601903 A 1986
- (15) George; Macromolecular Sequencing and Synthesis 1988, P127
- (16) Gerlach; Infection and Immunity 1992, V60, P3253 HCAPLUS
- (17) Ghayeb; The Embo J 1984, V3, P2437 HCAPLUS
- (18) Goeddel; Nature 1979, V281, P544 HCAPLUS
- (19) Goldberg; Mol Microbiology 1992, V6, P2407 HCAPLUS
- (20) Gordon; US 4496538 A 1985 HCAPLUS
- (21) Gray-Owen; Infect Immun 1995, V63(4), P1201 HCAPLUS
- (22) Gray-Owen; Microbial Pathogenesis 1993, V14 MEDLINE
- (23) Griffiths; Fems Microbiol Lett 1993, V109(1), P85 HCAPLUS
- (24) Harkness; J Bacteriol 1992, V174, P2425 HCAPLUS
- (25) Holland; FEMS, Microbiology Letters 1991, V77, P283 HCAPLUS
- (26) Holland; Infection and Immunity 1992, V60, P2986 HCAPLUS
- (27) Hopp, T; Journal of Immunological Methods 1986, V88, P1 HCAPLUS
- (28) Itakura; Science 1977, V198, P1058
- (29) Jaroalk; Infection and Immunity 1994, V62, P2470
- (30) Legrain; Gene 1993, V130, P73 HCAPLUS
- (31) Lockhoff; US 4855283 A 1989 HCAPLUS
- (32) Lockhoff; Chem Int Ed Engl 1991, V30, P1611
- (33) McGeoch; J Gen Virol 1988, V69, P1531 HCAPLUS
- (34) Mickelsen; Infect Immun 1981, V33, P555 HCAPLUS
- (35) Millich; US 4599230 A 1986 HCAPLUS
- (36) Millich; US 4599231 A 1986 HCAPLUS
- (37) Moloney; US 4258029 A 1981 HCAPLUS
- (38) Morton; Infection and Immunity 1993, V61, P4033 HCAPLUS
- (39) Murdin; J Gen Viral 1992, V73, P607 HCAPLUS
- (40) Murdin; Microbial Pathogenesis 1991, V10, P27 HCAPLUS
- (41) Nixon-George; J Immunol 1990, V14, P4798
- (42) Ogunnariwo; Avian Dis 1992, V38, P855
- (43) O'Hagen; Clin Pharmacokinet 1992, V22, P1
- (44) Panazutti; Infection and Immunity 1993, V61, P1867
- (45) Poulsen; Molecular Microbiology 1992, V6, P895 HCAPLUS
- (46) Roosi-Campos; Vaccine 1992, V10, P512
- (47) Sambrook; Molecular Biology, A Laboratory Manual 2nd Ed 1989, V3, P16.2
- (48) Schryvers; US 5141743 A 1992 HCAPLUS
- (49) Schryvers; Can J Microbiol 1989, V35, P409 HCAPLUS
- (50) Schryvers; J Infect Dis 1992, V165(suppl 1), P5103
- (51) Schryvers; Molec Microbiol 1988, V2, P467 HCAPLUS
- (52) Schyvers; Med Microbiol 1989, V29, P121
- (53) Short; Nucl Acids Res 1988, V16, P7533
- (54) Stevenson; Infection and Immunity 1992, V60(8), P2391
- (55) Studier; US 4952498 A 1990
- (56) Thomas; Methods in Enzymology 1990, V182, P499 HCAPLUS

Search done by Noble Jarrell

- (57) Ulmer; Curr Opin Invest Drugs 1993, V2(9), P983  
 (58) Vandarwerf; Proc Natl Acad Sci 1986, V83, P2330  
 (59) Vyae; US 4596792 A 1986 HCAPLUS  
 (60) Weismuller; Vaccine 1989, V8, P29

L30 ANSWER 3 OF 3 HCAPLUS COPYRIGHT 2004 ACS on STN  
 AN 2001:63846 HCAPLUS  
 DN 134:120915  
 ED Entered STN: 26 Jan 2001  
 TI Multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis  
 IN Loosmore, Sheena M.; Yang, Yan-Ping; Klein, Michel H.; Sasaki, Ken  
 PA Connaught Laboratories Limited, Can.  
 SO PCT Int. Appl., 58 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM A61K039-00  
 CC 63-3 (Pharmaceuticals)  
 Section cross-reference(s): 15

## FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001005424	A2	20010125	WO 2000-CA811	20000711
WO 2001005424	A3	20010802		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
US 6391313	B1	20020521	US 1999-353617	19990715
CA 2378862	AA	20010125	CA 2000-2378862	20000711
EP 1200122	A2	20020502	EP 2000-945494	20000711
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				
AU 767096	B2	20031030	AU 2000-59586	20000711
PRAI US 1999-353617	A	19990715		
WO 2000-CA811	W	20000711		

## CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
------------	-------	------------------------------------

WO 2001005424	ICM	A61K039-00
US 6391313	ECLA	A61K039/116

AB A multi-valent immunogenic composition confers protection on an immunized host against infection caused by both Haemophilus influenzae and Moraxella catarrhalis. Such composition comprises at least four antigens comprising at least one antigen from Haemophilus influenzae, and at least one antigen from Moraxella catarrhalis. Three of the antigens are adhesins. High mol. weight (HMW) proteins and Haemophilus influenzae adhesin (Hia) proteins of non-typeable Haemophilus and a 200 kDa outer membrane protein of Moraxella catarrhalis comprise the adhesin components while the other antigen is a non-proteolytic analog of Hin47 protein. Each component does not impair the immunogenicity of the others. The multi-valent immunogenic composition may be combined with DTP component vaccines, which may also include non-virulent poliovirus and PRP-T, to provide a component vaccine without impairment of the immunogenic properties of the other antigens.

ST adhesin antigen vaccine Haemophilus Moraxella

IT Proteins, specific or class

RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)

(HMW1; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

IT Proteins, specific or class

RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)

(HMW2; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

*Applicants*

- IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (Hin47; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (Hsf (Haemophilus surface fibril); multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (OMP (outer membrane protein); multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Immunostimulants  
 (adjuvants; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Agglutinins and Lectins  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
 (agglutinogens; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Adhesins  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (antigenic; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Toxoids  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
 (diphtheria; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Organelle  
 (fibril, surface; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Hemagglutinins  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
 (filamentous; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Chinchilla  
 Haemophilus influenzae  
 Molecular cloning  
 Molecular weight distribution  
 Moraxella catarrhalis  
 Polyacrylamide gel electrophoresis  
 Vaccines  
 (multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Antigens  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Heat-shock proteins  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)  
 (non-proteolytic; multicomponent vaccine to protect against disease

- caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Human poliovirus  
(non-virulent; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Ear  
(otitis, otitis media; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Agglutinins and Lectins  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(pertactins; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Toxoids  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(pertussis; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Mutation  
(substitution; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT Toxoids  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(tetanus; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT 9001-92-7, Proteinase  
RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence)  
(activity levels; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT 7784-30-7, Aluminum phosphate 21645-51-2, Aluminum hydroxide, biological studies  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(adjuvant; multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)
- IT 151-21-3, Sds, analysis  
RL: ARU (Analytical role, unclassified); ANST (Analytical study)  
(multicomponent vaccine to protect against disease caused by Haemophilus influenzae and Moraxella catarrhalis)

=> b wpix

FILE 'WPIX' ENTERED AT 13:10:45 ON 16 DEC 2004

COPYRIGHT (C) 2004 THE THOMSON CORPORATION

FILE LAST UPDATED: 13 DEC 2004 <20041213/UP>  
MOST RECENT DERWENT UPDATE: 200480 <200480/DW>  
DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,  
PLEASE VISIT:  
[http://www.stn-international.de/training\\_center/patents/stn\\_guide.pdf](http://www.stn-international.de/training_center/patents/stn_guide.pdf) <<<

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE  
<http://thomsonderwent.com/coverage/latestupdates/> <<<

>>> FOR INFORMATION ON ALL DERWENT WORLD PATENTS INDEX USER  
GUIDES, PLEASE VISIT:  
<http://thomsonderwent.com/support/userguides/> <<<

>>> NEW! FAST-ALERTING ACCESS TO NEWLY-PUBLISHED PATENT  
DOCUMENTATION NOW AVAILABLE IN DERWENT WORLD PATENTS INDEX  
FIRST VIEW - FILE WPIFV.  
FOR FURTHER DETAILS: <http://www.thomsonderwent.com/dwpifv> <<<

>>> NEW DISPLAY FORMAT HITSTR ADDED ALLOWING DISPLAY OF  
HIT STRUCTURES WITHIN THE BIBLIOGRAPHIC DOCUMENT <<<

>>> SMILES and ISOSMILES strings are no longer available as  
Derwent Chemistry Resource display fields <<<

Search done by Noble Jarrell



=> d all 154 tot

L54 ANSWER 1 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2004-774372 [76] WPIX

DNC C2004-271112

TI New immunogenic composition comprises a major outer membrane protein of a strain of *Chlamydia pneumoniae*, and a 76 kDa protein of a strain of *C. pneumoniae*, useful as a vaccine for treating or preventing *Chlamydia* infections.

DC B04 D16

IN DUNN, P L; MURDIN, A D

PA (AVET) AVENTIS PASTEUR LTD

CYC 1

PI US 6811783 B1 20041102 (200476)\* 31 A61K039-02 <--

ADT US 6811783 B1 US 1999-391606 19990907

PRAI US 1999-391606 19990907

IC ICM A61K039-02

ICS A61K039-00; C07H021-04; C07K001-00

AB US 6811783 B UPAB: 20041125

NOVELTY - An immunogenic composition comprises a first plasmid vector comprising a first nucleotide sequence encoding a major outer membrane protein (MOMP) of a strain of *Chlamydia pneumoniae*, and a second plasmid vector comprising a second nucleotide sequence encoding a 76 kDa protein of a strain of *C. pneumoniae*, is new.

DETAILED DESCRIPTION - An immunogenic composition comprises a first plasmid vector comprising a first nucleotide sequence encoding a major outer membrane protein of a strain of *C. pneumoniae*, the first nucleotide sequence is selected from 3 sequences comprising 1426, 1301, or 1101 bp (SEQ ID NO. 12, 13, or 14) or encoding a MOMP having an amino acid sequence comprising 394 or 367 amino acids (SEQ ID NO. 15 or 16), and a first promoter sequence operatively coupled to the first nucleotide sequence for expression of the MOMP in a host; and a second plasmid vector comprising a second nucleotide sequence encoding a 76 kDa protein of a strain of *C. pneumoniae*, the second nucleotide sequence is selected from 4 sequences comprising 2545, 651, 1470, or 1389 bp (SEQ ID NO. 1, 2, 3, or 4), and a second promoter sequence operatively coupled to the second nucleotide sequence for expression of the 76 kDa protein in a host; and a pharmaceutical carrier.

All sequences are defined in the specification.

ACTIVITY - Antibacterial.

MECHANISM OF ACTION - Vaccine.

Groups of 7-9 week old male Balb/c mice were immunized intramuscularly (i.m.) and intranasally (i.n.) with plasmids pCA76kDa and pCAMOMP containing the coding sequences of *C. pneumoniae* 76 kDa and MOMP, respectively. Saline or plasmid vectors containing non-protective inserted chlamydial genes were given to groups of control animals. Results showed an increased protection afforded by the combination of the two constructs. It showed that mice immunized intramuscularly and intranasally with both pCA76kDa and pCAMOMP had chlamydial lung titers less than 6700 in 6 of 6 cases, where the range of values for control mice with saline were 15000-106100 IFU/lung in 20 out of 23 cases, and 12600-80600 IFU/lung in 11 out of 12 cases for mice immunized with the vectors containing non-protective genes.

USE - The immunogenic composition is useful as a vaccine for immunizing a host against disease caused by infection with a strain of *Chlamydia*. It is also useful for treating or preventing *Chlamydia* infections.

Dwg.0/5

FS CPI

FA AB; DCN

MC CPI: B04-E08; B14-A01A; B14-S03; B14-S09; B14-S11B;  
D05-H07

L54 ANSWER 2 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-648559 [74] WPIX

DNN N2001-484575 DNC C2001-191446

TI Novel polypeptides from *Chlamydia pneumoniae* and genes encoding the polypeptide, useful for immunization of host e.g. human against disease caused by infection by a strain of *Chlamydia*.

DC B04 D16 S03

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J

PA (AVET) AVENTIS PASTEUR LTD; (DUNN-I) DUNN P; (MURD-I) MURDIN A

D; (OOME-I) OOMEN R P; (WANG-I) WANG J

CYC 95

PI WO 2001075114 A2 20011011 (200174)\* EN 90 C12N015-31

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
 NL OA PT SD SE SL SZ TR TZ UG ZW  
 W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK  
 DM DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ  
 LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD  
 SE SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW  
 AU 2001048178 A 20011015 (200209) C12N015-31  
 US 2002082402 A1 20020627 (200245) C07H021-02  
 US 2003224004 A1 20031204 (200380) A61K039-00 <--  
 ADT WO 2001075114 A2 WO 2001-CA462 20010404; AU 2001048178 A AU 2001-48178  
 20010404; US 2002082402 A1 Provisional US 2000-194477P 20000404, US  
 2001-824588 20010403; US 2003224004 A1 Provisional US 2000-194477P  
 20000404, Cont of US 2001-824588 20010403, US 2003-359289 20030206  
 FDT AU 2001048178 A Based on WO 2001075114  
 PRAI US 2000-194477P 20000404; US 2001-824588 20010403;  
 US 2003-359289 20030206  
 IC ICM A61K039-00; C07H021-02; C12N015-31  
 ICS A61K039-118; A61K039-38; A61K039-40;  
 C07H021-04; C07K001-00; C07K014-00; C07K014-295; C07K016-12;  
 C07K017-00; C12N015-11; C12N015-62; C12Q001-68; G01N033-53;  
 G01N033-68  
 AB WO 200175114 A UPAB: 20011217  
 NOVELTY - A transmembrane polypeptide from Chlamydia, preferably C.  
 pneumoniae comprising a 579 residue amino acid sequence, fully defined in  
 the specification, an immunogenic fragment of at least 12 consecutive  
 amino acids of S1, or a polypeptide modified without loss of  
 immunogenicity and having at least 75 % identity to them, is new.  
 DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the  
 following:  
 (1) a nucleic acid molecule (II) comprising a sequence encoding (I),  
 a 1940 nucleotide sequence (S2), fully defined in the specification, a  
 sequence encoding a polypeptide encoded by S2, a sequence comprising at  
 least 38 consecutive nucleotides of them, or a sequence encoding a  
 polypeptide having at least 75 % identity to a polypeptide encoded by S2;  
 (2) a nucleic acid molecule (IIa) comprising a sequence which is  
 antisense to (II);  
 (3) a nucleic acid molecule (IIb) comprising a sequence encoding a  
 fusion protein (FP) comprising a polypeptide encoded by (II) and a second  
 polypeptide;  
 (4) a vaccine (IIIa) comprising a vaccine vector and at least one  
 first nucleic acid encoding (I) or FP, which is capable of being  
 expressed, and optionally the vaccine comprises a second nucleic acid  
 encoding and capable of expressing an additional polypeptide which  
 enhances the immune response to the polypeptide expressed by the first  
 nucleic acid;  
 (5) a vaccine (IIIb) comprising (II)-(IIb) and a vaccine vector,  
 where (II)-(IIb) is expressed as a polypeptide, and optionally the vaccine  
 comprises a second nucleic acid encoding an additional polypeptide which  
 enhances the immune response to the polypeptide expressed by (II)-(IIb);  
 (6) a pharmaceutical composition (PC) comprising (II)-(IIb), (IIIa)  
 or (IIIb);  
 (7) a unicellular host (IV) transformed with (II)-(IIb);  
 (8) an isolated nucleic acid probe of 5-100 nucleotides which  
 hybridizes under stringent conditions to S2, or its complement or  
 antisense sequence;  
 (9) an isolated primer of 10-40 nucleotides which hybridizes under  
 stringent conditions to S2, or its complement or antisense sequence;  
 (10) a polypeptide (Ia) encoded by (II)-(IIb);  
 (11) a fusion protein (FP) comprising (I) or (Ia), and a second  
 polypeptide;  
 (12) producing (I) and FP;  
 (13) an antibody (Ab) against (I) or FP;  
 (14) a vaccine (IIIC) comprising (I), a polypeptide encoded by (II),  
 or FP comprising (I) and a second polypeptide, and optionally comprising  
 an additional polypeptide which enhances the immune response to the first  
 polypeptide;  
 (15) a vaccine (IIId) comprising at least one first polypeptide  
 selected from (I) or FP, and optionally comprising an additional  
 polypeptide which enhances the immune response to the first polypeptide;  
 (16) a pharmaceutical composition comprising (I), FP, (IIIC) or Ab;  
 (17) a diagnostic kit comprising instructions for use and a component  
 selected from (I), (II), FP and Ab;  
 (18) identifying (I) or FP which induces an immune response effective  
 to prevent or lessen the severity of Chlamydia infection in a mammal  
 previously immunized with polypeptide, by immunizing a mouse with (I) or  
 FP, and inoculating the immunized mouse with Chlamydia, where (I) or FP

are identified;

(19) an expression plasmid pCACPNM643 given in the specification;

(20) a nucleic acid molecule comprising a sequence (S7); and

(21) a peptide comprising a sequence (S8).

(S7) is ataagaatgcggccgccaccatgcagaagcatccttccttttattc or gcgcccggatcccagatccttgacagcggg.

(S8) is AlaLysTyrArgLysLysGlnGluAlaSerValLysLysTyrGln or TyrLeuPhePheProGlyTyrTyrThr.

ACTIVITY - Antibacterial.

MECHANISM OF ACTION - Vaccine (claimed); gene therapy.

Groups of 7-9 week old male Balb/c mice (8-10 per group) were immunized intramuscularly (i.m.) and intranasally (i.n.) with plasmid DNA containing Chlamydia pneumoniae transmembrane protein gene. Saline or plasmid vector lacking an inserted Chlamydial gene was given to groups of control animals. At week 8, immunized mice were inoculated i.n. with 5 multiply 105 infection forming units (IFU) of C. pneumoniae strain AR39 to test their ability to limit the growth of a sublethal C. pneumoniae challenge. Lungs were taken from mice at day 9 post-challenge and immediately homogenized for analyzing the presence of Chlamydial inclusions using convalescent sera from rabbits infected with C. pneumoniae and metal-enhanced DAB as a peroxidase substrate. The results showed that mice immunized with pCACPNM643 had Chlamydial lung titers less than 60000 in 5/6 cases at day 9 (mean 37993), and values for control mice sham immunized with saline was 53100-315200 IFU/lung (mean 141593) at day 9.

USE - (I), (II), (III), PC and Ab are useful for preventing or treating Chlamydia infection. (I), (II) and Ab are useful for detecting Chlamydia infection, by assaying a body fluid of a mammal to be tested (claimed). (I) and (II) are useful as vaccines. The probes are used in diagnostic tests as capture or detection probes and in hybridization techniques, and primers are useful in amplification techniques for use in diagnostic methods. (I) is useful for detecting the presence of anti-Chlamydia antibodies in blood sample.

Dwg.0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-C01B; B04-C01C; B04-E01; B04-E03F; B04-E05; B04-E08; B04-F0100E; B04-F10A; B04-G01; B04-N03A0E; B04-P01A; B11-C08; B11-C08E2; B11-C08E5; B12-K04A4; B12-K04E; B12-K04F; B14-A01A; B14-S03; B14-S11B; D05-C12; D05-H07; D05-H09; D05-H12A; D05-H12D1; D05-H12E; D05-H14; D05-H17A6  
EPI: S03-E14H; S03-E14H4

L54 ANSWER 3 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-648558 [74] WPIX

DNN N2001-484574 DNC C2001-191445

TI Novel Chlamydia myosin heavy chain homolog polypeptide and polynucleotide for preventing, detecting and treating Chlamydia infections in mammals, in particular humans.

DC B04 D16 S03

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J

PA (AVET) AVENTIS PASTEUR LTD; (DUNN-I) DUNN P; (MURD-I) MURDIN A D; (OOMEN-I) OOMEN R P; (WANG-I) WANG J

CYC 95

PI WO 2001075113 A2 20011011 (200174)\* EN 83 C12N015-31  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TR TZ UG ZW  
W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK  
DM DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ  
LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD  
SE SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2001048177 A 20011015 (200209) C12N015-31

US 2002132994 A1 20020919 (200264) C07H021-02

ADT WO 2001075113 A2 WO 2001-CA461 20010404; AU 2001048177 A AU 2001-48177 20010404; US 2002132994 A1 Provisional US 2000-194475P 20000404, US 2001-824568 20010403

FDT AU 2001048177 A Based on WO 2001075113

PRAI US 2000-194475P 20000404; US 2001-824568 20010403

IC ICM C07H021-02; C12N015-31

ICS A61K039-118; A61K039-40; A61K048-00; C07H021-04;  
C07K014-295; C07K016-12; C12N015-11; C12N015-62; C12Q001-68;  
G01N033-53; G01N033-68

AB WO 200175113 A UPAB: 20021031

NOVELTY - An isolated myosin heavy chain homolog polypeptide (I) from Chlamydia, especially C. pneumoniae having a 254 residue amino acid sequence (S1), fully defined in the specification, its immunogenic

fragment comprising at least 12 consecutive amino acids or a polypeptide which has been modified without loss of immunogenicity and which has 75 % sequence identity to (I), is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:

(1) a nucleic acid molecule (II) comprising a nucleic acid sequence chosen from a sequence which encodes (I), a 965 base pair sequence (S2), fully defined in the specification, a sequence which encodes a polypeptide encoded by (S2), a sequence comprising 38 consecutive nucleotides from (II) and a sequence which encodes a polypeptide which is 75 % identical in amino acid sequence to the polypeptide encoded by (S2);

(2) a nucleic acid molecule (III) comprising a nucleic acid sequence which is anti-sense to (II);

(3) a fusion protein (IV) comprising (I) and a second polypeptide;

(4) a nucleic acid molecule (V) comprising a nucleic acid sequence which encodes (IV);

(5) a nucleic acid molecule chosen from (II), (III) and (V) operatively linked to one or more expression control sequences;

(6) a vaccine (VI), comprising:

(a) a vaccine vector and (II), (III) or (V), where each nucleic acid is capable of being expressed and the vaccine optionally comprises a second nucleic acid encoding and capable of expressing an additional polypeptide which enhances the immune response to the polypeptide expressed by the nucleic acids; or

(b) (I) or (IV), optionally comprising an additional polypeptide which enhances the immune response to (I) or (IV);

(7) a unicellular host (VII) transformed with (II), (III) or (V);

(8) an isolated nucleic acid probe of 5-100 nucleotides which hybridizes under stringent conditions to (II), its complement or anti-sense sequence;

(9) an isolated primer of 10-40 nucleotides which hybridizes under stringent conditions to (II), its complement or anti-sense sequence;

(10) a polypeptide encoded by (II) or (V);

(11) producing (I) or (IV), comprising culturing (VII);

(12) an antibody (VIII) against (I) or (IV);

(13) a pharmaceutical composition (IX) comprising (II), (III), (V), (I), (VI) or (VIII);

(14) a diagnostic kit comprising instructions for use and (II), (III), (V), (I), (IV) or (VIII);

(15) identifying (I) or (IV) which induces an immune response effective to prevent or lessen the severity of Chlamydia infection in a mammal previously immunized with polypeptide, by immunizing a mouse with (I) or (IV) and inoculating the immunized mouse with Chlamydia, where (I) or (IV) which prevents or lessens the severity of Chlamydia infection in the immunized mouse compared to a non-immunized control mouse is identified;

(16) expression plasmid pCACPNM559 containing the myosin heavy chain homolog gene, as shown in the specification;

(17) a nucleic acid molecule of sequence (S7); and

(18) a peptide having the sequence (S8).

(S7) is ATAAGAATGCGGCCGCCACCATGCATGACGCACTTCTAAGCA or GCGCGGATCCCTACAGCTGCGGACGACGACG.

(S8) is ArgValLysLysGluHisGlnLysGluLeu, LysMetAspGluPheAsnAlaLeuThr, TrpGlnGluSerGlnValAsnAlaGlnGluAsnSerThrAlaLysArgArgArgArg, AlaLeuLeuGluGlnArgThrGluLeu or IleLeuTyrTrpGlnGluSerGlnVal.

ACTIVITY - Antibiotic.

MECHANISM OF ACTION - Vaccine.

The effect of *C. pneumoniae* myosin heavy chain homolog gene in protecting mice against an intranasal challenge of *C. pneumoniae* was studied. Strain AR-39 was used in Balb/c mice as a challenge infection model to examine the capacity of Chlamydia gene products delivered as naked DNA to elicit a protective response against a sublethal *C. pneumoniae* lung infection. Protective immunity was defined as an accelerated clearance of pulmonary infection. Groups of 7-9 week old male Balb/c mice were immunized intramuscularly (i.m.) plus intranasally (i.n.) with plasmid DNA containing the *C. pneumoniae* myosin heavy chain homolog gene (pCACPNM559). Saline or the plasmid vector lacking an inserted chlamydial gene was given to groups of control animals. For i.m. immunization, alternate left and right quadriceps were injected with 100 micro g of DNA in 50 micro l of phosphate buffered saline (PBS) on three occasions at 0, 3 and 6 weeks. For i.n. immunization, anesthetized mice were aspirated 50 micro l of PBS containing 50 micro g DNA on three occasions at 0, 3 and 6 weeks. At week 8, immunized mice were inoculated i.n. with 5 multiply 105 infection forming units (IFU) of *C. pneumoniae*, strain AR39 in 100 micro l of SPG buffer to test their ability to limit the growth of a sublethal *C. pneumoniae* challenge. Lungs were taken from

mice at day 9 post-challenge and immediately homogenized. Dilutions of the homogenate were assayed for the presence of infectious Chlamydia. The results showed that the mice immunized i.n. and i.m. with pCACP559 had chlamydial lung titers less than 49000 in 5 of 6 cases at day 9 and for control mice sham immunized with saline the value was 53100-315200 IFU/lung at day 9.

USE - (I)-(V) and (VIII) are useful for detecting Chlamydia infection by assaying a body fluid of a mammal with the components. (VI) and (IX) are useful for preventing and treating Chlamydia infection (claimed), in mammals, such as humans. The nucleic acid molecules are useful for producing (I), in the construction of vaccine vectors such as poxviruses, which are further useful for preventing and/or treating Chlamydia infection and in the construction of attenuated Chlamydia strains that can over-express the nucleic acid molecules or express it in a non-toxic, mutated form. (VI) is effective in preventing and/or treating Chlamydia infection for e.g. infection caused by *C. trachomatis*, *C. psittaci*, *C. pneumoniae* or *C. pecorum*. Probes which hybridize to (II) are useful in diagnostic tests, as capture of detection probes. (VIII) is useful in affinity chromatography for purifying (I) and in prophylactic or therapeutic passive immunization methods.

Dwg.0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-C01B; B04-C01D; B04-C01G; B04-E01; B04-E03F; B04-E05; B04-E08; B04-F10A; B04-G07; B04-N03A0E; B04-P01A; B11-C08; B11-C08E2; B11-C08E5; B12-K04A4; B12-K04E; B12-K04F; B14-A01A; B14-S03; B14-S11B; D05-C12; D05-H07; D05-H09; D05-H12A; D05-H12D1; D05-H12E; D05-H14; D05-H17A6  
EPI: S03-E14H; S03-E14H4

L54 ANSWER 4 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-648557 [74] WPIX

DNN N2001-484573 DNC C2001-191444

TI Novel Chlamydia glutamate binding protein and polynucleotide for preventing, detecting and treating Chlamydia infections in mammals, in particular humans.

DC B04 D16 S03

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J

PA (AVET) AVENTIS PASTEUR LTD; (DUNN-I) DUNN P; (MURD-I) MURDIN A

D; (OOMEN-I) OOMEN R P; (WANG-I) WANG J

CYC 95

PI WO 2001075112 A2 20011011 (200174)\* EN 86 C12N015-31  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TR TZ UG ZW  
W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK  
DM DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ  
LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD  
SE SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW  
AU 2001048176 A 20011015 (200209) C12N015-31  
US 2002094965 A1 20020718 (200254) A61K048-00

ADT WO 2001075112 A2 WO 2001-CA460 20010404; AU 2001048176 A AU 2001-48176 20010404; US 2002094965 A1 Provisional US 2000-194472P 20000404, US 2001-824206 20010403

FDT AU 2001048176 A Based on WO 2001075112

PRAI US 2000-194472P 20000404; US 2001-824206 20010403

IC ICM A61K048-00; C12N015-31

ICS A61K039-118; A61K039-40; C07H021-04; C07K014-295;  
C07K016-12; C07K019-00; C12N001-21; C12N015-62; C12N015-74;  
C12Q001-68; G01N033-569

AB WO 200175112 A UPAB: 20011217

NOVELTY - An isolated glutamate binding protein (I) from Chlamydia, especially *C. pneumoniae* having a 250 residue amino acid sequence (S1), fully defined in the specification, its immunogenic fragment comprising at least 12 consecutive amino acids or a polypeptide which has been modified without loss of immunogenicity and which has 75 % sequence identity to (I), is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:

(1) a nucleic acid molecule (II) comprising a nucleic acid sequence chosen from a sequence which encodes (I), a 953 base pair sequence (S2), fully defined in the specification, a sequence which encodes a polypeptide encoded by (S2), a sequence comprising 38 consecutive nucleotides from (II) and a sequence which encodes a polypeptide which is 75 % identical in amino acid sequence to the polypeptide encoded by (S2);

(2) a nucleic acid molecule (III) comprising a nucleic acid sequence which is anti-sense to (II);

- (3) a fusion protein (IV) comprising (I) and a second polypeptide;
  - (4) a nucleic acid molecule (V) comprising a nucleic acid sequence which encodes (IV);
  - (5) a nucleic acid molecule chosen from (II), (III) and (V) operatively linked to one or more expression control sequences;
  - (6) a vaccine (VI), comprising:
    - (a) a vaccine vector and any one of the above nucleic acids, where each nucleic acid is capable of being expressed and the vaccine optionally comprises a second nucleic acid encoding and capable of expressing an additional polypeptide which enhances the immune response to the polypeptide expressed by the nucleic acid; or
    - (b) (I) or (IV), optionally comprising an additional polypeptide which enhances the immune response to (I) or (IV);
  - (7) a unicellular host (VII) transformed with (II), (III) or (V);
  - (8) an isolated nucleic acid probe of 5-100 nucleotides which hybridizes under stringent conditions to (II), its complement or anti-sense sequence;
  - (9) an isolated primer of 10-40 nucleotides which hybridizes under stringent conditions to (II), its complement or anti-sense sequence;
  - (10) a polypeptide encoded by (II) or (V);
  - (11) producing (I) or (IV), comprising culturing (VII);
  - (12) an antibody (VIII) against (I) or (IV);
  - (13) a pharmaceutical composition (IX) comprising (II), (III), (V), (I), (VI) or (VIII);
  - (14) a diagnostic kit comprising instructions for use and (II), (III), (V), (I), (IV) or (VIII);
  - (15) identifying (I) or (IV) which induces an immune response effective to prevent or lessen the severity of Chlamydia infection in a mammal previously immunized with polypeptide, by immunizing a mouse with (I) or (IV) and inoculating the immunized mouse with Chlamydia, where (I) or (IV) which prevents or lessens the severity of Chlamydia infection in the immunized mouse compared to a non-immunized control mouse is identified;
  - (16) expression plasmid pCACPNM653 containing the glutamate binding protein gene;
  - (17) a nucleic acid molecule of sequence (S7); and
  - (18) a peptide having the sequence (S8).
- (S7) is ATAAGATGCGCGCCGCCACCATGAAGATAAAATTTCTTGAAGG or GCGCCGATCCCGGAAGACGATACCGCTGTTTT. (S8) is GluAsnLeuAspAspLysLysThrGlnGly, LysThrArgArgSerGlyLysTyrAspAlaIleLysGlnArgTyrArgLeuPro, AlaLeuLeuAlaProValIleGluVal or PheLeuAsnAspLeuValSerGluIle.

ACTIVITY - Antibiotic.

MECHANISM OF ACTION - Vaccine.

The effect of *C. pneumoniae* glutamate binding protein gene in protecting mice against an intranasal challenge of *C. pneumoniae* was studied. Strain AR-39 Grayston et al (1990) Journal of Infectious Diseases 161:618-625 was used in Balb/c mice as a challenge infection model to examine the capacity of Chlamydia gene products delivered as naked DNA to elicit a protective response against a sublethal *C. pneumoniae* lung infection. Protective immunity was defined as an accelerated clearance of pulmonary infection. Groups of 7-9 week old male Balb/c mice were immunized intramuscularly (i.m.) plus intranasally (i.n.) with plasmid DNA containing the *C. pneumoniae* glutamate binding protein gene (pCACPNM653). Saline or the plasmid vector lacking an inserted chlamydial gene was given to groups of control animals. For i.m. immunization, alternate left and right quadriceps were injected with 100 micro g of DNA in 50 micro l of phosphate buffered saline (PBS) on three occasions at 0, 3 and 6 weeks. For i.n. immunization, anesthetized mice were aspirated 50 micro l of PBS containing 50 micro g DNA on three occasions at 0, 3 and 6 weeks. At week 8, immunized mice were inoculated i.n. with 5 multiply 10<sup>5</sup> infection forming units (IFU) of *C. pneumoniae*, strain AR39 in 100 micro l of SPG buffer to test their ability to limit the growth of a sublethal *C. pneumoniae* challenge. Lungs were taken from mice at day 9 post-challenge and immediately homogenized. Dilutions of the homogenate were assayed for the presence of infectious Chlamydia. The results showed that the mice immunized i.n. and i.m. with pCACPNM653 had chlamydial lung titers less than 60000 in 4 of 6 cases at day 9 and for control mice sham immunized with saline the value was 53100-315200 IFU/lung at day 9.

USE - (I)-(V) and (VIII) are useful for detecting Chlamydia infection by assaying a body fluid of a mammal with the components (claimed). (VI) and (IX) are useful for preventing or treating Chlamydia infection (claimed), in mammals, such as humans. The nucleic acid molecules are useful for producing (I), in the construction of vaccine vectors such as poxviruses, which are further useful for preventing and/or treating Chlamydia infection and in the construction of attenuated Chlamydia strains that can over-express the nucleic acid molecules or express it in

a non-toxic, mutated form. (VI) is effective in preventing and/or treating Chlamydia infection for e.g. infection caused by *C. trachomatis*, *C. psittaci*, *C. pneumoniae* or *C. pecorum*. Probes which hybridize to (II) are useful in diagnostic tests, as capture or detection probes. (VIII) is useful in affinity chromatography for purifying (I) and in prophylactic or therapeutic passive immunization methods.

Dwg.0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-C01B; B04-C01D; B04-C01G; B04-E01; B04-E03F; B04-E05; B04-E08; B04-F0100E; B04-F10A; B04-N03A0E; B11-C07A; B11-C08E2; B11-C08E5; B12-K04A4; B12-K04E; B12-K04F; B14-A01A; B14-S03; B14-S11B; D05-C12; D05-H07; D05-H09; D05-H11; D05-H12A; D05-H12D1; D05-H12E; D05-H14; D05-H17A6  
EPI: S03-E14H4

L54 ANSWER 5 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-648556 [74] WPIX

DNN N2001-484572 DNC C2001-191443

TI Novel isolated myosin heavy chain polypeptide from *Chlamydia pneumoniae* and polynucleotides encoding them, useful for treating or preventing *Chlamydia* infection in mammals.

DC B04 D16 S03

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J

PA (AVET) AVENTIS PASTEUR LTD; (DUNN-I) DUNN P; (MURD-I) MURDIN A

D; (OOME-I) OOMEN R P; (WANG-I) WANG J

CYC 95

PI WO 2001075111 A2 20011011 (200174)\* EN 83 C12N015-31

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TR TZ UG ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK  
DM DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ  
LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD  
SE SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2001048172 A 20011015 (200209) C12N015-31

US 2003100706 A1 20030529 (200337) A61K039-02 <--

ADT WO 2001075111 A2 WO 2001-CA456 20010404; AU 2001048172 A AU 2001-48172  
20010404; US 2003100706 A1 Provisional US 2000-194471P 20000404, US  
2001-824584 20010403

FDT AU 2001048172 A Based on WO 2001075111

PRAI US 2000-194471P 20000404; US 2001-824584 20010403

IC ICM A61K039-02; C12N015-31

ICS A61K039-118; A61K039-40; A61K048-00; C07H021-04;  
C07K014-195; C07K014-295; C07K016-12; C07K019-00; C12N001-21;  
C12N015-62; C12N015-74; C12Q001-68; G01N033-569

AB WO 200175111 A UPAB: 20011217

NOVELTY - An isolated myosin heavy chain polypeptide (I) from *Chlamydia pneumoniae*, comprising 168 residue amino acid sequence (S2), fully defined in the specification, an immunogenic fragment having 12 consecutive amino acids of (S2), or a sequence of (S2) or its fragment which has been modified without loss of immunogenicity and having 75 % identity to above mentioned polypeptide sequences, is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:

(1) a nucleic acid molecule (II) comprising a nucleic acid sequence which encodes (I), comprising:

(a) a 707 nucleotide sequence (S1), fully defined in specification;  
(b) a sequence which encodes a polypeptide encoded by (S1);  
(c) a sequence comprising at least 38 consecutive nucleotides of (a) or (b), or a sequence which encodes a polypeptide that is 75 % identical in amino acid sequence to polypeptide encoded by (S1);

(2) a nucleic acid molecule (III) comprising a nucleic acid sequence which is antisense to (II);

(3) a nucleic acid molecule (IV) comprising a nucleic acid sequence which encodes fusion protein that comprises a polypeptide encoded by (II) and a second polypeptide;

(4) a nucleic acid molecule ((I)-(IV)) operatively linked to one or more expression control sequences;

(5) a vaccine (V) comprising a vaccine vector and (II);

(6) a vaccine (VI) comprising a vaccine vector and at least one first nucleic acid encoding a fusion protein that comprises a polypeptide encoded by (S1), a polypeptide encoded by a nucleic acid comprising at least 38 consecutive nucleotides from (S1), a polypeptide which is 75 % identical to the amino acid sequence to the polypeptide encoded by (S1), or is (I); and

(7) a vaccine (VII) comprising (II), (III), or (V) operatively linked

to expression control sequences, as first nucleic acid and a vaccine vector, the vaccine optionally comprises a second nucleic acid encoding an additional polypeptide which enhances the immune response to the polypeptide expressed by the first;

(8) a unicellular host (VIII) transformed with a nucleic acid molecule ((II)-(IV)) operatively linked to expression control sequences;

(9) an isolated nucleic acid probe (IX) of 5-100 nucleotides which hybridizes under stringent conditions to (S1);

(10) an isolated primer (X) of 10-40 nucleotides which hybridizes under stringent conditions to (S1);

(11) a polypeptide encoded by (II), (III), (IV) or a nucleic acid molecule ((II)-(IV)) operatively linked to expression control sequences;

(12) a fusion protein (XI) comprising (I) and a second polypeptide;

(13) preparation of (I) or (XI);

(14) an antibody (XII) against (I) or (XI);

(15) a vaccine (XIII) comprising at least one first polypeptide (FP1) encoded by (S1), a polypeptide encoded by a nucleic acid comprising at least 38 consecutive nucleotides of (S1), a polypeptide which is 75 % identical to the amino acid sequence to the polypeptide encoded by (S1), or is (I), where the vaccine optionally comprises an additional polypeptide which enhances the immune response to FP1;

(16) a vaccine (XIV) comprising a fusion protein which comprises FP1 and a second polypeptide, where the vaccine optionally comprises an additional polypeptide which enhances the immune response to FP1;

(17) a vaccine (XV) comprising (I) or (XI) as the first polypeptide, and an additional polypeptide which enhances the immune response to the first polypeptide;

(18) a diagnostic kit comprising instructions for use and a component (II), (III), (V) operatively linked to expression control sequences, (I), (XI) or (XII);

(19) identifying (I) or (XI) which prevents or lessens the severity of Chlamydia infection in a mammal previously immunized with polypeptide involves immunizing a mouse with the polypeptide or fusion protein and inoculating the immunized mouse with Chlamydia;

(20) expression plasmid pCACPNM760;

(21) a nucleic acid molecule having a sequence (S7); and

(22) a peptide having a sequence (S8).

(S7) is ataagaatgcggccgccaccatggcaaaatatccactagagcc or gcgccgatcccgcttccccctgattcacg.

(S8) is LysArgArgLysGluGluGluLysThrArgLeuHisLysGluGluTrpMet, LeuArgGlnLysLysLysArgGlySer or GlnLeuSerGluGluGluGluLysVal.

ACTIVITY - Antibiotic.

MECHANISM OF ACTION - Vaccine.

Strain AR-39 was used in Balb/c mice as a challenge infection model to examine the capacity of Chlamydia gene products delivered as naked DNA to elicit a protective response against sublethal *C. pneumoniae* lung infections. Groups of 7-9 week old male Balb/c mice were immunized intramuscularly (i.m.) plus intranasally (i.n.) with plasmid DNA containing the *C. pneumoniae* myosin heavy chain gene. For i.m. immunization, alternate left and right quadriceps were injected with 100 micro g of DNA. For i.n. immunization, anesthetized mice were aspirated 50 micro l of phosphate buffered saline (PBS) containing 50 micro g DNA on three occasions at 0, 3 and 6 weeks. At week 8, immunized mice were inoculated i.n. with 5 multiply 105 infection forming units (IFU) of *C. pneumoniae*, strain AR39 in 100 micro l of SPG buffer to test their ability to limit the growth of a sublethal *C. pneumoniae* challenge. Lungs were taken from mice at day 9 post-challenge and homogenized in SPG buffer. Dilutions of the homogenate were assayed for Chlamydia by inoculation onto monolayers of susceptible cells. The inoculum was centrifuged onto the cells, then the cells were incubated for three days at 35 deg. C in the presence of 1 micro g/ml cycloheximide. After incubation the monolayers were fixed and stained using convalescent sera from rabbits infected with *C. pneumoniae*. Results showed that mice immunized with i.n. and i.m. with pCACPNM760 had chlamydial lung titers less than 40000 in 3 of 6 cases at day 9, whereas the range of values for control mice sham immunized with saline was 20800-323300 IFU/lung at day 9.

USE - (II), (III), (IV) or a nucleic acid molecule ((II), (III), (V)) operatively linked to expression control sequences, the vaccines and pharmaceutical compositions are useful for treating or preventing Chlamydia infection. (II), (III), (IV) or a nucleic acid molecule ((II)-(IV)) operatively linked to expression control sequences, (I), (XI) or (XII) is also useful for detecting Chlamydia infection. (All claimed). (I) is useful for detecting the presence of anti-Chlamydia antibodies in a biological sample. (II) is useful for producing (I), for constructing vaccine vectors, and as a vaccine agent, or in the construction of attenuated Chlamydia strains that can overexpress (II). (IX) is useful as



capture or detection probe. (IX) and (X) are useful for detecting and/or identifying the presence of Chlamydia in a biological material. (XII) is useful for purifying (I) by antibody-based affinity chromatography. (XII) can also be used in therapeutic and prophylactic passive immunization methods. (XII) used for detecting Chlamydia in biological sample.

Dwg.0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-C01B; B04-C01C; B04-C01D; B04-C01G; B04-E01; B04-E03F; B04-E05; B04-E08; B04-F0100E; B04-F10A; B04-G07; B04-N03A0E; B11-C08; B11-C08E2; B11-C08E5; B12-K04A4; B12-K04E; B12-K04F; B14-A01A; B14-S03; B14-S11B; D05-C12; D05-H07; D05-H09; D05-H12A; D05-H12D1; D05-H12E; D05-H14; D05-H17A6  
EPI: S03-E14H4

L54 ANSWER 6 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-343797 [36] WPIX

DNC C2001-106482

TI A Chlamydia polypeptide, an amino acid transporter gene, for the treatment and prevention of Chlamydia infection.

DC B04 C06 D16

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J

PA (AVET) AVENTIS PASTEUR LTD

CYC 94

PI WO 2001036457 A2 20010525 (200136)\* EN 81 C07K014-00

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TR TZ UG ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM  
DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC  
LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE  
SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2001013757 A 20010530 (200152) C07K014-00

ADT WO 2001036457 A2 WO 2000-CA1346 20001110; AU 2001013757 A AU 2001-13757 20001110

FDT AU 2001013757 A Based on WO 2001036457

PRAI US 1999-165615P 19991115

IC ICM C07K014-00

AB WO 200136457 A UPAB: 20010628

NOVELTY - A Chlamydia polypeptide which is encoded by (I), a 468 amino acid (aa) sequence, given in the specification, is new

DETAILED DESCRIPTION - The polypeptide may also be a fusion protein comprising (I) and an additional polypeptide.

INDEPENDENT CLAIMS are included for the following:

(1) a nucleic acid molecule which encodes a polypeptide, a C. pneumoniae, an amino acid transporter gene comprising:

(i) a 1607 base pair (bp) nucleic acid sequence defined in the specification;

(ii) an immunogenic fragment comprising at least 12 aa from a polypeptide encoded by (a); and

(iii) a polypeptide of (a) or (b) which has been modified to improve its immunogenicity, where the peptide is at least 75% identical in aa sequence to (a) or (b).

(2) a nucleic acid sequence selected from:

(i) 1564 bp sequence; a sequence including (a);

(ii) a sequence which encodes a polypeptide encoded by (i);

(iii) a sequence comprising at least 38 consecutive nucleotides from (i) or (ii);

(iv) a sequence which encodes a polypeptide which is at least 75% identical in aa sequence to the polypeptide encoded by (i);

(3) a nucleic acid molecule comprising a nucleic acid sequence which is antisense to (1);

(4) a nucleic acid molecule comprising a sequence encoding a fused protein which is encoding a nucleic acid (1) and an additional polypeptide;

(5) a vaccine comprising a nucleic acid of (1) and a vaccine vector where each nucleic acid is expressed as a polypeptide. The vaccine optionally comprises a second nucleic acid encoding an additional polypeptide which enhances the immune response to the polypeptide is expressed by the first;

(6) a unicellular host transformed with the nucleic acid molecule (4);

(7) a nucleic acid probe of 5 to 100 nucleotides which hybridizes under stringent conditions to the nucleic acid molecule (1) or to its homolog or complementary anti-sense sequence;

(8) a primer of 10 to 40 nucleotides which hybridizes under stringent conditions to the nucleic acid molecule of (1) or to its homolog or

complementary anti-sense sequence;

(9) the production of the polypeptide (I) comprising the culturing of (6);

(10) an antibody against polypeptide(s) of the invention;

(11) a vaccine comprising at least one first polypeptide of the invention, and optionally a second polypeptide which enhances the immune response to the first;

(12) the treatment or prevention of Chlamydia infection using:

(i) a nucleic acid of (1-4);

(ii) a vaccine of (5) or (11);

(iii) a polypeptide of the invention; and/or

(iv) (11);

(13) the detection of Chlamydia comprising the step of assaying a body fluid of a mammal with a component selected from 12 (i), (iii) and/or (11);

(14) a diagnostic kit comprising instructions for use and 12 (i), (iii) or (11);

(15) the identification of a polypeptide of the invention which induces a response effective to prevent or lessen the extent of Chlamydia infection in a mammal previously immunized with a polypeptide comprising:

(i) immunizing a mouse with the polypeptide; and

(ii) innoculating the immunized mouse with Chlamydia; where the polypeptide which prevents or lessens the severity of Chlamydia infection in the immunized mouse compared to a non-immunized control mouse

(16) expression plasmid pCABk297.

ACTIVITY - Antibacterial.

MECHANISM OF ACTION - Vaccine. Gene therapy.

USE - The polypeptide, an amino acid transporter is useful for the treatment, prevention and diagnosis of Chlamydia infection, preferably Chlamydia pneumoniae infection (claimed), in human and veterinary applications.

ADVANTAGE - A protective vaccine against Chlamydia pneumoniae is useful to prevent infection which leads to chronic bronchitis and sinusitis. There is also a correlation between infection and atherosclerosis, with epidemiological studies showing connections between the incidence of heart attack, coronary artery and carotid artery disease with organisms being detected in the fatty streaks of the coronary, carotid, peripheral arteries and aorta. The infection may also be linked with the high incidence of lower respiratory tract infections and mortality in infants and children in tropical regions of the world. The preventative vaccine reduces the need for antibiotic treatment.

Dwg.0/3

FS CPI

FA AB; DCN

MC CPI: B04-C01G; B04-E02F; B04-E03F; B04-E05; B04-E06; B04-E08;  
B04-G01; B04-N03A; B11-C08E; B12-K04A4; B12-K04F;  
B14-A01A; C04-C01G; C04-E02F; C04-E03F; C04-E05; C04-E06;  
C04-E08; C04-G01; C04-N03A; C11-C08E; C12-K04A4; C12-K04F;  
C14-A01A; D05-C11; D05-H07; D05-H11; D05-H12D; D05-H12D1;  
D05-H12E; D05-H14; D05-H17A; D05-H17B

L54 ANSWER 7 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-343796 [36] WPIX

DNC C2001-106481

TI A Chlamydia polypeptide, OppB, for the treatment and prevention of Chlamydia infection.

DC B04 C06 D16

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J

PA (AVET) AVENTIS PASTEUR LTD

CYC 94

PI WO 2001036456 A2 20010525 (200136)\* EN 75 C07K014-00

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TR TZ UG ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM  
DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC  
LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE  
SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2001013756 A 20010530 (200152) C07K014-00

ADT WO 2001036456 A2 WO 2000-CA1345 20001110; AU 2001013756 A AU 2001-13756 20001110

FDT AU 2001013756 A Based on WO 2001036456

PRAI US 1999-164918P 19991115

IC ICM C07K014-00

AB WO 200136456 A UPAB: 20010628

NOVELTY - A Chlamydia polypeptide which is encoded by (I) a 314 amino acid (aa) sequence, given in the specification, is new.

DETAILED DESCRIPTION - The polypeptide may also be a fusion protein comprising (I) and an additional polypeptide.

INDEPENDENT CLAIMS are included for the following:

(1) a nucleic acid molecule which encodes a polypeptide, a C. pneumoniae, OppB gene comprising:

(i) a 1145 base pair (bp) nucleic acid sequence defined in the specification;

(ii) an immunogenic fragment comprising at least 12 aa from a polypeptide encoded by (a); and

(iii) a polypeptide of (a) or (b) which has been modified to improve its immunogenicity, where the peptide is at least 75% identical in aa sequence to (a) or (b).

(2) a nucleic acid sequence selected from

(i) a sequence including (a);

(ii) a sequence which encodes a polypeptide encoded by (i);

(iii) a sequence comprising at least 38 consecutive nucleotides from (i) or (ii);

(iv) a sequence which encodes a polypeptide which is at least 75% identical in aa sequence to the polypeptide encoded by (i);

(3) a nucleic acid molecule comprising a nucleic acid sequence which is antisense to (1);

(4) a nucleic acid molecule comprising a sequence encoding a fused protein which is encoding a nucleic acid (1) and an additional polypeptide;

(5) a vaccine comprising a nucleic acid of (1) and a vaccine vector where each nucleic acid is expressed as a polypeptide. The vaccine optionally comprises a second nucleic acid encoding an additional polypeptide which enhances the immune response to the polypeptide is expressed by the first;

(6) a unicellular host transformed with the nucleic acid molecule (3);

(7) a nucleic acid probe of 5 to 100 nucleotides which hybridizes under stringent conditions to the nucleic acid molecule (a) or to its homolog or complementary anti-sense sequence;

(8) a primer of 10 to 40 nucleotides which hybridizes under stringent conditions to the nucleic acid molecule of (a) or to its homolog or complementary anti-sense sequence;

(9) the production of the polypeptide (I) comprising the culturing of (6);

(10) antibody against polypeptide(s) of the invention;

(11) a vaccine comprising at least one first polypeptide of the invention, and optionally a second polypeptide which enhances the immune response to the first;

(12) the treatment of Chlamydia infection using:

(i) a nucleic acid of (1-4);

(ii) a vaccine of (5) or (12);

(iii) a polypeptide of the invention; and/or

(iv) (10);

(13) the detection of Chlamydia comprising the step of assaying a body fluid of a mammal with a component selected from 12 (i), (iii) and/or (iv);

(14) a diagnostic kit comprising instructions for use and 12 (i), (iii) or (iv);

(15) the identification of a polypeptide of the invention which induces a response effective to prevent or lessen the extent of Chlamydia infection in a mammal previously immunized with a polypeptide comprising:

(i) immunizing a mouse with the polypeptide; and

(ii) innoculating the immunized mouse with Chlamydia;

(16) the expression plasmid pCAI434.

ACTIVITY - Antibacterial.

MECHANISM OF ACTION - Vaccine. Groups of 7-9 week old male Balb/c mice (n = 8-10 / group) were immunized intramuscularly (i.m) and intranasally (i.n.) with plasmid DNA containing the Chlamydia pneumonia OppB gene. Saline or the plasmid vector without the insert was given to the control animals. For i.m immunization, alternate left and right quadriceps were injected with 100 micro g of DNA in phosphate buffered saline (PBS) at three timepoints, 0, 3 and 6 weeks. For i.n immunization, anaesthetized mice were aspirated with 50 micro g of DNA in PBS at the three timepoints. A 8 weeks immunized mice were innoculated i.n with 5 X 10<sup>5</sup> IFU of Chlamydia pneumonia, strain AR39 in 100 micro l of SPG buffer. Lungs were taken from mice at day 9 post challenge, homogenised and the homogenate examined for the presence of Chlamydial inclusions. The mean bacterial load (inclusion forming units per lung) was 83378.6 for the saline control; 77000 for pCAI1021 (p = 0.7671) ; and 27450 for pCAI434 (p = 0.0028), where pCAI1021 and pCAI434 are active constructs.

USE - The polypeptide of the invention is useful for the treatment,

prevention and diagnosis of Chlamydia infection (claimed), preferably Chlamydia pneumonia infection. In human and veterinary applications.

ADVANTAGE - A protective vaccine against Chlamydia pneumonia is useful to prevent infection which leads to chronic bronchitis and sinusitis. There is also a correlation between infection and atherosclerosis, with epidemiological studies showing connections between the incidence of heart attack, coronary artery and carotid artery disease with organisms being detected in the fatty streaks of the coronary, carotid, peripheral arteries and aorta.

Dwg.0/4

FS CPI

FA AB; DCN

MC CPI: B04-C01G; B04-E02F; B04-E03F; B04-E05; B04-E06; B04-E08; B04-F0100E; B04-G01; B04-N03A; B11-C08E; B12-K04A4; B14-A01A; C04-C01G; C04-E02F; C04-E03F; C04-E05; C04-E06; C04-E08; C04-F01; C04-G01; C04-N03A; C11-C08E; C12-K04A4; C12-K04F; D05-C11; D05-H07; D05-H09; D05-H11; D05-H12A; D05-H12B; D05-H12D1; D05-H12D2; D05-H12E; D05-H14; D05-H17A; D05-H17B

L54 ANSWER 8 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-328102 [34] WPIX

DNN N2001-236077 DNC C2001-100610

TI New lpxB polypeptides useful for treating, preventing or diagnosing Chlamydia infections, particularly infections caused by Chlamydia pneumonia, e.g. bronchitis, cough, asthma.

DC B04 D16 S03

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J

PA (AVET) AVENTIS PASTEUR LTD

CYC 94

PI WO 2001021810 A1 20010329 (200134)\* EN 80 C12N015-54

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TZ UG ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM  
DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC  
LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE  
SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2000073982 A 20010424 (200141) C12N015-54

ADT WO 2001021810 A1 WO 2000-CA1085 20000915; AU 2000073982 A AU 2000-73982 20000915

FDT AU 2000073982 A Based on WO 2001021810

PRAI US 1999-154461P 19990917

IC ICM C12N015-54

ICS A61K031-711; A61K038-45; A61K039-40; C07K016-40;  
C12N009-10; C12N015-62; C12N015-85; G01N033-53

AB WO 200121810 A UPAB: 20011217

NOVELTY - A novel polypeptide (I) comprises:

(A) a fully defined sequence (IIa) of 604 amino acids (aa) given in the specification;

(B) an immunogenic fragment (IIb) comprising at least 12 consecutive aa from (IIa);

(C) (IIa) or (IIb) which has been modified to improve its immunogenicity and is at least 75% identical to (IIa) or (IIb);

(D) a sequence encoded by a sequence antisense to those in (A) - (C);

or

(E) a polypeptide of (A) - (C) and an additional polypeptide.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:

(1) a nucleic acid (II) encoding a polypeptide comprising:

(a) a fully defined sequence (IIa) of 604 amino acids (aa) given in the specification;

(b) an immunogenic fragment (IIb) comprising at least 12 consecutive aa from (IIa); or

(c) (IIa) or (IIb) which has been modified to improve its immunogenicity and is at least 75% identical to (IIa) or (IIb), is new.

(II) has a sequence of 2023 base pairs (bp) fully defined in the specification, or at least 38 consecutive nucleotides (nt) of this sequence.

(2) a nucleic acid (III) comprising a sequence antisense to (II);

(3) a nucleic acid (IV) encoding a fusion protein comprising a polypeptide encoded by (II) and an additional polypeptide;

(4) vaccines (V) comprising:

(a) at least one (II) a vaccine vector, and optionally a second nucleic acid encoding an additional polypeptide that enhances the immune response to the polypeptide expressed by the first nucleic acid; or

(b) at least one (I) and optionally a second polypeptide that enhances the immune response to the first polypeptide;

- (5) a unicellular host (VI) transformed with (II);
- (6) a nucleic acid probe (VIIa) of 5-100 nt or a primer (VIIb) of 10-40 nt, which hybridizes under stringent conditions to a 2023-bp sequence, or its homologue, complement, or antisense sequence;
- (7) producing (I) by culturing (VI);
- (8) an antibody (VIII) immunospecific for (I);
- (9) preventing or treating (M1) Chlamydia infection using the nucleic acids, vaccines, pharmaceutical composition, polypeptides or antibodies of the invention;
- (10) detecting (M2) Chlamydia infection by assaying a body fluid of a mammal with the nucleic acids, polypeptides or antibody of the invention;
- (11) a diagnostic kit (IX) comprising instructions for use and the nucleic acids, polypeptides or antibodies of the invention;
- (12) identifying (M3) a (I) that induces immune response effective to prevent or lessen the severity of Chlamydia infection in a mammal previously immunized with the polypeptide by:
  - (a) immunizing a mouse with the polypeptide; and
  - (b) inoculating the immunized mouse with Chlamydia, where the polypeptide prevents or lessens the severity of Chlamydia infection in the immunized mouse compared to a non-immunized;
- (13) expression plasmid (X), pCABk1043;
- (14) a nucleic acid (XI) with a 45 or 34 bp sequence given in the specification; and
- (15) polypeptide lpxB (XII) from Chlamydia.

## ACTIVITY - Antimicrobial.

MECHANISM OF ACTION - Vaccine. 7-9 week old male Balb/c mice were immunized intramuscularly plus intranasally with plasmid DNA containing the coding sequence of C. pneumonia lpxB or plasmid vector lacking an inserted chlamydial gene. Immunization was given at 0, 3 and 6 weeks, and at week 8, mice were inoculated with 5 multiply 10<sup>5</sup> IFU of C. pneumoniae strain AR39. 9 days post-challenge, lungs were taken and homogenized in SPG buffer. Dilutions of homogenate were assayed for the presence of infectious chlamydia by inoculation onto monolayers of susceptible cells. Cells were incubated for 3 days, and monolayers were fixed with formalin and methanol, and immunoperoxidase stained for the presence of chlamydial inclusions using convalescent sera from rabbits infected with C. pneumoniae. Mice immunized with pCABk1043 had chlamydial lung titers less than 37,000 in 5 of 6 cases at day 9, while the range of values for the controls was 13,600-458,100 IFU/lung.

USE - (I), (II), (V) and (VIII) are useful as pharmaceutical compositions (claimed). The nucleic acids encoding the Chlamydia lpxB polypeptides are useful as a vaccine in preventing, treating or diagnosing Chlamydia infections, particularly those caused by C. pneumoniae, including respiratory diseases, e.g. cough, sore throat, bronchitis, asthma. The polynucleotides, including DNA or RNA may be used in producing the encoded polypeptide in a recombinant host system, in the construction of vaccine vectors such as pox viruses, as vaccine agent, and in constructing attenuated Chlamydia strains that can over-express a polynucleotide or express it in a non-toxic mutated form. The polypeptides may also be used as diagnostic reagent for detecting the presence of anti-Chlamydia antibodies, and in the preparation of a medicament for treating or preventing Chlamydia infection.

Dwg.0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-C01G; B04-E03F; B04-E05; B04-E06; B04-E08; B04-F01; B04-G07;  
 B04-N03A; B11-C07A; B11-C08E2; B11-C08E5; B12-K04A4; B12-K04F;  
 B14-A01A; B14-S11B; D05-C11; D05-H04; D05-H07;  
 D05-H08; D05-H11; D05-H12A; D05-H12C; D05-H12D1; D05-H12D2; D05-H12E;  
 D05-H14; D05-H17A5; D05-H17C

EPI: S03-E14H4

L54 ANSWER 9 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-328101 [34] WPIX

DNN N2001-236076 DNC C2001-100609

TI New general secretion pathway protein E polypeptides and nucleic acids encoding the polypeptides useful for treating, preventing or diagnosing Chlamydia infections, particularly infections caused by Chlamydia pneumoniae.

DC B04 D16 S03

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J

PA (AVET) AVENTIS PASTEUR LTD

CYC 94

PI WO 2001021805 A1 20010329 (200134)\* EN 79 C12N015-31

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
 NL OA PT SD SE SL SZ TZ UG ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM  
 DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC  
 LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE  
 SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2000073986 A 20010424 (200141) C12N015-31

ADT WO 2001021805 A1 WO 2000-CA1089 20000915; AU 2000073986 A AU 2000-73986  
 20000915

FDT AU 2000073986 A Based on WO 2001021805

PRAI US 1999-154595P 19990917

IC ICM C12N015-31

ICS A61K031-711; A61K039-118; A61K039-40;  
 C07K014-295; C07K016-12; C12N015-62; C12N015-85; G01N033-53

AB WO 200121805 A UPAB: 20010620

NOVELTY - A nucleic acid (I) encoding a polypeptide comprising:

(a) a fully defined sequence of 496 amino acids given in the  
 specification;

(b) an immunogenic fragment comprising at least 12 consecutive amino  
 acids from (a); or

(c) (a) or (b) which has been modified to improve its immunogenicity  
 and which is at least 75% identical to (a) or (b), is new.

DETAILED DESCRIPTION - The nucleic acid (I) has a sequence of 1691 bp  
 fully defined in the specification, or has at least 38 consecutive  
 nucleotides of this sequence.

INDEPENDENT CLAIMS are also included for the following:

(1) a nucleic acid comprising a sequence antisense to (I);  
 (2) a nucleic acid encoding a fusion protein comprising a polypeptide  
 encoded by (I) and an additional polypeptide;

(3) vaccines comprising at least one first nucleic acid expressed as  
 a polypeptide, a vaccine vector, and optionally a second nucleic acid  
 encoding an additional polypeptide that enhances the immune response to  
 the polypeptide expressed by the first nucleic acid;

(4) a unicellular host transformed with the nucleic acid;

(5) a nucleic acid probe of 5-100 nucleotides or a primer of 10-40  
 nucleotides, which hybridizes under stringent conditions to a 1691-bp  
 sequence, or its homologue, complement, or antisense sequence;

(6) a polypeptide encoded by the nucleic acids;

(7) vaccines comprising at least one first polypeptide and optionally  
 a second polypeptide that enhances the immune response to the first  
 polypeptide;

(8) a fusion polypeptide comprising a polypeptide of (6) and an  
 additional polypeptide;

(9) a method of producing a polypeptide of (6) by culturing a  
 unicellular host of (4);

(10) an antibody against the polypeptide of (6);

(11) pharmaceutical compositions comprising a polypeptide or an  
 antibody;

(12) a method of preventing or treating Chlamydia infection using the  
 above nucleic acids, vaccines, pharmaceutical composition, polypeptides or  
 antibodies;

(13) a method of detecting Chlamydia infection by assaying a body  
 fluid of a mammal with the above nucleic acids, polypeptides or antibody;

(14) a diagnostic kit comprising instructions for use and the above  
 nucleic acids, polypeptides or antibodies;

(15) a method for identifying a polypeptide that induces immune  
 response effective to prevent or lessen the severity of Chlamydia  
 infection in a mammal previously immunized with the polypeptide by:

(a) immunizing a mouse with the polypeptide; and

(b) inoculating the immunized mouse with Chlamydia, where the  
 polypeptide prevents or lessens the severity of Chlamydia infection in the  
 immunized mouse compared to a non-immunized;

(16) expression plasmid pCAI284;

(17) the nucleic acid

(I) ATAAGAATGC GGCCGCCACC ATGGCTGCTA GTATTTTAT;

(II) CCCCAAGCTT CATCACAGCG CTGGTAAC.

(18) having a 39 or 29 bp sequence given in the specification; and

(19) general secretion pathway protein E from Chlamydia, preferably

C. pneumoniae.

ACTIVITY - Antimicrobial.

MECHANISM OF ACTION - Vaccine. 7-9 week old male Balb/c mice were  
 immunized intramuscularly plus intranasally with plasmid DNA containing  
 the coding sequence of C. pneumoniae general secretion pathway protein E  
 or plasmid vector lacking an inserted chlamydial gene. Immunization was  
 given at 0, 3 and 6 weeks, and at week 8, mice were inoculated with 5  
 multiply 105 IFU of C. pneumoniae strain AR39. 9 days post-challenge,  
 lungs were taken and homogenized in SPG buffer. Dilutions of homogenate  
 were assayed for the presence of infectious chlamydia by inoculation onto

monolayers of susceptible cells. Cells were incubated for 3 days, and monolayers were fixed with formalin and methanol, and immunoperoxidase stained for the presence of chlamydial inclusions using convalescent sera from rabbits infected with *C. pneumoniae*. Mice immunized with pCAI284 had chlamydial lung titers less than 50,000 in 5 of 6 cases at day 9, while the range of values for the controls was 18,200-247,100 IFU/lung.

USE - The nucleic acids encoding the Chlamydia general secretion pathway protein E polypeptides are useful as a vaccine in preventing, treating or diagnosing Chlamydia infections, particularly those caused by *C. pneumoniae*, including respiratory diseases, e.g. cough, sore throat, bronchitis, asthma. The polynucleotides, including DNA or RNA may be used in producing the encoded polypeptide in a recombinant host system, in the construction of vaccine vectors such as poxviruses, as vaccine agent, and in constructing attenuated Chlamydia strains that can over-express a polynucleotide or express it in a non-toxic mutated form. The polypeptides may also be used as diagnostic reagent for detecting the presence of anti-Chlamydia antibodies, and in the preparation of a medicament for treating or preventing Chlamydia infection.

Dwg.0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-C01G; B04-E02F; B04-E03F; B04-E04; B04-E06; B04-E08; B04-F01; B04-F02; B04-G01; B04-G21; B04-G22; B11-C07; B11-C08; B11-C08E5; B12-K04A; B12-K04F; B14-A01A; B14-K01; B14-S11; D05-C07; D05-C11; D05-H07; D05-H08; D05-H09; D05-H12A; D05-H12B; D05-H12C; D05-H12D1; D05-H12D2; D05-H12D6; D05-H12E; D05-H14B2; D05-H17A1; D05-H17B; D05-H17B6; D05-H17C  
EPI: S03-E14H4

L54 ANSWER 10 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-316102 [33] WPIX

DNN N2001-227243 DNC C2001-097308

TI New Npt2cp (ADP/ATP translocase) polypeptides and nucleic acids encoding the polypeptides useful for treating, preventing or diagnosing Chlamydia infections, particularly infections caused by *Chlamydia pneumoniae*.

DC B04 D16 S03

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J

PA (AVET) AVENTIS PASTEUR LTD

CYC 95

PI WO 2001021803 A1 20010329 (200133)\* EN 79 C12N015-31  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TZ UG ZW  
W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM  
DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC  
LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE  
SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW  
AU 2000073984 A 20010424 (200141) C12N015-31  
EP 1220924 A1 20020710 (200253) EN C12N015-31  
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

ADT WO 2001021803 A1 WO 2000-CA1087 20000915; AU 2000073984 A AU 2000-73984 20000915; EP 1220924 A1 EP 2000-962124 20000915; WO 2000-CA1087 20000915

FDT AU 2000073984 A Based on WO 2001021803; EP 1220924 A1 Based on WO 2001021803

PRAI US 1999-154326P 19990917

IC ICM C12N015-31

ICS A61K031-711; A61K039-118; A61K039-40;

C07K014-295; C07K016-12; C12N015-62; C12N015-85; G01N033-53

AB WO 200121803 A UPAB: 20010615

NOVELTY - A nucleic acid (I) encoding a polypeptide comprising:

(a) a fully defined sequence of 540 amino acids given in the specification;

(b) an immunogenic fragment comprising at least 12 consecutive amino acids from (a); or

(c) (a) or (b) which has been modified to improve its immunogenicity and which is at least 75% identical to (a) or (b), is new.

DETAILED DESCRIPTION - The nucleic acid has 1823 bp sequence given in the specification, or comprises at least 38 consecutive nucleotides of this sequence.

INDEPENDENT CLAIMS are also included for the following:

(1) a nucleic acid comprising a sequence antisense to (I);

(2) a nucleic acid encoding a fusion protein comprising a polypeptide encoded by (I) and an additional polypeptide;

(3) vaccines comprising at least one first nucleic acid expressed as a polypeptide, a vaccine vector, and optionally a second nucleic acid encoding an additional polypeptide that enhances the immune response to

the polypeptide expressed by the first nucleic acid;

(4) a unicellular host transformed with the nucleic acid;

(5) a nucleic acid probe of 5-100 nucleotides or a primer of 10-40 nucleotides, which hybridizes under stringent conditions to an 1823-bp sequence, or to its homologue, complement, or antisense sequence;

(6) a polypeptide encoded by the nucleic acids;

(7) a fusion polypeptide comprising a polypeptide of (6) and an additional polypeptide;

(8) a method of producing a polypeptide of (6) by culturing a unicellular host of (4);

(9) an antibody against the polypeptide of (6);

(10) pharmaceutical compositions comprising a polypeptide or an antibody;

(11) a method of preventing or treating Chlamydia infection using the above nucleic acids, vaccines, pharmaceutical composition, polypeptides or antibodies;

(12) a method of detecting Chlamydia infection by assaying a body fluid of a mammal with the above nucleic acids, polypeptides or antibody;

(13) a diagnostic kit comprising instructions for use and the above nucleic acids, polypeptides or antibodies;

(14) a method for identifying a polypeptide that induces immune response effective to prevent or lessen the severity of Chlamydia infection in a mammal previously immunized with the polypeptide by: (a) immunizing a mouse with the polypeptide; and (b) inoculating the immunized mouse with Chlamydia, where the polypeptide prevents or lessens the severity of Chlamydia infection in the immunized mouse compared to a non-immunized;

(15) expression plasmid pCABk663;

(16) a nucleic acid having a 42 or 33 bp sequence given in the specification; and (17) Npt2cp (ADP/ATP translocase) from Chlamydia pneumoniae.

#### ACTIVITY - Antimicrobial.

MECHANISM OF ACTION - Vaccine. 7-9 week old male Balb/c mice were immunized intramuscularly plus intranasally with plasmid containing the coding sequence of Chlamydia pneumoniae Npt2cp (ADP/ATP translocase) or plasmid vector lacking an inserted chlamydial gene. Immunization was given at 0, 3 and 6 weeks, and at week 8, mice were inoculated with 5 multiply 105 IFU of C. pneumoniae strain AR39. 9 days post-challenge, lungs were taken and homogenized in SPG buffer. Dilutions of homogenate were assayed for the presence of infectious chlamydia by inoculation onto monolayers of susceptible cells. Cells were incubated for 3 days, and monolayers were fixed with formalin and methanol, and immunoperoxidase stained for the presence of chlamydial inclusions using convalescent sera from rabbits infected with C. pneumoniae. Mice immunized with pCABk663 had chlamydial lung titers less than 36,000 in 5 of 6 cases at day 9, while the range of values for the controls was 13,600-458,100 IFU/lung.

USE - The nucleic acids encoding the Chlamydia Npt2cp (ADP/ATP translocase) polypeptides are useful as a vaccine in preventing, treating or diagnosing Chlamydia infections, particularly those caused by C. pneumoniae, including respiratory diseases, e.g. cough, sore throat, bronchitis, asthma. The polynucleotides, including DNA or RNA may be used in producing the encoded polypeptide in a recombinant host system, in the construction of vaccine vectors such as poxviruses, as vaccine agent, and in constructing attenuated Chlamydia strains that can over-express a polynucleotide or express it in a non-toxic mutated form. The polypeptides may also be used as diagnostic reagent for detecting the presence of anti-Chlamydia antibodies, and in the preparation of a medicament for treating or preventing Chlamydia infection.

Dwg.0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-C01G; B04-E02; B04-E02E; B04-E03E; B04-E06; B04-E08; B04-F01;

B04-G01; B04-G03; B04-L01; B11-C08E; B12-K04A4;

B14-A01A; D05-C03; D05-C07; D05-H07; D05-H09; D05-H11;

D05-H12A; D05-H12B; D05-H12C; D05-H12D2; D05-H12E; D05-H14; D05-H17A;

D05-H17B; D05-H17C

EPI: S03-E14H4

L54 ANSWER 11 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-257992 [26] WPIX

DNN N2001-183971 DNC C2001-077792

TI Novel Chlamydia pneumoniae lpdA protein and polynucleotides encoding them useful as component of vaccines for treating Chlamydia infections, and for detecting Chlamydia infection in the body fluid of a mammal.

DC B04 D16 S03

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J



PA (AVET) AVENTIS PASTEUR LTD

CYC 94

PI WO 2001021802 A1 20010329 (200126)\* EN 78 C12N015-31

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TZ UG ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM  
DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC  
LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE  
SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2000073983 A 20010424 (200141) C12N015-31

ADT WO 2001021802 A1 WO 2000-CA1086 20000915; AU 2000073983 A AU 2000-73983  
20000915

FDT AU 2000073983 A Based on WO 2001021802

PRAI US 1999-154325P 19990917

IC ICM C12N015-31

ICS A61K039-118; C07K014-295; C07K016-12; C07K019-00;

C12N005-10; C12N015-62; C12N015-63; C12Q001-68; G01N033-53

AB WO 200121802 A UPAB: 20010515

NOVELTY - A polypeptide (I) which is (i) a polypeptide having fully defined *Chlamydia pneumoniae* lpdA protein sequence of 461 amino acids (S2) given in the specification, (ii) an immunogenic fragment of (S2) comprising 12 consecutive amino acids or (iii) polypeptide of (i) or (ii) which has been modified to improve its immunogenicity, and having 75% identity to amino acid sequence of (i) or (ii), is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:

(1) a nucleic acid molecule (II) comprising a nucleic acid sequence which encodes (I);

(2) a nucleic acid molecule (III) comprising a nucleic acid sequence which is antisense to (II);

(3) a nucleic acid molecule (IV) comprising a nucleic acid sequence which encodes a fusion protein, comprising (I) encoded by (II) and an additional polypeptide;

(4) a vaccine (V) comprising (I), (II) or (IV) and a vaccine vector, where each nucleic acid is expressed as a polypeptide. The vaccine optionally comprising a second nucleic acid encoding an additional polypeptide which enhances the immune response to the polypeptide expressed by the above mentioned nucleic acid;

(5) a pharmaceutical composition comprising (II), (II) or (IV) and a carrier;

(6) a unicellular host transformed with (II), (III) or (IV) which is operatively linked to one or more expression control sequences;

(7) a nucleic acid probe (VI) of 5 to 100 nucleotides which hybridizes under stringent conditions to a fully defined *C.pneumoniae* lpdA gene sequence of 1586 nucleotides (S1) as given in the specification, its homolog or complementary or anti-sense sequence;

(8) a primer of 10 to 40 nucleotides which hybridizes under stringent conditions to (S1), or to a homolog or complementary or anti-sense sequence of the nucleic acid molecule;

(9) a polypeptide encoded by (II) or (IV);

(10) a fusion polypeptide (VII) comprising (I) and an additional polypeptide;

(11) preparation of (I);

(12) an antibody (VIII) against (I);

(13) a vaccine (IX) comprising (I) or (VII), and a carrier and optionally comprising a second polypeptide which enhances the immune response to (I);

(14) a pharmaceutical composition comprising (I), (VII), (IX) or (VIII) and a carrier;

(15) a diagnostic kit comprising instructions for use and (II), (III), (IV), (I), (VII) or (VIII);

(16) identifying (I) or (VII) which induces an immune response effective to prevent or lessen the severity of *Chlamydia* infection in a mammal previously immunized with polypeptide involves immunizing a mouse with (I) or (VII) and inoculating the immunized mouse with *Chlamydia*;

(17) expression plasmid pCABk892;

(18) a nucleic acid molecule having a fully defined sequence of  
ataagaatgcggccgccaccatgacccaagaatttgattgtgttg (S3) or  
cggggtaccgtgacttaggaggggaagtgttaaag (S4); and

(19) lpdA protein from *C.pneumoniae*.

ACTIVITY - Antibacterial. The biological activity of (I) was tested in mice. Groups of 7 to 9 week old male Balb/c mice (6 to 10 per group) were immunized intramuscularly (i.m.) plus intranasally (i.n.) with plasmid DNA containing the coding sequence of *C.pneumoniae* lpdA. At week 8, immunized mice were inoculated i.n. with 5 multiply 105 IFU of *C.pneumoniae*, strain AR39 in 100 µl of SPG buffer to test their ability

to limit the growth of a sublethal *C.pneumoniae* challenge. Lungs were taken from the mice at day 9 post-challenge and immediately homogenized in SPG buffer (7.5% sucrose, 5 mM glutamate, 12.5 mM phosphate pH 7.5). Dilutions of the homogenate were assayed for the presence of infectious *Chlamydia* by inoculation onto monolayers of susceptible cells. The inoculum was centrifuged onto the cells, then the cells were incubated for three days at 35 deg. C in the presence of 1  $\mu$ g/ml cycloheximide. After incubation the monolayers were fixed with formalin and methanol then immunoperoxidase stained for the presence of chlamydial inclusions using convalescent sera from rabbits infected with *C.pneumoniae* and metal-enhanced DAB as a peroxidase substrate. Results showed that mice immunized i.n. and i.m. with pCABk892 had chlamydial lung titers less than 25,000 in 6 of 6 cases at day 9 whereas the range of values for control mice sham immunized with saline was 13,600-458,100 IFU/lung at day 9.

#### MECHANISM OF ACTION - Vaccine; Gene therapy.

USE - (II), (III), (IV), (V), (VIII) or (IX) or the pharmaceutical compositions as described above are useful for preventing or treating *Chlamydia* (*C.trachomatis*, *C.psittaci*, *C.pneumonia* or *C.pecorum*) infection. (I), (II), (III), (IV), (VII) or (VIII) is useful as diagnostic reagents for detecting *Chlamydia* infection which involves assaying a body fluid of a mammal to be tested for the above mentioned components. (II) is useful for producing (I) (claimed). The vaccine vectors, (I), (II), (VIII) are useful in the preparation of a medicament for preventing and/or treating *Chlamydia* infection. (VI) is useful in diagnostic tests as capture or detection probes. (VI) is thus useful as an agent for detecting and/or identifying presence of *Chlamydia* in the biological material. The primers derived from (II) are also useful for detecting and/or identifying *Chlamydia* in the biological material. (VIII) is also useful as a reagent for purifying (I) from a biological sample which involves carrying out antibody-based affinity chromatography with the biological sample.

Dwg.0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-C01G; B04-E03F; B04-E04; B04-E05; B04-E06; B04-E08; B04-F01; B04-F10A; B04-G09; B04-N0300E; B04-N03A; B11-C07A; B11-C08E5; B12-K04A4; B12-K04F; B14-A01A; B14-S03; B14-S11B; D05-C11; D05-H09; D05-H11; D05-H12A; D05-H12C; D05-H12D1; D05-H12D2; D05-H12D5; D05-H12E; D05-H14A1; D05-H17A6; D05-H17C1; D05-H18 EPI: S03-E14H4

L54 ANSWER 12 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2001-168447 [17] WPIX

DNC C2001-050284

TI Novel multivalent immunogenic composition for conferring protection against infection caused by *Hameophilus influenzae* and *Moraxella catarrhalis* comprises four antigens derived from each of the two microorganisms.

DC B04 D16

IN KLEIN, M H; LOOSMORE, S M; SASAKI, K; YANG, Y

PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD

CYC 95

PI WO 2001005424 A2 20010125 (200117)\* EN 58 A61K039-00 <--  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TZ UG ZW  
W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM  
DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC  
LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE  
SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW  
AU 2000059586 A 20010205 (200128) A61K039-00 <--  
EP 1200122 A2 20020502 (200236) EN A61K039-116 <--  
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

US 6391313 B1 20020521 (200239) A61K039-116 <--

AU 767096 B 20031030 (200382) A61K039-00 <--

NZ 516819 A 20031219 (200404) A61K039-00 <--

ADT WO 2001005424 A2 WO 2000-CA811 20000711; AU 2000059586 A AU 2000-59586 20000711; EP 1200122 A2 EP 2000-945494 20000711; WO 2000-CA811 20000711; US 6391313 B1 US 1999-353617 19990715; AU 767096 B AU 2000-59586 20000711; NZ 516819 A NZ 2000-516819 20000711; WO 2000-CA811 20000711

FDT AU 2000059586 A Based on WO 2001005424; EP 1200122 A2 Based on WO 2001005424; AU 767096 B Previous Publ. AU 2000059586, Based on WO 2001005424; NZ 516819 A Based on WO 2001005424

PRAI US 1999-353617 19990715

IC ICM A61K039-00; A61K039-116

ICS A61P031-04

AB WO 200105424 A UPAB: 20010328

NOVELTY - A multivalent immunogenic composition (I) for conferring protection in a host against disease caused by both *Hameophilus influenzae* (HI) and *Moraxella catarrhalis* (MC) comprising four different antigens, of which at least one antigen is from HI and one antigen is from MC, is new. Additionally three of the antigens of (I) are adhesins, and one is from MC.

ACTIVITY - Auditory; antibacterial.

MECHANISM OF ACTION - Vaccine.

Groups of five BALB/C mice were immunized subcutaneously on days 1, 29 and 43 with one of the mouse H91A Hin47 + rHMW + rHia + r200 kDa vaccines. Blood samples were taken on days 0, 14, 28, 42 and 56. Groups of five Hartley outbred guinea pigs were immunized intramuscularly on days 1, 29 and 43 with the vaccine as described above. Blood samples were taken on days 0, 14, 28, 42 and 56. Anti-H91A Hin47, anti-rHMW, anti-rHia and anti-r200 kDa IgG antibody titers were determined by antigen specific enzyme linked immunosorbant assays (ELISAs). The results of the immunogenicity studies showed that the final bleed sera obtained from mice immunized with 0.3 mu g, or 3.0 mu g each of H91A Hin47 + rHMW + rHia with 0, 0.3, 1.0, 3.0 or 10.0 mu g of added r200 kDa, all had high antibody titers to H91A Hin47 component. The final bleed sera obtained from the mice immunized with 3.0 mu g each of H91A Hin47 + rHMW + rHia with 0, 0.3, 1.0, 3.0 or 10.0 mu g of added r200 kDa, all had high titer antibodies to the rHMW apparent enhancing or inhibiting effect on the anti-rHMW response with the addition of the r200 kDa component. Mice immunized with 0.3 mu g each of H91A Hin 47 + HMW + rHia with 0, 0.3, 1.0, 3.0 or 10.0 mu g of added r200 kDa, all had high titer antibodies to the rHia component. There was no apparent enhancing or inhibiting effect on the anti-rHia response with the addition of the r200 kDa component. The final bleed sera obtained from guinea pigs immunized with 25 mu g or 50 mu g each of H91A Hin47 + rHMW + rHia with 0, 25, 50 or 100 mu g of added r200 kDa, all had high titer antibodies to the H91A Hin47 component. Also final bleed sera obtained from guinea pigs immunized with 25 mu g or 50 mu g each of H91A Hin47 + rHMW + rHia with 0, 25, 50 or 100 mu g of added r200 kDa, all had titer antibodies to the rHMW component. There was no apparent enhancing or inhibiting effect on the anti-rHMW response upon the addition of the r200 kDa antigen.

USE - (I) is useful for immunizing a host against infection caused by both HI and MC including otitis media (claimed).

ADVANTAGE - The multivalent vaccine can confer protection against encapsulated and unencapsulated HI and MC diseased in a safe and efficient manner.

Dwg.0/14

FS CPI

FA AB; DCN

MC CPI: B04-B04C1; B14-A01; B14-A01A; B14-N02;  
B14-S11B; D05-C02; D05-H07; D05-H12F

L54 ANSWER 13 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2000-687542 [67] WPIX

DNC C2000-209327

TI Nucleic acids encoding a 76 kDa protein from *Chlamydia pneumoniae*, useful for vaccinating against *Chlamydia* infections.

DC B04 D16

IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J

PA (AVET) AVENTIS PASTEUR LTD; (DUNN-I) DUNN P; (MURD-I) MURDIN A

D; (OOMEN-I) OOMEN R P; (WANG-I) WANG J

CYC 93

PI WO 2000066739 A2 20001109 (200067)\* EN 90 C12N015-31

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ TZ UG ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM DZ  
EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK  
LR LS LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI  
SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2000043885 A 20001117 (200111) C12N015-31

EP 1177301 A2 20020206 (200218) EN C12N015-31

R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

JP 2002542827 W 20021217 (200312) 110 C12N015-09

US 2003095973 A1 20030522 (200336) A61K039-40 <--

NZ 515674 A 20031219 (200404) C12N015-31

US 2004086525 A1 20040506 (200430) C07H021-04

ADT WO 2000066739 A2 WO 2000-CA511 20000503; AU 2000043885 A AU 2000-43885  
20000503; EP 1177301 A2 EP 2000-925004 20000503; WO 2000-CA511 20000503;  
JP 2002542827 W JP 2000-615762 20000503; WO 2000-CA511 20000503; US

2003095973 A1 Provisional US 1999-132270P 19990503, Provisional US 1999-141276P 19990630, US 2000-564479 20000503; NZ 515674 A NZ 2000-515674 20000503, WO 2000-CA511 20000503; US 2004086525 A1 Provisional US 1999-132270P 19990503, Provisional US 1999-141276P 19990630, Cont of US 2000-564479 20000503, US 2003-608559 20030630

FDT AU 2000043885 A Based on WO 2000066739; EP 1177301 A2 Based on WO 2000066739; JP 2002542827 W Based on WO 2000066739; NZ 515674 A Based on WO 2000066739

PRAI US 1999-141276P 19990630; US 1999-132270P 19990503; US 2000-564479 20000503; US 2003-608559 20030630

IC ICM A61K039-40; C07H021-04; C12N015-09; C12N015-31

ICS A61K031-70; A61K039-00; A61K039-02; A61K039-118; A61K039-38; A61K039-39; A61K039-395; A61K048-00; A61P009-10; A61P011-00; A61P011-02; A61P011-06; A61P031-04; C07K014-295; C07K016-12; C07K019-00; C12N001-15; C12N001-19; C12N001-21; C12N005-10; C12N015-11; C12N015-62; C12N015-85; C12P021-02; C12Q001-68; G01N033-53; G01N033-566; G01N033-569

AB WO 2000066739 A UPAB: 20001223

NOVELTY - Nucleic acids (NAM1) encoding a 76 kDa protein (PEP1) from Chlamydia pneumoniae, is new. NAM1 and PEP1 have defined nucleotide and amino acid sequences ((I)-(VIII)) given in the specification.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are included for the following:

(1) a nucleic acid molecule (NAM1) comprising a nucleic acid sequence which encodes a polypeptide selected from:

(a) one of 3 defined amino acid sequences ((I)-(III)) given in the specification;

(b) a immunogenic fragment comprising at least 12 consecutive amino acids from (I)-(III); and

(c) the polypeptides of (a) and (b) which have been modified to improve their immunogenicity (the modified polypeptide is at least 75% identical in sequence to the corresponding polypeptides of (a) and (b);

(2) a nucleic acid molecule (I') comprising a sequence antisense to NAM1;

(3) a nucleic acid molecule (NAM2) which encodes a fusion protein that comprises a polypeptide encoded by NAM1 and an additional polypeptide;

(4) a vaccine (VAC1) comprising NAM1 and/or NAM2 and a vaccine vector (each nucleic acid molecule is expressed as a polypeptide and the vaccine may comprise additional nucleic acids encoding other polypeptides which enhance the immune response to the polypeptide expressed from NAM1 and/or NAM2);

(5) a unicellular host (UCH) transformed with NAM1 and NAM2 operatively linked to at least 1 expression control sequence;

(6) a nucleic acid probe of 5-100 nucleotides which hybridizes under stringent conditions to (I) (or homolog, complementary or antisense sequences of (I));

(7) a polypeptide (PEP1) encoded by NAM1 or NAM2;

(8) a fusion polypeptide (PEP2) comprising PEP1 and an additional polypeptide;

(9) a method for producing PEP1 comprising culturing UCH;

(10) an antibody (Ab) against PEP1 and/or PEP2;

(11) a vaccine (VAC2) comprising PEP1 and/or PEP2 (the vaccine may comprise additional polypeptides which enhance the immune response to PEP1 and/or PEP2);

(12) a diagnostic kit comprising NAM1, NAM2, PEP1, PEP2 and/or Ab and instructions for use;

(13) a method for identifying polypeptides (either PEP1 or PEP) which induce an immune response that prevents or reduces the severity of Chlamydia infections in mammals previously immunized with the polypeptide, comprising:

(a) immunizing a mouse with the polypeptide; and

(b) inoculating the immunized mouse with Chlamydia (the polypeptide which prevents or lessens the severity of the Chlamydia infection in the immunized mouse compared to a non-immunized control mouse is identified);

(14) an expression plasmid selected from pCACPNM555a, pCAI555, pCAD76kDa; and

(15) an isolated 76 kDa protein (PEP3) from Chlamydia.

ACTIVITY - Bactericidal.

MECHANISM OF ACTION - Vaccine.

Mouse immunized intranasally and intramuscularly with pCACPNM555a had Chlamydial lung titers less than 30000 IFU/lung in 5 of 6 cases at day 9 the range of values for control mice sham immunized with saline were 20800-323300 IFU/lung.

USE - NAM1, NAM2, PEP1, PEP2, VAC1, VAC2 and Ab may be used as

antigens for preventing and treating Chlamydia infection by vaccination. NAM1, NAM2, PEP1, PEP2 and Ab may also be used to detect Chlamydia infection in mammals by using them to assay body fluid (claimed) (e.g. in DNA hybridization assays and immunoassays).  
Dwg.0/9

FS  
CPI

FA  
AB; DCN

MC  
CPI: B04-B04C1; B04-C01; B04-E03F; B04-E04; B04-E05; B04-E06;  
B04-E08; B04-F0100E; B04-F10A; B04-G07; B04-N03A0E; B11-A; B11-C07A;  
B11-C08E; B11-C09; B12-K04A4; B12-K04E; B12-K04F; B14-A01A;  
B14-S11B; D05-A01A4; D05-A01B; D05-C12; D05-H04; D05-H07;  
D05-H08; D05-H09; D05-H11; D05-H12A; D05-H12C; D05-H12D; D05-H12E;  
D05-H14; D05-H17A5; D05-H17C; D05-H18

L54 ANSWER 14 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2000-431500 [37] WPIX

DNC C2000-131168

TI New immunogenic composition for conferring protection in a host against a disease caused by Haemophilus influenzae, comprises two different antigens of H. influenzae, where one of the antigens is an adhesin.

DC B04 D16

IN KLEIN, M H; LOOSMORE, S M; YANG, Y

PA (CONN-N) CONNAUGHT LAB LTD; (KLEI-I) KLEIN M H; (LOOS-I) LOOSMORE S M;  
(YANG-I) YANG Y; (AVET) AVENTIS PASTEUR LTD

CYC 91

PI WO 2000035477 A2 20000622 (200037)\* EN 44 A61K039-102 <--  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ TZ UG ZW  
W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES  
FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS  
LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL  
TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2000015439 A 20000703 (200046)

EP 1140158 A2 20011010 (200167) EN A61K039-102 <--  
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

JP 2002532433 W 20021002 (200279) 55 A61K039-102 <--

NZ 512679 A 20030829 (200365) A61K039-102 <--

AU 772882 B2 20040513 (200462) A61K039-102 <--

ADT WO 2000035477 A2 WO 1999-CA1189 19991215; AU 2000015439 A AU 2000-15439  
19991215; EP 1140158 A2 EP 1999-957822 19991215; WO 1999-CA1189 19991215;  
JP 2002532433 W WO 1999-CA1189 19991215; JP 2000-587796 19991215; NZ  
512679 A NZ 1999-512679 19991215; WO 1999-CA1189 19991215; AU 772882 B2 AU  
2000-15439 19991215

FDT AU 2000015439 A Based on WO 2000035477; EP 1140158 A2 Based on WO  
2000035477; JP 2002532433 W Based on WO 2000035477; NZ 512679 A Based on  
WO 2000035477; AU 772882 B2 Previous Publ. AU 2000015439, Based on WO  
2000035477

PRAI US 1998-210995 19981215

IC ICM A61K039-102

ICS A61K039-05; A61K039-08; A61K039-10;  
A61K039-116; A61K039-13; A61K039-295;  
A61K039-39; A61P027-16; A61P031-04; C07K014-285

ICA C12N015-09

AB WO 2000035477 A UPAB: 20000807

NOVELTY - A new immunogenic composition (I) for conferring protection in a host against a disease caused by Haemophilus influenzae, comprises at least two different antigens of Haemophilus influenzae, where at least one of the antigens is an adhesin.

DETAILED DESCRIPTION - An INDEPENDENT CLAIM is included for a method of immunizing a host against disease caused by infection with Haemophilus influenzae, including otitis media, comprising administering to the host an immunoeffective amount of (I).

ACTIVITY - Antibacterial.

MECHANISM OF ACTION - Vaccine.

H91A Hin47 is partially protective in the chinchilla model of otitis media, as described in the US Patent Number 5,506,139.

In this model, 1 to 2 year old chinchillas (Moulton Chinchilla Ranch, Rochester, Minnesota) were immunized intramuscularly (i.m.) on days 0, 14 and 28 with 30 micro g of H91A Hin47 adsorbed to alum, and challenged on day 44 with 50 to 350 colony forming units (cfu) of live organisms delivered into the middle ear space via the epitympanic bulla. Animals were monitored by tympanometry and middle ear fluid was collected 4 days post challenge, mixed with 200 micro l of BHI (undefined) medium and dilutions plated onto chocolate agar plates that were incubated for 24 hours at 37 deg. C. Convalescent animals or those mock-immunized with alum

alone, were used as controls. For the multi-component vaccine study, 50 micro g of H91A Hin47 was mixed with 50 micro g of recombinant HMW (rHMW) and chinchillas were immunized as described above.

The results of the protection study indicate that there was still partial protection afforded in the intrabulla challenge model by the combination of H91A Hin47 and rHMW.

USE - The two different antigens of H. influenzae, at least one of which is an adhesin, are useful in the manufacture of a vaccine for conferring protection against disease caused by infection with H. influenzae, including otitis media. (I) is used as a vaccine (all claimed) against diseases caused by H. influenzae infection.

Dwg.0/12

FS CPI  
FA AB; DCN  
MC CPI: B04-N0300E; B14-A01A; B14-S11B; D05-H07

L54 ANSWER 15 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2000-412339 [35] WPIX

DNN N2000-308180 DNC C2000-125066

TI Nucleic acids encoding polypeptide antigens from Chlamydia useful for preventing, diagnosing and treating diseases such as community acquired pneumonia, bronchitis, sinusitis and asthmatic bronchitis, adult-onset asthma.

DC B04 C06 D16 S03

IN MURDIN, A D; OOMEN, R P; WANG, J; JACOBSON, E L; JACOBSON, M K

PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD; (JACO-I)

JACOBSON E L; (JACO-I) JACOBSON M K

CYC 91

PI WO 2000032794 A2 20000608 (200035)\* EN 173 C12N015-62  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ TZ UG ZW  
W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES  
FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS  
LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL  
TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2000015405 A 20000619 (200044)

EP 1135509 A2 20010926 (200157) EN C12N015-62

R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

US 2002123517 A1 20020905 (200260) A61K031-44

JP 2002531095 W 20020924 (200278) 175 C12N015-00

NZ 512308 A 20040130 (200414) C12N015-62

MX 2001005617 A1 20030401 (200415) A61K039-118 <--

ADT WO 2000032794 A2 WO 1999-CA1147 19991201; AU 2000015405 A AU 2000-15405  
19991201; EP 1135509 A2 EP 1999-957785 19991201, WO 1999-CA1147 19991201;  
US 2002123517 A1 Provisional US 1998-110428P 19981201, CIP of US  
1999-452617 19991201, Div ex US 2000-549691 20000414, US 2002-113681  
20020508; JP 2002531095 W WO 1999-CA1147 19991201, JP 2000-585425  
19991201; NZ 512308 A NZ 1999-512308 19991201, WO 1999-CA1147 19991201; MX  
2001005617 A1 WO 1999-CA1147 19991201, MX 2001-5617 20010601

FDT AU 2000015405 A Based on WO 2000032794; EP 1135509 A2 Based on WO  
2000032794; US 2002123517 A1 CIP of US 6337065, Div ex US 6403619; JP  
2002531095 W Based on WO 2000032794; NZ 512308 A Based on WO 2000032794;  
MX 2001005617 A1 Based on WO 2000032794

PRAI US 1998-110438P 19981201; US 1998-110339P 19981201;  
US 1998-110340P 19981201; US 1998-110427P 19981201;  
US 1998-110428P 19981201; US 1999-452617 19991201;  
US 2000-549691 20000414; US 2002-113681 20020508

IC ICM A61K031-44; A61K039-118; C12N015-00; C12N015-62

ICS A61K031-7088; A61K039-385; A61K039-39;  
A61K039-395; A61K048-00; A61P031-00; C07K007-08; C07K014-295;  
C07K014-705; C07K016-12; C07K019-00; C12N001-15; C12N001-19;  
C12N001-21; C12N005-10; C12P021-02; C12Q001-68; G01N033-15;  
G01N033-50; G01N033-53; G01N033-569

AB WO 200032794 A UPAB: 20000725

NOVELTY - Nucleic acids (NAM1) encoding polypeptide (PEP1) antigens from Chlamydia, are new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are included for the following:

(1) a nucleic acid molecule (NAM1) comprising a sequence selected from:

(a) nucleic acid sequences (N1)-(N10), which are defined sequences given in the specification;

(b) a sequence encoding a polypeptide encoded by (N1)-(N10);

(c) a sequence comprising at least 38 consecutive nucleotides from (N1)-(N10); and/or

(d) a sequence which encodes a polypeptide at least 75% identical in amino acid sequence to the polypeptides encoded by (N1)-(N10);

(2) a nucleic acid molecule (NAM1') that comprises an antisense sequence to NAM1;

(3) a nucleic acid molecule (NAM2) comprising a sequence encoding a fusion protein comprising the polypeptide encoded by NAM1 and an additional polypeptide;

(4) a vaccine composition (VAC1) comprising a vaccine vector and NAM1 and/or NAM2 expressed as a polypeptide (the vaccine may comprise an additional polypeptide that enhances the immune response to the polypeptide expressed by NAM1);

(5) a unicellular host (CELL1) transformed with either NAM1 and/or NAM2;

(6) a nucleic acid probe of 5-100 nucleotides which hybridizes under stringent conditions to (N1)-(N10) (or homologs, complementary and/or antisense sequences of them);

(7) a primer of 10-40 nucleotides which hybridizes under stringent conditions to (N1)-(N10) (or homologs, complementary and/or antisense sequences of them);

(8) a polypeptide (PEP1) encoded by NAM1 and/or NAM2;

(9) a fusion peptide (PEP2) comprising PEP1 and an additional polypeptide;

(10) a method (METH1) for producing PEP1 and/or PEP2 comprising culturing CELL1;

(11) an antibody (ANB1) against PEP1 and/or PEP2;

(12) a method (METH2) for preventing and/or treating Chlamydia infections using NAM1 and/or NAM2, VAC1, PEP1 and/or PEP2 or ANB1;

(13) a method (METH3) for detecting Chlamydia infection comprising assaying a body fluid of a mammal to be tested with either NAM1 and/or NAM2, PEP1 or ANB1;

(14) a diagnostic kit comprising instructions for use and either NAM1 and/or NAM2, PEP1 or ANB1; and

(15) a method (METH4) for identifying a PEP1 and/or PEP2 which induces an immune response that prevents or lessens the severity of a Chlamydia infection in a mammal previously immunized with the polypeptide, comprising:

(a) immunizing a mouse with the polypeptide; and

(b) inoculating the immunized mouse with Chlamydia (the polypeptide which lessens or prevents the severity of the Chlamydia infection in the immunized mouse compared to a non-immunized control mouse is identified).

ACTIVITY - Bactericide.

No data given.

MECHANISM OF ACTION - Vaccine.

USE - The nucleic acids (and complementary sequences) may be used as diagnostic agents for detecting the presence of nucleic acids encoding Chlamydia antigens in samples according to standard methods, and therefore, for diagnosing Chlamydia infections. For example, they may be used as primers and probes for diagnostic polymerase chain reaction (PCR) assays. Antisense sequences may be used to down regulate expression of the proteins and may be used to treat infections. The nucleic acids may also be used to produce the protein antigens they encode according to standard recombinant DNA methodologies. The proteins may then be used as antigens for the production of antibodies (i.e. as vaccines) for preventing infection by Chlamydia. The antibodies may also be used as diagnostic reagents for detecting infections. Chlamydia is a pathogen implicated in the development of (for example) community acquired pneumonia, upper respiratory tract disease (especially bronchitis and sinusitis, asthmatic bronchitis, adult-onset asthma and acute exacerbations of asthma in adults.

Dwg. 0/0

FS CPI EPI

FA AB; DCN

MC CPI: B04-B04C1; B04-C01; B04-E03F; B04-E04; B04-E05; B04-E06; B04-E08; B04-F0100E; B04-F10A; B04-G07; B04-N03A0E; B04-P01B; B11-A; B11-C07A4; B11-C08E; B11-C09; B12-K04A4; B12-K04F; B12-M05; B14-A01A; B14-S11B; B14-S12; C04-B04C1; C04-C01; C04-E03F; C04-E04; C04-E05; C04-E06; C04-E08; C04-F0100E; C04-F10A; C04-G07; C04-N03A0E; C04-P01B; C11-A; C11-C07A4; C11-C08E; C11-C09; C12-K04A4; C12-K04F; C12-M05; C14-A01A; C14-S11B; C14-S12; D05-A01A4; D05-A01B; D05-C12; D05-H04; D05-H07; D05-H08; D05-H09; D05-H11; D05-H12A; D05-H12C; D05-H12D; D05-H12E; D05-H14; D05-H17A5; D05-H17C; D05-H18  
EPI: S03-E14H4

L54 ANSWER 16 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN  
AN 2000-412330 [35] WPIX

Search done by Noble Jarrell

DNN N2000-308175 DNC C2000-125057  
 TI New polynucleotide encoding the Chlamydia 98 kiloDalton outer membrane protein, useful for preventing or treating Chlamydia infection.  
 DC B04 D16 S03  
 IN DUNN, P; MURDIN, A D; OOMEN, R P; WANG, J  
 PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD; (DUNN-I) DUNN P; (MURD-I) MURDIN A D; (OOMEN-I) OOMEN R P; (WANG-I) WANG J  
 CYC 91  
 PI WO 2000032784 A1 20000608 (200035)\* EN 94 C12N015-31  
 RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
 OA PT SD SE SL SZ TZ UG ZW  
 W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES  
 FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS  
 LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL  
 TJ TM TR TT TZ UA UG UZ VN YU ZA ZW  
 AU 2000037909 A 20000619 (200044)  
 EP 1135501 A1 20010926 (200157) EN C12N015-31  
 R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
 RO SE SI  
 US 2002094340 A1 20020718 (200254) A61K039-118 <--  
 JP 2002531093 W 20020924 (200278) 93 C12N015-09  
 US 2003157124 A1 20030821 (200356) A61K039-02 <--  
 MX 2001005616 A1 20030401 (200415) A61K039-118 <--  
 NZ 529361 A 20040625 (200445) C12N015-63  
 ADT WO 2000032784 A1 WO 1999-CA1148 19991201; AU 2000037909 A AU 2000-37909 19991201; EP 1135501 A1 EP 1999-957786 19991201; WO 1999-CA1148 19991201; US 2002094340 A1 Provisional US 1998-113439P 19981223, Provisional US 1999-132272P 19990503, US 1999-452380 19991201; JP 2002531093 W WO 1999-CA1148 19991201, JP 2000-585415 19991201; US 2003157124 A1 Provisional US 1998-110439P 19981201, Provisional US 1999-132272P 19990503, Cont of US 1999-452380 19991201, US 2002-324129 20021220; MX 2001005616 A1 WO 1999-CA1148 19991201, MX 2001-5616 20010601; NZ 529361 A Div ex NZ 2000-512354 20000114, NZ 2000-529361 20000114  
 FDT AU 2000037909 A Based on WO 2000032784; EP 1135501 A1 Based on WO 2000032784; JP 2002531093 W Based on WO 2000032784; MX 2001005616 A1 Based on WO 2000032784; NZ 529361 A Div ex NZ 512354  
 PRAI US 1999-132272P 19990503; US 1998-110439P 19981201;  
 US 1998-113439P 19981223; US 1999-452380 19991201;  
 US 2002-324129 20021220  
 IC ICM A61K039-02; A61K039-118; C12N015-09; C12N015-31;  
 C12N015-63  
 ICS A61K031-711; A61K038-00; A61K039-39; A61K039-395;  
 A61K048-00; A61P031-04; C07H021-04; C07K014-295; C07K014-705;  
 C07K016-12; C07K019-00; C12N001-15; C12N001-19; C12N001-21;  
 C12N005-10; C12N015-11; C12N015-62; C12N015-74; C12P021-02;  
 C12Q001-68; G01N033-15; G01N033-50; G01N033-53; G01N033-569  
 AB WO 200032784 A UPAB: 20000725  
 NOVELTY - Isolated polynucleotide (N1) encoding the Chlamydia 98 kiloDalton (kDa) outer membrane protein, known as CPN100640, is new.  
 DETAILED DESCRIPTION - Isolated polynucleotide (N1) encoding the Chlamydia 98 kiloDalton (kDa) outer membrane protein, is new.  
 N1 comprises a nucleic acid sequence selected from:  
 (a) the 3050 (I) or 2808 (II) nucleotide sequence defined in the specification;  
 (b) a sequence which encodes a polypeptide encoded by (I) or (II);  
 (c) a sequence comprising at least 38 consecutive nucleotides from any one of the nucleic acid sequences of (a) and (b); and  
 (d) a sequence which encodes a polypeptide which is at least 75% identical in amino acid sequence to the polypeptides encoded by (I) or (II).  
 INDEPENDENT CLAIMS are also included for the following:  
 (1) a nucleic acid molecule (N2) comprising a nucleic acid sequence which encodes a polypeptide selected from:  
 (a) the 936 (III) or 925 (IV) amino acid sequence defined in the specification;  
 (b) an immunogenic fragment comprising at least 12 consecutive amino acids from a polypeptide of (a); and  
 (c) a polypeptide of (a) or (b) which has been modified to improve its immunogenicity, where the modified polypeptide is at least 75% identical in amino acid sequence to the corresponding polypeptide of (a) or (b);  
 (2) a nucleic acid molecule (N3) comprising a nucleic acid sequence which is anti-sense to the nucleic acid molecule of (1) or N1;  
 (3) a nucleic acid molecule (N4) comprising a nucleic acid sequence which encodes a fusion protein comprising a polypeptide encoded by N1 and an additional polypeptide;



(4) a vaccine comprising at least one first nucleic acid of N1, N2 or N4 and a vaccine vector, where each first nucleic acid is expressed as a polypeptide, the vaccine optionally comprising a second nucleic acid encoding an additional polypeptide which enhances the immune response to the polypeptide expressed by the first nucleic acid;

(5) a unicellular host transformed with a nucleic acid (N1, N2, N3 or N4) operatively linked to one or more expression control sequences;

(6) a nucleic acid probe of 5 to 100 nucleotides which hybridizes under stringent conditions to N1 or N2, or to a homolog or complementary or anti-sense sequence of the nucleic acid molecule;

(7) a primer of 10 to 40 nucleotides which hybridizes under stringent conditions to N1 or N2, or to a homolog or complementary or anti-sense sequence of the nucleic acid molecule;

(8) a polypeptide (P1) encoded by N1, N2 or N4;

(9) a polypeptide (P2) comprising an amino acid sequence selected from:

(a) (III) or (IV);

(b) an immunogenic fragment comprising at least 12 consecutive amino acids from a polypeptide of (a); and

(c) a polypeptide of (a) or (b) which has been modified to improve its immunogenicity, where the modified polypeptide is at least 75% identical in amino acid sequence to the corresponding polypeptide of (a) or (b);

(10) a fusion polypeptide (P3) comprising P1 or P2, and an additional polypeptide;

(11) a method for producing P1 or P2, comprising culturing the unicellular host of (5);

(12) an antibody against P1, P2 or P3;

(13) a vaccine comprising at least one first polypeptide of P1, P2 or P3, and optionally comprising a second polypeptide which enhances the immune response to the first polypeptide;

(14) a method of detecting Chlamydia infection comprising the step of assaying a body fluid of a mammal to be tested, with a component selected from:

(a) N1, N2, N3 or N4;

(b) P1, P2 or P3; or

(c) the antibody of (12);

(15) a diagnostic kit comprising a component selected from those defined in the method of (14);

(16) a method for identifying a polypeptide of P1, P2 or P3 which induces an immune response effective to prevent or lessen the severity of Chlamydia infection in a mammal previously immunized with polypeptide, comprising:

(a) immunizing a mouse with the polypeptide; and

(b) inoculating the immunized mouse with Chlamydia, where the immunized mouse is compared with a non-immunized mouse control to identify the polypeptide; and

(17) expression plasmid pCAI640.

#### ACTIVITY - Antibacterial.

Groups of 7 to 9 week old male Balb/c mice (8 to 10 per group) were immunized intramuscularly (i.m.) plus intranasally (i.n.) with plasmid DNA containing the coding sequence of *C. pneumoniae* 98 kDa outer membrane protein gene. Saline or the plasmid vector lacking an inserted chlamydial gene was given to groups of control animals.

For i.m. immunization alternate left and right quadriceps were injected with 100 micro g of DNA in 50 micro l of phosphate buffered saline (PBS) on three occasions at 0, 3 and 6 weeks. For i.n. immunization, anaesthetized mice aspirated 50 micro l of PBS containing 50 micro g DNA on three occasions at 0, 3 and 6 weeks. At week 8, immunized mice were inoculated i. n. with 5 x 10<sup>5</sup> IFU (undefined) of *C. pneumoniae*, strain AR39 in 100 micro l of SPG (7.5 % sucrose, 5 mM glutamate, 12.5 mM phosphate pH 7.5) buffer to test their ability to limit the growth of a sublethal *C. pneumoniae* challenge.

Lungs were taken from mice at days 5 and 9 post-challenge and immediately homogenized in SPG buffer. Dilutions of the homogenate were assayed for the presence of infectious Chlamydia by inoculation onto monolayers of susceptible cells. The inoculum was centrifuged onto the cells at 3000 rotations per minute (rpm) for 1 hour, then the cells were incubated for three days at 35 deg. C in the presence of 1 micro g/ml cycloheximide. After incubation the monolayers were fixed with formalin and methanol then immunoperoxidase stained for the presence of chlamydial inclusions using convalescent sera from rabbits infected with *C. pneumoniae* and metal-enhanced DAB (undefined) as a peroxidase substrate.

Mice immunized i.n. and i.m. with pCAI640 had chlamydial lung titers less than 255,000 in 4 of 4 cases at day 5 and less than 423,200 in 4 of 4 cases at day 9 while the range of values for control mice immunized with

saline was 227,000-934,200 IFU/lung (mean 685,240) at day 5 and 96,000-494,000 IFU/lung (mean 238,080) at day 9.

# MECHANISM OF ACTION - Vaccine.

USE - The nucleic acids, proteins, antibodies and vaccines are useful for preventing or treating Chlamydia infection (claimed).

Dwg. 0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-C01G; B04-E02F; B04-E03F; B04-E05; B04-E06; B04-E08; B04-F0100E; B04-F1100E; B04-G01; B04-G07; B04-N03A0E; B11-C08E; B12-K04A4; B12-K04F; B14-A01A; B14-S11B; D05-H07; D05-H09; D05-H11; D05-H12A; D05-H12B2; D05-H12D1; D05-H12D2; D05-H12E; D05-H14A; D05-H17A6; D05-H17B6  
EPI: S03-E14H4

L54 ANSWER 17 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2000-352521 [31] WPIX

DNC C2000-107478

TI Novel multivalent vaccine composition for use solely as a booster against in pre-sensitized subjects, to protect against e.g., diphtheria, poliomyelitis and tetanus.

DC B04 D16

IN CARTIER, J R; LAROCHE, P

PA (INMR) PASTEUR MERIEUX MSD; (AVET) AVENTIS PASTEUR; (AVET)

AVENTIS PASTEUR MSD

CYC 88

PI EP 1004314 A1 20000531 (200031)\* FR 11 A61K039-295 <--  
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

WO 2000030678 A1 20000602 (200033) FR  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ TZ UG ZW  
W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB  
GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU  
LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR  
TT UA UG US UZ VN YU ZA ZW

AU 2000013901 A 20000613 (200043)  
EP 1131094 A1 20010912 (200155) FR A61K039-295 <--  
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

CZ 2001001860 A3 20020116 (200215) A61K039-295 <--

ZA 2001003634 A 20020731 (200271) 44 A61K000-00

NZ 511933 A 20020927 (200272) A61K039-295 <--

AU 759221 B 20030410 (200337) A61K039-295 <--

JP 2003525858 W 20030902 (200358) 31 A61K039-05 <--

EP 1131094 B1 20041103 (200475) FR A61K039-295 <--

R: AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE

ADT EP 1004314 A1 EP 1998-122373 19981126; WO 2000030678 A1 WO 1999-FR2913 19991125; AU 2000013901 A AU 2000-13901 19991125; EP 1131094 A1 EP 1999-972544 19991125; WO 1999-FR2913 19991125; CZ 2001001860 A3 WO 1999-FR2913 19991125; CZ 2001-1860 19991125; ZA 2001003634 A ZA 2001-3634 20010504; NZ 511933 A NZ 1999-511933 19991125; WO 1999-FR2913 19991125; AU 759221 B AU 2000-13901 19991125; JP 2003525858 W WO 1999-FR2913 19991125; JP 2000-583561 19991125; EP 1131094 B1 EP 1999-972544 19991125; WO 1999-FR2913 19991125

FDT AU 2000013901 A Based on WO 2000030678; EP 1131094 A1 Based on WO 2000030678; CZ 2001001860 A3 Based on WO 2000030678; NZ 511933 A Based on WO 2000030678; AU 759221 B Previous Publ. AU 2000013901, Based on WO 2000030678; JP 2003525858 W Based on WO 2000030678; EP 1131094 B1 Based on WO 2000030678

PRAI EP 1998-122373 19981126

IC ICM A61K000-00; A61K039-05; A61K039-295

ICS A61K039-08; A61K039-10; A61K039-13;

A61K039-29; A61P025-00; A61P031-04

AB EP 1004314 A UPAB: 20040511

NOVELTY - Vaccine formulated for exclusive use as a repeat ('booster') vaccine in a population already having received a primary vaccine and/or sensitized against at least the poliovirus, Corynebacterium diphtheriae and/or C. tetani, is new.

DETAILED DESCRIPTION - Vaccine (I) formulated for exclusive use as a repeat ('booster') vaccine in a population already having received a primary vaccine and/or sensitized against at least the poliovirus, Corynebacterium diphtheriae and/or C. tetani, comprises,

- (1) at least 1.2 mg/ml of aluminum salt;
- (2) antigens derived from at least the poliovirus; and
- (3) a quantity of diphtheria anatoxin (DA) used as an antigen of C.

diphtheriae comprising between 4-16 Flocculation units (Fu);

ACTIVITY - Immunostimulant; anti-viral.

MECHANISM OF ACTION - Vaccine. Three lots of 0.5 ml per vaccine were used in a study of 31 adults, the lots differed only in the quantity of diphtheria anatoxin, 2 (lot A), 5 (B) and 8 (C) Fu/ml, the adults were divided into 3 groups of 10, one for each lot. Each subject was injected into their deltoid muscle. No systemic reactions were reported for any of the lots, although localized reactions were noted over the first week in 8 subjects in lot A, 6 subjects in lot B and 8 subjects in lot C. Symptoms included redness and swellings around the site of injection, although all symptoms disappeared without treatment and did not affect the quality of life of the subjects. No reactions were observed beyond the first week and no serious reactions were observed in any subjects. In addition the immune responses to the 5 antigens were excellent for all three lots, despite initially elevated titers due to the young age of the subjects and because of recent vaccinations, a booster effect was obtained for each antigen.

ADVANTAGE - The vaccine (I) is specifically designed for use as a booster vaccine only, and as such avoids the reduced immunogenicity that occurs when administering reduced dosages of normal primary vaccines. The quantity of immunogenic diphtheria anatoxin used (10 Fu/ml) and allows optimal immunogenic protection while minimizing undesirable side-effects, such as allergic reactions to the antigens.

Dwg.0/0

FS

CPI

FA

MC

AB; DCN  
CPI: B04-B04M; B04-F11; B14-A01A; B14-A01A1; B14-A01B;  
B14-A02A4; B14-A02A5; B14-S11A; B14-S11B;  
D05-H07; D05-H08; D05-H14

L54 ANSWER 18 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN  
AN 2000-350742 [30] WPIX  
DNN N2000-262745 DNC C2000-106768  
TI Isolated polynucleotide encoding a Chlamydia polypeptide useful to treat,  
diagnose and prevent disease caused by Chlamydia infection.  
DC B04 D16 S03  
IN DUNN, P L; MURDIN, A D; OOMEN, R P  
PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD; (DUNN-I)  
DUNN P L; (MURD-I) MURDIN A D; (OOM-I) OOMEN R P  
CYC 91  
PI WO 2000024901 A1 20000504 (200030)\* EN 88 C12N015-31  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ TZ UG ZW  
W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES  
FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS  
LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL  
TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW  
AU 9963593 A 20000515 (200039)  
EP 1124964 A1 20010822 (200149) EN C12N015-31  
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI  
US 6403101 B1 20020611 (200244) A61K039-118 <--  
US 2002091096 A1 20020711 (200248) A61K048-00  
JP 2002528081 W 20020903 (200273) 104 C12N015-09  
MX 2001004356 A1 20030701 (200366) A61K039-02 <--  
US 6642025 B2 20031104 (200374) C12P021-06  
NZ 511886 A 20031219 (200404) C12N015-31  
AU 770905 B2 20040304 (200453) C12N015-31  
ADT WO 2000024901 A1 WO 1999-GB3565 19991028; AU 9963593 A AU 1999-63593  
19991028; EP 1124964 A1 EP 1999-951017 19991028; WO 1999-GB3565 19991028;  
US 6403101 B1 Provisional US 1998-106037P 19981028, Provisional US  
1999-154658P 19990920, US 1999-427501 19991026; US 2002091096 A1  
Provisional US 1998-106037P 19981028, Provisional US 1999-154658P  
19990920, Div ex US 1999-427501 19991026, US 2001-905119 20010713; JP  
2002528081 W WO 1999-GB3565 19991028, JP 2000-578453 19991028; MX  
2001004356 A1 WO 1999-GB3565 19991028, MX 2001-4356 20010430; US 6642025  
B2 Provisional US 1998-106037P 19981028, Provisional US 1999-154658P  
19990920, Div ex US 1999-427501 19991026, US 2001-905119 20010713; NZ  
511886 A NZ 1999-511886 19991028, WO 1999-GB3565 19991028; AU 770905 B2 AU  
1999-63593 19991028  
FDT AU 9963593 A Based on WO 2000024901; EP 1124964 A1 Based on WO 2000024901;  
JP 2002528081 W Based on WO 2000024901; MX 2001004356 A1 Based on WO  
2000024901; US 6642025 B2 Div ex US 6403101; NZ 511886 A Based on WO  
2000024901; AU 770905 B2 Previous Publ. AU 9963593, Based on WO 2000024901  
PRAI US 1999-427501 19991026; US 1998-106037P 19981028;  
US 1999-154658P 19990920; US 2001-905119 20010713  
IC ICM A61K039-02; A61K039-118; A61K048-00; C12N015-09;

C12N015-31; C12P021-06

ICS A61K038-00; A61K039-00; A61K039-39; A61P009-10;  
A61P011-00; A61P011-06; C07H021-04; C07K001-22; C07K014-295;  
C07K016-12; C12N001-15; C12N001-19; C12N001-21; C12N005-10;  
C12N015-62; C12P021-02; C12Q001-68; G01N033-53; G01N033-543;  
G01N033-566; G01N033-569

ICI C12N001-21; C12N015-09; C12P021-02; C12R001:01; C12R001:01; C12R001:01  
AB WO 200024901 A UPAB: 20000624

NOVELTY - An isolated polynucleotide (N1) encoding a lorf2 protein of a strain of *Chlamydia pneumoniae*, is new.

DETAILED DESCRIPTION - An isolated polynucleotide (N1) has a nucleotide sequence which comprises:

(a) a defined nucleotide sequence (I) of 1550 base pairs or functional fragments of (I);

(b) a nucleotide sequence encoding a polypeptide with a sequence at least 75% homologous to (II) which has a defined protein sequence of 422 amino acids, or functional fragments; or

(c) a sequence capable of hybridizing under stringent conditions to a sequence comprising (I), or functional fragments.

INDEPENDENT CLAIMS are also included for the following:

(1) an isolated polypeptide (P1) with a sequence at least 75% homologous to (II), or functional fragments of (II);

(2) a polypeptide P2 comprising P1 linked to a fusion polypeptide;

(3) an expression cassette comprising N1 operably linked to a promoter;

(4) an expression vector comprising the expression cassette of (3);

(5) a host cell comprising the expression cassette of (3);

(6) a method of producing a recombinant polypeptide with sequence (II) comprising culturing the host cell of (5) and recovering the polypeptide;

(7) a vaccine vector comprising the expression cassette of (3);

(8) a pharmaceutical composition containing P1 and one or more known *Chlamydia* antigens;

(9) a method for inducing an immune response in a mammal comprising administering the vaccine vector of (7) or a composition containing P1 to induce an immune response;

(10) a polynucleotide probe reagent capable of detecting the presence of *Chlamydia* in biological material comprising a polynucleotide that hybridizes to N1 under stringent conditions;

(11) a hybridization method for detecting the presence of *Chlamydia* in a sample comprising:

(a) obtaining polynucleotide from the sample;

(b) hybridizing the obtained polynucleotide with the polynucleotide probe reagent of (10) under conditions allowing hybridization of the probe and the sample; and

(c) detecting any hybridization occurring;

(12) an amplification method for detecting the presence of *Chlamydia* in a sample comprising:

(a) obtaining polynucleotide from the sample;

(b) amplifying the polynucleotide using one or more polynucleotide probe reagents of (10); and

(c) detecting the amplified polynucleotide;

(13) a method for detecting the presence of *Chlamydia* in a sample comprising contacting the sample with a detecting reagent that binds to P1 in the sample and detecting the formed complex;

(14) an affinity chromatography method for substantially purifying a polypeptide with sequence (II) comprises:

(a) contacting a sample containing (II) with a detecting reagent that binds to the polypeptide to form a complex;

(b) isolating the formed complex;

(c) dissociating the formed complex; and

(d) isolating the dissociated polypeptide; and

(15) an antibody that immunospecifically binds P1 or a fragment or derivative of the antibody containing its binding domain.

ACTIVITY - Antibacterial.

MECHANISM OF ACTION - Vaccine.

Balb/c mice (7-9 weeks old) were immunized intramuscularly and intranasally with plasmid DNA containing the coding sequence of *C. pneumoniae* lorf2 gene. Control animals were given saline or the plasmid vector without the chlamydial gene. The intramuscular immunization comprised 100 micro g DNA in 50 micro l phosphate buffered saline (PBS) at 0, 3 and 6 weeks and the intranasal immunization comprised 50 micro g DNA in 50 micro l PBS at 0, 3 and 6 weeks. At week 8, immunized mice were inoculated intranasally with 5x10<sup>5</sup> inclusion forming units (IFU) of *C. pneumoniae*, strain AR39 in 100 micro l SPG (sucrose, glutamate, phosphate) buffer. Lungs were taken from the mice at day 9 post challenge and

homogenized in SPG buffer, the homogenate was assayed for the presence of infectious chlamydia by inoculation onto monolayers of susceptible cells. After incubation the monolayers were fixed and immunoperoxidase stained for the presence of chlamydial inclusions using convalescent sera from rabbits infected with *C. pneumoniae* and metal-enhanced DAB (not defined) as a peroxidase substrate. Mice immunized with the plasmid containing the *lorf2* gene had an average chlamydial lung titer of 11050 IFU/lung compared to 111783 IFU/lung for the control mice immunized with saline.

USE - The polynucleotides and polypeptides can be used as a vaccine for humans to treat or prevent disease caused by Chlamydia infection and P1, N1 or an antibody to P1 can be used to diagnose a Chlamydia infection.

Dwg.0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-B04D5; B04-C01G; B04-E03F; B04-E05; B04-E08; B04-F0100E;

B04-F10A; B04-G01; B04-G21; B04-G22; B04-N03A; B11-C07A;

B11-C08E3; B11-C08E5; B12-K04A4; B12-K04F; B14-A01A;

B14-S11B; D05-H04; D05-H07; D05-H09; D05-H11A; D05-H11B;

D05-H12A; D05-H12D1; D05-H12E; D05-H14; D05-H17A6; D05-H18B

EPI: S03-E14H4

L54 ANSWER 19 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2000-350688 [30] WPIX

DNC C2000-106714

TI Chlamydia antigenes and the proteins they encode, useful for vaccinating against Chlamydia infections that affect the respiratory tract.

DC B04 D16

IN MURDIN, A D; OOMEN, R P; WANG, J; DUNN, P

PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD

CYC 91

PI WO 2000024765 A2 20000504 (200030)\* EN 165 C07K014-00

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL

OA PT SD SE SL SZ TZ UG ZW

W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES

FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS

LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL

TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2000012541 A 20000515 (200039)

EP 1129202 A2 20010905 (200151) EN C12N015-62

R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT

RO SE SI

JP 2002530052 W 20020917 (200276) 222 C12N015-09

MX 2001004291 A1 20030601 (200417) C07K014-00

ADT WO 2000024765 A2 WO 1999-CA992 19991028; AU 2000012541 A AU 2000-12541

19991028; EP 1129202 A2 EP 1999-955602 19991028, WO 1999-CA992 19991028;

JP 2002530052 W WO 1999-CA992 19991028, JP 2000-578335 19991028; MX

2001004291 A1 WO 1999-CA992 19991028, MX 2001-4291 20010427

FDT AU 2000012541 A Based on WO 2000024765; EP 1129202 A2 Based on WO

2000024765; JP 2002530052 W Based on WO 2000024765; MX 2001004291 A1 Based

on WO 2000024765

PRAI US 1998-107035P 19981102; US 1998-106034P 19981028;

US 1998-106039P 19981028; US 1998-106042P 19981028;

US 1998-106044P 19981028; US 1998-106072P 19981029;

US 1998-106073P 19981029; US 1998-106074P 19981029;

US 1998-106087P 19981029; US 1998-106587P 19981102;

US 1998-106588P 19981102; US 1998-106589P 19981102;

US 1998-107034P 19981102

IC ICM C07K014-00; C12N015-09; C12N015-62

ICS A61K038-00; A61K039-118; A61K039-395; A61K048-00;

A61P031-04; C07K014-295; C07K016-12; C12N001-15; C12N001-19;

C12N001-21; C12N005-10; C12Q001-68; G01N033-15; G01N033-50;

G01N033-53; G01N033-569

AB WO 200024765 A UPAB: 20000624

NOVELTY - Nucleic acids (A) encoding Chlamydia antigenes and the proteins

(B) they express, are new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are included for the following:

(1) a nucleic acid molecule (A) comprising a sequence encoding a polypeptide (B) selected from:

(a) one of 19 defined amino acid sequences ((IIa) - (IIs)) given in the specification;

(b) an immunogenic fragment comprising at least 12 consecutive amino acids from (IIa) - (IIs); and/or

(c) a modified form of the polypeptide sequences (IIa) - (IIs) which has been modified to improve its immunogenicity (the modified peptide is at least 75% identical to the corresponding amino acid sequence of (IIa) -

- (IIb));
- (2) a polypeptide (B) encoded by (A);
  - (3) a nucleic acid encoding a fusion protein comprising a polypeptide encoded by (A) and an additional polypeptide;
  - (4) a fusion protein comprising (B) and an additional polypeptide;
  - (5) a vaccine (C) comprising (A) and a vaccine vector which express (B) (and optionally comprising a second nucleic acid (D) encoding an additional polypeptide (J) which enhances the immune response to (A) and/or (B);
  - (6) a nucleic acid probe (E) (of 5 to 100 nucleotides) or primer (F) (of 10 to 40 nucleotides) which hybridizes under stringent conditions to (Ia) - (Iz) (or a homolog, complementary or antisense sequence of (Ia) - (Iz));
  - (7) a unicellular host (G) transformed with (A);
  - (8) a method for producing (B) comprising culturing (G);
  - (9) a vaccine (H) comprising (B) (and optionally comprising an additional polypeptide (J));
  - (10) an antibody (K) against (B);
  - (11) a method for preventing or treating Chlamydia infection, using:
    - (a) (A);
    - (b) (C) and/or (H);
    - (c) (B); and/or
    - (d) (K);
  - (12) a method for detecting Chlamydia infection comprising assaying a body fluid of a mammal with either:
    - (a) (A);
    - (b) (B); and/or
    - (c) (K); and
  - (13) a diagnostic kit comprising instructions for use and a component selected from:
    - (a) (A);
    - (b) (B); and/or
    - (c) (K).

ACTIVITY - Antiinflammatory; respiratory; antibacterial; anti-asthmatic; antiarteriosclerotic.

No biological data given.

MECHANISM OF ACTION - Vaccine.

USE - The nucleic acids may be used for the recombinant production of the Chlamydia polypeptides (either in vivo or in vitro) according to standard recombinant DNA methodologies. The polypeptides may then be used to vaccinate against Chlamydia infections in mammals. Chlamydia, such as *C. pneumoniae*, are pathogens responsible for upper respiratory tract infections such as community acquired pneumonia, acute respiratory disease and bronchitis and may be implicated in atherosclerotic changes and asthma.

The nucleic acids may also be used as probes for detecting the presence of Chlamydia nucleic acids in samples (and therefore diagnose infections) and the proteins may be used as antigens for the production of antibodies that may be used to detect Chlamydia proteins in samples (e.g. via enzyme linked immunosorbant assay (ELISA)).

Dwg.0/0

FS CPI

FA AB; DCN

MC CPI: B04-B03C; B04-B04C; B04-B04C1; B04-B04C7

B04-B04M; B04-C01G; B04-E03F; B04-E04; B04-E05; B04-E06; B04-E08; B04-F01; B04-G07; B04-N03A0E; B11-A; B11-C07A4; B11-C08E1; B11-C08E3; B11-C08E5; B12-K04A4; B12-K04F; B14-A01A; B14-C03; B14-K01A; B14-S03; B14-S11B; D05-A01A4; D05-A01B; D05-C12; D05-H04; D05-H07; D05-H08; D05-H09; D05-H11; D05-H12A; D05-H12D1; D05-H12D2; D05-H12D5; D05-H12E; D05-H14; D05-H17A5; D05-H18B

L54 ANSWER 20 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2000-303789 [26] WPIX

DNC C2000-092308

TI Nucleic acid molecule for producing recombinant high molecular weight proteins of *Haemophilus* which are used as a vaccine to provide protection against *Haemophilus* induced diseases in humans.

DC B04 D16

IN KLEIN, M H; LOOSMORE, S M; YANG, Y

PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD; (KLEI-I)

KLEIN M H; (LOOS-I) LOOSMORE S M; (YANG-I) YANG Y

CYC 90

PI WO 2000020609 A2 20000413 (200026)\* EN 307 C12N015-70

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ TZ UG ZW

W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES

FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS  
 LT LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ  
 TM TR TT UA UG US UZ VN YU ZA ZW  
 AU 9960736 A 20000426 (200036)  
 EP 1117807 A2 20010725 (200143) EN C12N015-70  
 R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
 RO SE SI

US 6432669 B1 20020813 (200255) C12P021-06  
 JP 2002532062 W 20021002 (200279) 307 C12N015-09  
 US 2003133943 A1 20030717 (200348) A61K039-02 <--  
 AU 765339 B 20030918 (200370) C12N015-70  
 NZ 511360 A 20040130 (200414) C12N015-70  
 JP 2004166710 A 20040617 (200440) 90 C12N015-09

ADT WO 2000020609 A2 WO 1999-CA938 19991007; AU 9960736 A AU 1999-60736  
 19991007; EP 1117807 A2 EP 1999-947153 19991007; WO 1999-CA938 19991007;  
 US 6432669 B1 CIP of US 1998-167568 19981007, US 1998-206942 19981208; JP  
 2002532062 W WO 1999-CA938 19991007, JP 2000-574704 19991007; US  
 2003133943 A1 Cont of US 1998-167568 19981007, US 2002-193764 20021211; AU  
 765339 B AU 1999-60736 19991007; NZ 511360 A NZ 1999-511360 19991007, WO  
 1999-CA938 19991007; JP 2004166710 A Div ex JP 2000-574704 19991007, JP  
 2004-37346 20040213

FDT AU 9960736 A Based on WO 2000020609; EP 1117807 A2 Based on WO 2000020609;  
 JP 2002532062 W Based on WO 2000020609; AU 765339 B Previous Publ. AU  
 9960736, Based on WO 2000020609; NZ 511360 A Based on WO 2000020609

PRAI US 1998-206942 19981208; US 1998-167568 19981007;  
 US 2002-193764 20021211

IC ICM A61K039-02; C12N015-09; C12N015-70; C12P021-06  
 ICS A61K009-127; A61K009-14; A61K009-48; A61K031-70; A61K038-16;  
 A61K039-102; A61P031-04; C07H021-04; C07K014-195;  
 C07K014-285; C12N001-12; C12N001-21; C12N015-31; C12N015-74;  
 C12P021-02

ICI C12P021-02; C12R001-93

AB WO 200020609 A UPAB: 20000531

NOVELTY - A nucleic acid molecule (I) comprising a promoter functional in  
 Escherichia coli and operatively coupled to a modified operon of a  
 non-typeable strain of Haemophilus comprising A, B and C genes, where the  
 A gene only contains a nucleic acid sequence encoding a mature high  
 molecular weight protein (HMW) of the non-typeable strain of Haemophilus,  
 is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the  
 following:

- (1) a vector adapted for transformation of a host comprising (I);
- (2) a strain of E. coli transformed by the vector of (1) expressing a  
 protective HMW protein of a non-typeable strain of Haemophilus;
- (3) a recombinant protective HMW protein of a non-typeable strain of  
 Haemophilus or immunogenic fragment or analog, produced by the transformed  
 E. coli strain of (2);
- (4) a plasmid vector (II) for expression of a HMW protein of a  
 non-typeable strain of Haemophilus comprising the T7 promoter, a cloning  
 site for insertion of a nucleic acid molecule into the plasmid vector and  
 portions B and C of the operon of a non-typeable Haemophilus strain;
- (5) an isolated and purified HMW 1 protein of a non-typeable strain  
 of Haemophilus free from contamination by HMW 2 of the same strain of  
 non-typeable Haemophilus;
- (6) an isolated and purified HMW 2 protein of a non-typeable strain  
 of Haemophilus free from contamination by HMW 1 of the same strain of  
 non-typeable Haemophilus;
- (7) an immunogenic composition comprising at least one  
 immunogenically-active component which is (I), the recombinant protective  
 HMW protein of (3) or the HMW proteins of (5) or (6) and a carrier;
- (8) a method for inducing protection against disease caused by  
 Haemophilus comprising administering to a susceptible host the composition  
 of (7); and
- (9) a method for producing a protective HMW protein of a non-typeable  
 strain of Haemophilus comprising transforming E. coli with the vector of  
 (1), growing the E. coli to express the encoded mature HMW protein and  
 isolating and purifying the expressed HMW protein.

ACTIVITY - Antibacterial.

Groups of 8-9 chinchillas were immunized three times intramuscularly  
 with 30 micro g of purified rHMW1 or rHMW2, 2 x 10<sup>9</sup> colony forming units  
 (cfu) of heat inactivated (56 deg. C for 10 minutes) H. influenzae (NTHi)  
 whole cells in alum or alum alone on days 0, 14 and 28. Serum samples and  
 nasal wash samples were taken on day 42 for measurement of anti-HMW1 or  
 anti-HMW2 antibody titers by ELISAs (enzyme linked immunosorbent assays).  
 On day 44, animals were lightly anesthetized using xylazine/ketamine  
 hydrochloric acid by intramuscular injection. Intranasal inoculations were

performed by passive inhalation (0.1 ml per animal) of freshly cultured streptomycin-resistant NTHi strain 12 in BHI (not defined) medium supplemented with hemin and nicotinamide adenine dinucleotide (NAD) both at 2 micro g ml<sup>-1</sup>. Dose of bacterial challenge was 1 x 10<sup>8</sup> cfu per animal. Nasopharyngeal lavages were performed 4 days post inoculation on chinchillas. 67-88% of control animals immunized with alum only had culture positive nasal lavage fluids but 67-80% of animals immunized with the rHMW1 protein purified from constructs abc (pDS-1046-1-1), a/abc (pBK86-1-1) or abc/cer (pBK-76-1-1) were largely protected. Animals immunized with constructs that did not contain intact ABC genes were 70-90% infected. Similar results were achieved with rHMW2 protein.

#### MECHANISM OF ACTION - Vaccine.

USE - The nucleic acids and vectors are used for the production of recombinant H. influenzae HMW proteins which can be used as vaccines to mediate a humoral or cell-mediated immune response to provide protection against H. influenzae induced diseases in humans.

The HMW proteins are also useful as antigens in immunoassays for detecting antibacterial, Haemophilus, HMW and/or peptide antibodies. The nucleotide sequences encoding the HMW proteins can be used to isolate and clone hmw genes from other non-typeable strains of Haemophilus in hybridization reactions.

ADVANTAGE - Including the cer gene of E. coli enhances the level of expression of mature HMW protein by the vectors.

Dwg.0/235

FS CPI

FA AB; DCN

MC CPI: B04-E03F; B04-E08; B04-F10A3E; B04-N03A0E; **B14-A01A;**  
**B14-S11B;** D05-H07; D05-H12A; D05-H12E; D05-H14A1; D05-H17A6

L54 ANSWER 21 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2000-224701 [19] WPIX

DNC C2000-068762

TI Nucleic acid molecule encoding an inclusion membrane protein C of a strain of Chlamydia, useful as a vaccine for immunizing against Chlamydia infection.

DC B04 D16

IN DUNN, P L; MURDIN, A D; OOMEN, R P

PA (CONN-N) CONNAUGHT LAB LTD; (DUNN-I) DUNN P L; (MURD-I) MURDIN A D;  
(OOMEN-I) OOMEN R P; (AVET) AVENTIS PASTEUR LTD

CYC 89

PI WO 2000011181 A1 20000302 (200019)\* EN 62 C12N015-31  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ UG ZW  
W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES  
FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS  
LT LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ  
TM TR TT UA UG US UZ VN YU ZA ZW  
AU 9953660 A 20000314 (200031) C12N015-31  
EP 1105490 A1 20010613 (200134) EN C12N015-31  
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

US 6521745 B1 20030218 (200317) C07H021-04

US 6686339 B1 20040203 (200413) A61K048-00

US 2004228874 A1 20041118 (200477) C12Q001-68

ADT WO 2000011181 A1 WO 1999-CA766 19990819; AU 9953660 A AU 1999-53660  
19990819; EP 1105490 A1 EP 1999-939280 19990819; WO 1999-CA766 19990819;  
US 6521745 B1 Provisional US 1998-97199P 19980820, Provisional US  
1999-132961P 19990507, US 1999-377399 19990820; US 6686339 B1 Provisional  
US 1998-97199P 19980820, Provisional US 1999-132961P 19990507, WO  
1999-CA766 19990819, US 2001-763063 20010615; US 2004228874 A1 Provisional  
US 1998-97199P 19980820, Provisional US 1999-132961P 19990507, Div ex WO  
1999-CA766 19990819, Div ex US 2001-763063 20010615, US 2004-756320  
20040114

FDT AU 9953660 A Based on WO 2000011181; EP 1105490 A1 Based on WO 2000011181;  
US 6686339 B1 Based on WO 2000011181; US 2004228874 A1 Div ex US 6686339

PRAI US 1999-132961P 19990507; US 1998-97199P 19980820;  
US 1999-377399 19990820; US 2001-763063 20010615;  
US 2004-756320 20040114

IC ICM A61K048-00; C07H021-04; C12N015-31; C12Q001-68

ICS A61K035-66; A61K039-02; A61K039-118; C07K014-295;  
C07K016-12; C12N015-63

AB WO 200011181 A UPAB: 20000419

NOVELTY - An isolated and purified nucleic acid molecule (750 base pairs (bp)) (I) encoding an inclusion membrane protein C (203 amino acids) (II) of a strain of Chlamydia (both sequences given in the specification), is new.



DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:

- (1) an expression cassette containing (I);
  - (2) an expression vector containing the expression cassette of (1);
- and
- (3) a vaccine vector containing (I).

#### ACTIVITY - Antibacterial.

Groups of 7 to 9 week old male Balb/c mice (8 to 10 per group) were immunized intramuscularly (i.m.) and intranasally (i.n.) with plasmid DNA containing (I). Saline was given to groups of control animals. For i.m. immunization, alternate left and right quadriceps were injected with 100 micro g of DNA in 50 micro l of phosphate buffered saline (PBS) on three occasions at 0, 3 and 6 weeks. For i.n. immunization, anaesthetized mice aspirated 50 micro l of PBS containing 50 micro g DNA on three occasions at 0, 3 and 6 weeks. At week 8, immunized mice were inoculated i.n. with 5 multiply 10 inclusion forming units (IFU) of Chlamydia pneumoniae, strain AR39 in 100 micro l of buffer to test their ability to limit the growth of a sublethal C. pneumoniae challenge. Lungs were taken from mice at days 5 and 9 post-challenge and homogenized. The homogenate was frozen at -70 deg. C until assay. Dilutions of the homogenate were assayed for the presence of infectious Chlamydia by inoculation into monolayers of susceptible cells. The inoculum was centrifuged onto the cells at 300 revolutions per minute (rpm) for 1 hour, the cells were then incubated for three days at 35 deg. C in the presence of 1 micro g/ml cycloheximide. After incubation, the monolayers were fixed with formalin and methanol then immunoperoxidase stained for the presence of chlamydial inclusions using convalescent sera from rabbits infected with C. pneumoniae. Mice immunized i.n. and i.m. with pCAI115 had chlamydial lung titers less than 262500 in 4 of the 4 cases at day 5 and less than 250000 in 4 of the 4 cases at day 9. In contrast, mice sham immunized with saline had 202400 to 886800 IFU/lung (mean 429800) at day 5 and 78400-284600 IFU/lung (mean 157080) at day 9.

#### MECHANISM OF ACTION - Vaccine.

USE - (I) is useful for protecting against Chlamydia infection.

Dwg.0/4

FS CPI

FA AB; DCN

MC CPI: B04-B03C; B04-C01G; B04-E03; B04-E04; B04-E05; B04-E08;  
B04-G01; B04-N03A; B11-C08D2; B11-C08E; B12-K04A4;  
B14-A01A; B14-S11B; D05-H07; D05-H09; D05-H11;  
D05-H12A; D05-H12D1; D05-H12D5; D05-H12E; D05-H17A

L54 ANSWER 22 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2000-224700 [19] WPIX

DNC C2000-068761

TI New nucleic acid encoding POMP91A protein from a strain of Chlamydia useful for preventing, treating and diagnosing Chlamydia infection.

DC B04 D16

IN DUNN, P L; MURDIN, A D; OOMEN, R P

PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD

CYC 89

PI WO 2000011180 A1 20000302 (200019)\* EN 98 C12N015-31

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ UG ZW

W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES  
FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS  
LT LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ  
TM TR TT UA UG US UZ VN YU ZA ZW

AU 9953659 A 20000314 (200031) C12N015-31

EP 1105489 A1 20010613 (200134) EN C12N015-31

R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

US 6693087 B1 20040217 (200413) A61K039-395 <--

ADT WO 2000011180 A1 WO 1999-CA765 19990819; AU 9953659 A AU 1999-53659

19990819; EP 1105489 A1 EP 1999-939279 19990819; WO 1999-CA765 19990819;

US 6693087 B1 Provisional US 1998-97198P 19980820, US 1999-377850 19990820

FDT AU 9953659 A Based on WO 2000011180; EP 1105489 A1 Based on WO 2000011180

PRAI US 1998-97198P 19980820; US 1999-377850 19990820

IC ICM A61K039-395; C12N015-31

ICS A61K031-70; A61K048-00; C07H021-04; C07K014-295

AB WO 2000011180 A UPAB: 20000419

NOVELTY - Isolated and purified nucleic acid molecule (I) encoding a POMP91A protein or polypeptide fragment of POMP91A from a strain of Chlamydia, is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:

Search done by Noble Jarrell

- (1) an expression cassette containing (I) under the control of elements required for expression of (I);
- (2) an expression vector containing the expression cassette of (1);
- (3) a vaccine vector comprising (I) under the control of elements required for expression of (I); and
- (4) an antibody that specifically binds to a polypeptide with a protein sequence (II) of 947 amino acids or a polypeptide fragment containing the binding domain of (II).

ACTIVITY - Antibacterial.

MECHANISM OF ACTION - Vaccine.

Male Balb/c mice (7-9 weeks old) were immunized intramuscularly (100 micro g DNA in 50 micro l phosphate buffered saline) and intranasally (50 micro g DNA in 50 micro l phosphate buffered saline) with plasmid DNA pCAI327 containing the coding sequence of C. pneumonia POMP91A at 0, 3 and 6 week intervals. Control animals were given saline and plasmids pCAI116 and pCAI178 which express non-protective chlamydial antigens. After 8 weeks the immunized mice were inoculated intranasally with 5 x 10<sup>5</sup> IFU of C. pneumonia strain AR39 in 100 micro l SPG (sucrose, phosphate, glutamate) buffer. The lungs were taken from the mice at days 5 and 9 post challenge and homogenized in SPG buffer (7.5% sucrose, 5 mM glutamate, 12.5 mM phosphate pH 7.5). Dilutions of the homogenate were assayed for the presence of infectious Chlamydia by inoculation onto monolayers of susceptible cells. The cells were incubated for 3 days at 35 deg. C in the presence of 1 micro g/ml cycloheximide and then fixed with formalin and methanol then immunoperoxidase stained using convalescent sera from rabbits infected with C. pneumonia and metal enhanced DAB (not defined) as peroxidase substrate. Mice immunized with pCAI327 had chlamydial lung titers less than 21500 in 5 of 6 cases at day 9 but for saline immunized mice the average titer was 49069 IFU/lung.

USE - (I) is used to prevent, treat and diagnose Chlamydia infection.

Vaccine vectors containing (I) are used to induce an immune response against Chlamydia. (I) or a monoclonal antibody specific to POMP91A can be used to diagnose the presence of Chlamydia in a biological sample.

Dwg.0/41

FS CPI

FA AB; DCN

MC CPI: B04-C01G; B04-E03F; B04-E08; B04-F0100E; B04-F10A; B04-G01; B04-N03A; B11-C07A; B11-C08E5; B12-K04A4; B12-K04F; B14-A01A; B14-S11B; D05-H09; D05-H11A; D05-H12A; D05-H12E; D05-H14; D05-H17A6

L54 ANSWER 23 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2000-096387 [08] WPIX

CR 1995-194089 [25]; 1997-052329 [05]; 1998-100410 [09]; 1999-404437 [34]; 1999-404459 [34]; 1999-404487 [34]; 2000-181144 [16]

DNC C2000-027984

TI Antibodies specific for transferrin receptor proteins of Haemophilus influenzae, useful for treating otitis media, epiglottitis, pneumonia and tracheobronchitis.

DC B04 D16

IN CHONG, P; GRAY-OWEN, S; HARKNESS, R; KLEIN, M; LOOSMORE, S; MURDIN, A; SCHRYVERS, A; YANG, Y

PA (CONN-N) CONNAUGHT LAB LTD

CYC 1

PI US 6008326 A 19991228 (200008)\* 252 C07K016-12

ADT US 6008326 A CIP of US 1993-148968 19931108, CIP of US 1993-175116 19931229, US 1995-474671 19950607, Cont of US 1995-337483 19951108

PRAI US 1995-337483 19951108; US 1993-148968 19931108; US 1993-175116 19931229; US 1995-474671 19950607

IC ICM C07K016-12

ICS C07K016-28

AB US 6008326 A UPAB: 20000925

NOVELTY - An isolated and purified antibody (or monospecific antiserum) specific for a single transferrin receptor protein (or immunogenic fragment) of a strain of Haemophilus influenzae, is new.

ACTIVITY - Antibacterial; antiinflammatory; auditory; respiratory.

No relevant biological data given.

MECHANISM OF ACTION - Vaccine (antibody inhibition of bacterial growth and replication).

USE - The antibodies may be used for preventing and treating infections and disorders caused by H. influenzae, these include bacterial meningitis, otitis media, epiglottitis, pneumonia and tracheobronchitis. The antibodies may also be used to detect the presence of H. influenzae proteins in samples according to standard methodologies (e.g. enzyme linked immunosorbant assay (ELISA)) and hence diagnose infections.

ADVANTAGE - The use of antibodies to treat bacterial infections

avoids the risk of antibiotic resistant bacterial strains developing. In the treatment of otitis media, the use of antibodies avoids the need for extensive and costly surgery (e.g. tonsillectomies, adenoidectomies and the insertion of tympanostomy tubes) to rectify hearing problems.

Dwg.0/30

FS CPI

FA AB; DCN

MC CPI: B04-B04C1; B04-B04C7; B04-C01; B04-E03D;  
B04-F0100E; B04-G04; B04-G07; B04-G21; B04-K01T; B04-N0300E; B11-A;  
B11-C07A1; B11-C07A4; B11-C08E; B11-C09; B12-K04A4; B12-M05;  
B14-A01A; B14-C03; B14-K01; B14-N02; B14-N05;  
B14-S11B; D05-A01A4; D05-A01B; D05-C12; D05-H04; D05-H07;  
D05-H09; D05-H11A; D05-H12A; D05-H14; D05-H17A4; D05-H17A5

L54 ANSWER 24 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 1999-620376 [53] WPIX

CR 1997-457533 [42]

DNC C1999-181129

TI Nucleic acid encoding transferrin binding protein 2 of *Moraxella* catarrhalis, useful for diagnostics, immunization and recombinant protein production.

DC B04 D16

IN DU, R; HARKNESS, R E; KLEIN, M H; LOOSMORE, S M;

MYERS, L E; SCHRYVERS, A B; YANG, Y

PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD

CYC 83

PI WO 9952947 A2 19991021 (199953)\* EN 113 C07K014-79  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ UG ZW  
W: AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GE  
GH GM HR HU ID IL IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD MG  
MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT UA UG  
US UZ VN YU ZW

AU 9931350 A 19991101 (200013)

EP 1071715 A2 20010131 (200108) EN C07K014-79

R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI

BR 9909576 A 20011016 (200170) C07K014-79

JP 2002511490 W 20020416 (200242) 122 C07K014-705

US 6440701 B1 20020827 (200259) C12N015-31

AU 761008 B 20030529 (200346) C07K014-79

NZ 507978 A 20030725 (200357) C07K014-79

ADT WO 9952947 A2 WO 1999-CA307 19990412; AU 9931350 A AU 1999-31350 19990412;  
EP 1071715 A2 EP 1999-913049 19990412; WO 1999-CA307 19990412; BR 9909576  
A BR 1999-9576 19990412; WO 1999-CA307 19990412; JP 2002511490 W WO  
1999-CA307 19990412; JP 2000-543503 19990412; US 6440701 B1 CIP of US  
1996-613009 19960308, CIP of US 1997-778570 19970103, CIP of WO 1997-CA163  
19970307, US 1998-59584 19980414; AU 761008 B AU 1999-31350 19990412; NZ  
507978 A NZ 1999-507978 19990412; WO 1999-CA307 19990412

FDT AU 9931350 A Based on WO 9952947; EP 1071715 A2 Based on WO 9952947; BR  
9909576 A Based on WO 9952947; JP 2002511490 W Based on WO 9952947; AU  
761008 B Previous Publ. AU 9931350, Based on WO 9952947; NZ 507978 A Based  
on WO 9952947

PRAI US 1998-59584 19980414; US 1996-613009 19960308;

US 1997-778570 19970103; WO 1997-CA163 19970307

IC ICM C07K014-705; C07K014-79; C12N015-31

ICS A61K039-02; A61K048-00; C07K014-22; C12N001-15; C12N001-19;

C12N001-21; C12N005-10; C12N015-09; C12N015-63; C12P021-02;

C12Q001-68

ICI C07K014-705; C12N015-09; C12R001:01

AB WO 9952947 A UPAB: 20030906

NOVELTY - Purified, isolated nucleic acid (I) encoding a transferrin binding protein (Tbp2) (II) from *Moraxella* catarrhalis strains M35, 3 or LES1, is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:

(a) vectors containing (I);

(b) transformed host cells containing the vector of (a);

(c) recombinant production of (II);

(d) recombinant (II) produced this way;

(e) an immunogenic composition containing (I) or recombinant (II)

plus a carrier;

(f) a method for detecting *Moraxella* nucleic acid that

encodes transferrin receptor protein by the formation of a hybrid with (I); and

(g) diagnostic kits for the method of (f).

ACTIVITY - Antibacterial; cytostatic; auditory.

MECHANISM OF ACTION - Tbp binding blocker.

(I) and (II) generate an immune response that includes anti-Tbp antibodies and opsonizing and/or bactericidal antibodies. By blocking binding to Tbp, the antibodies stop the bacterium from acquiring essential iron.

USE - (I) is used to produce recombinant (II); for identification or diagnosis of *Moraxella*, or for cloning related species, using hybridization assays; and for genetic immunization against *Moraxella* infections, e.g. otitis media. (II) are useful as antigens, either in vaccines (including components of conjugate vaccines that contain antigens from other bacteria or from tumors, in which case they elicit production of antitumor antibodies that may be coupled to chemotherapeutic agents or biologically active agents) or to raise antibodies (for use as diagnostic reagents and for treating *Moraxella* infections), also for detecting *Moraxella* antibodies.

Dwg.0/9

FS CPI

FA AB; DCN

MC CPI: B04-C01G; B04-E02F; B04-E05; B04-E08; B04-F0100E; B04-G01; B04-K01T; B11-C07A; B11-C07A1; B11-C08E5; B12-K04A; B12-K04E; B12-K04F; B14-A01; B14-H01; B14-N02; D05-H04; D05-H07; D05-H09; D05-H11; D05-H12A; D05-H12E; D05-H14; D05-H17A4; D05-H17A6; D05-H18B

LS4 ANSWER 25 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 1997-373222 [35] WPIX

DNN N1997-309929 DNC C1997-120314

TI Lactoferrin receptor protein isolated from bacterial pathogen - used as, e.g. vaccine, carrier for antigens and immunogens and diagnostic agents.

DC A96 B04 D16 S03

IN BONNAH, R A; SCHRYVERS, A B

PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD

CYC 2

PI CA 2162193 A 19970503 (199735)\* 50 C12P021-08  
US 6048539 A 20000411 (200025) A61K039-095 <--  
US 6211343 B1 20010403 (200120) C08H001-00  
US 6344200 B1 20020205 (200211) A61K039-02 <--  
US 6348198 B1 20020219 (200221) A61K039-095 <--

ADT CA 2162193 A CA 1995-2162193 19951106; US 6048539 A US 1995-552232 19951102; US 6211343 B1 Div ex US 1995-552232 19951102, US 1999-370869 19990810; US 6344200 B1 Div ex US 1995-552232 19951102, US 1999-371126 19990810; US 6348198 B1 Div ex US 1995-552232 19951102, US 1999-371127 19990810

FDT US 6211343 B1 Div ex US 6048539; US 6344200 B1 Div ex US 6048539; US 6348198 B1 Div ex US 6048539

PRAI US 1995-552232 19951102; US 1999-370869 19990810; US 1999-371126 19990810; US 1999-371127 19990810

IC ICM A61K039-02; A61K039-095; C08H001-00; C12P021-08  
ICS C07K001-36; C07K014-22; C07K016-12; G01N033-566; G01N033-569

AB CA 2162193 A UPAB: 19970828

Lactoferrin receptor protein (I) is isolated and purified from a bacterial protein and has molecular weight (MW) 70-90 kDa as determined by SDS-PAGE.

USE - The immunogenic composition can be used as a vaccine to a bacterial pathogen selected from *Neisseria meningitidis*, *N-gonorrhoeae*, *Moraxella catarrhalis*, *M. movis* and *M. lacunata* (all claimed).

The proteins can be used in the diagnosis of and vaccination against diseases caused by bacterial pathogens that produce lactoferrin receptor proteins or proteins capable of raising antibodies reactive with lactoferrin receptor proteins. The proteins can be used as antigens, immunogenic preparations including vaccines, carriers for other antigens and immunogens and the generation of diagnostic reagents.

The bacterial pathogen may also be selected from *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Escherichia coli*, *Salmonella typhi*, *Streptococcus mutans*, *Cryptococcus neoformans*, *Klebsiella* sp., *Staphylococcus aureus* and *Pseudomonas aeruginosa*.

(I; Lbp2) may also be used to induce immunity toward abnormal polysaccharides of tumour cells and to produce antitumour antibodies that can be conjugated to chemotherapeutic and bioactive agents.

Dwg.0/4

FS CPI EPI

FA AB

MC CPI: A12-V01; A12-V03C2; B04-G01; B04-N03; B11-C07A; B12-K04A; B14-S11B; D05-H07; D05-H09; D05-H13  
EPI: S03-E14H4

L54 ANSWER 26 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN  
 AN 1996-342057 [34] WPIX  
 DNN N1996-287919 DNC C1996-108616  
 TI New immunogenic conjugate molecules - comprise a capsular polysaccharide of Streptococcus linked to an outer membrane protein of Haemophilus.  
 DC B04 D16 S03  
 IN FAHIM, R E F; GISONNI, L; KANDIL, A; KLEIN, M H; YANG, Y  
 ; FAHIM, R E  
 PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD  
 CYC 71  
 PI WO 9621465 A2 19960718 (199634)\* EN 64 A61K039-09 <--  
 RW: AT BE CH DE DK EA ES FR GB GR IE IT KE LS LU MC MW NL OA PT SD SE  
 SZ UG  
 W: AL AM AT AU AZ BB BG BR BY CA CH CN CZ DE DK EE ES FI GB GE HU IS  
 JP KE KG KP KR KZ LK LR LS LT LU LV MD MG MK MN MW MX NO NZ PL PT  
 RO RU SD SE SG SI SK TJ TM TR TT UA UG US UZ VN  
 AU 9643254 A 19960731 (199645) A61K039-09 <--  
 WO 9621465 A3 19961010 (199648) A61K039-09 <--  
 US 5681570 A 19971028 (199749) 21 A61K039-385 <--  
 EP 805691 A1 19971112 (199750) EN A61K039-09 <--  
 R: AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL PT SE  
 US 6177085 B1 20010123 (200107) A61K039-09 <--  
 US 6329512 B1 20011211 (200204) A23J001-00  
 ADT WO 9621465 A2 WO 1996-CA7 19960105; AU 9643254 A AU 1996-43254 19960105;  
 WO 9621465 A3 WO 1996-CA7 19960105; US 5681570 A US 1995-371965 19950112;  
 EP 805691 A1 EP 1996-900066 19960105; WO 1996-CA7 19960105; US 6177085 B1  
 Cont of US 1995-371965 19950112, US 1995-467884 19950606; US 6329512 B1  
 Cont of US 1995-371965 19950112, US 1995-467883 19950606  
 FDT AU 9643254 A Based on WO 9621465; EP 805691 A1 Based on WO 9621465; US  
 6177085 B1 Cont of US 5681570; US 6329512 B1 Cont of US 5681570  
 PRAI US 1995-371965 19950112; US 1995-467884 19950606;  
 US 1995-467883 19950606  
 REP 1.Jnl.Ref; EP 172107; EP 338265; EP 389925; EP 497524; EP 497525; US  
 4673574; US 5098997; WO 9106652  
 IC ICM A23J001-00; A61K039-09; A61K039-385  
 ICS A61K039-02; A61K039-102; A61K039-116;  
 C07K014-285; C07K017-10; G01N033-569  
 AB WO 9621465 A UPAB: 19960829  
 The following are claimed: (A) an immunogenic conjugate molecule (ICM) comprising at least a portion of a capsular polysaccharide (CP) of a Streptococcus strain linked to at least a portion of an outer membrane protein (OMP) of a Haemophilus strain, which are selected to provide in the ICM an enhanced immune response to the CP; (B) a diagnostic kit for determining the presence of antibodies in a sample specifically reactive with a CP of a Streptococcus strain or with an OMP of a Haemophilus strain; (C) a diagnostic kit for detecting the presence of a CP of a Streptococcus strain in a sample; (D) a diagnostic kit for detecting the presence of an OMP of a Haemophilus strain in a sample; (E) a process for individually isolating P1, P2 and P6 OMPs from a Haemophilus strain; and (F) an ICM comprising a P6 OMP of a Haemophilus strain linked to at least a portion of a CP of an encapsulated pathogen to provide in the ICM an enhanced immune response to the CP.  
 The ICM is pref. used with an adjuvant eg. AlPO<sub>4</sub>, Al(OH)<sub>3</sub>, QIL A, QS21, Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>, Ca(OH)<sub>2</sub>, Zn(OH)<sub>2</sub> a glycolipid analogue or an octadecyl ester of an amino acid. The OMPs of Haemophilus strains may be purified using eg. DEAE-Sephacel (RTM) and hydroxyapatite columns.  
 USE - The ICMs can be used in vaccines to confer protection against disease caused by the Streptococcus strain and the Haemophilus strain (claimed). The ICMs and antibodies produced using the ICMs can also be used in diagnostic procedures and kits.  
 ADVANTAGE - The ICMs provide an enhanced immune response to CPs of Streptococcus strains without the necessity to employ carrier proteins. They also provide an immune response to the OMP of the Haemophilus strain.  
 Dwg.0/8  
 FS CPI EPI  
 FA AB  
 MC CPI: B04-C02F; B04-G07; B04-N03; B11-C07A; B12-K04A4; B14-A01A;  
 B14-A01B2; B14-S11B; D05-H04; D05-H07; D05-H10  
 EPI: S03-E14H4

L54 ANSWER 27 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN  
 AN 1994-200269 [24] WPIX  
 DNC C1994-091569  
 TI Nucleic acid encoding D15 outer membrane protein - especially of Haemophilus influenzae, and related proteins, vectors, antisera etc. useful in vaccines, for diagnosis and for passive immunisation..

DC B04 D16  
IN CHONG, P; KLEIN, M; LOOSMORE, S; SIA, D Y C; THOMAS, W; YANG, Y; LOOSMORE, S M; SIA, D; YANG, Y P  
PA (CONN-N) CONNAUGHT LAB LTD; (AVET) AVENTIS PASTEUR LTD  
CYC 28  
PI WO 9412641 A1 19940609 (199424)\* 161 C12N015-31  
RW: AT BE CH DE DK ES FR GB GR IE IT LU MC NL PT SE  
W: AU BR CA FI JP KR NO NZ RU UA US  
AU 9455565 A 19940622 (199436) C12N015-31  
EP 668916 A1 19950830 (199539) EN C12N015-31  
R: AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL PT SE  
JP 08502417 W 19960319 (199644) 156 C12N015-09  
AU 683435 B 19971113 (199803) C12N015-31  
BR 9307510 A 19990601 (199927) C12N015-31  
JP 2907552 B2 19990621 (199930) 181 C12N015-09  
US 6013514 A 20000111 (200010) A61K039-102 <--  
US 6083743 A 20000704 (200036) A61K039-102 <--  
RU 2141528 C1 19991120 (200041) C12N015-31  
KR 216390 B1 19990816 (200104) C12N015-31  
US 6264954 B1 20010724 (200146) A61K039-102 <--  
ADT WO 9412641 A1 WO 1993-CA501 19931123; AU 9455565 A AU 1994-55565 19931123; EP 668916 A1 WO 1993-CA501 19931123; EP 1994-900671 19931123; JP 08502417 W WO 1993-CA501 19931123; JP 1994-512608 19931123; AU 683435 B AU 1994-55565 19931123; BR 9307510 A BR 1993-7510 19931123; WO 1993-CA501 19931123; JP 2907552 B2 WO 1993-CA501 19931123; JP 1994-512608 19931123; US 6013514 A WO 1993-CA501 19931123; US 1995-433522 19950912; US 6083743 A Cont of WO 1993-CA501 19931123, Cont of US 1995-433522 19950912, US 1998-135166 19980818; RU 2141528 C1 WO 1993-CA501 19931123, RU 1995-117238 19931123; KR 216390 B1 WO 1993-CA501 19931123, KR 1995-702081 19950523; US 6264954 B1 Cont of WO 1993-CA501 19931123, Div ex US 1995-433522 19950912, US 1997-942046 19971001  
FDT AU 9455565 A Based on WO 9412641; EP 668916 A1 Based on WO 9412641; JP 08502417 W Based on WO 9412641; AU 683435 B Previous Publ. AU 9455565, Based on WO 9412641; BR 9307510 A Based on WO 9412641; JP 2907552 B2 Previous Publ. JP 08502417, Based on WO 9412641; US 6013514 A Based on WO 9412641; RU 2141528 C1 Based on WO 9412641  
PRAI GB 1992-24584 19921123  
REP 01Jnl.Ref; EP 281673; EP 378929; US 5013664; WO 9106652  
IC ICM A61K039-102; C12N015-09; C12N015-31  
ICS A61K039-12; A61K039-395; C07H021-04; C07K013-00; C07K014-11; C07K014-195; C07K014-285; C07K016-12; C12N015-62  
ICA C12P021-02; G01N033-569  
ICI C12N015-31, C12R001:21; C12P021-02, C12R001:19; C12N015-31, C12R001:21  
AB WO 9412641 A UPAB: 19940803  
New nucleic acid (I) contains at least a portion coding for a D15 outer membrane protein (omp) and has a sequence which is (a) any of 5 (all about 3000bp) reproduced in the specification, or complementary sequences or (b) hybridisable under stringent conditions with such sequences. Also new are (1) recombinant plasmids containing a segment of (I) at least 18 bp long (and opt. expression control elements, (12) proteins (II) encoded by these plasmids; (3) purified D15 omp (III); (4) synthetic polypeptides with sequences corresp. to (II) or (III), or their variants and mutants which retain immunogenicity; (5) antisera or antibodies specific for (II), (III) or immunologous containing them; (6) chimeric molecules consisting of (II) or (III) bonded to another polypeptides, protein or polysaccharides.  
USE - (I), (II) and the synthetic polypeptides are useful in vaccines to protect against Haemophilus. D15 can also be used as a carrier for polysaccharide antigens to form conjugate vaccines against other bacteria; to induce immunity to abnormal polysaccharides or tumour cells and to generate anti-tumour antibodies, for coupling to toxins etc. (I), (II) synthetic peptides and antisera can also be used diagnostically (in hybridisation or immunoassay procedures) and antibodies can be used for passive immunisation.  
Dwg.0/11  
FS CPI  
FA AB  
MC CPI: B04-C01; B04-C02; B04-E02F; B04-E08; B04-G01; B04-N0300E; B04-N04; B12-K04; B14-A01A; B14-H01B; B14-S11A; B14-S11B; D05-H11; D05-H12A; D05-H12E; D05-H17A; D05-H17C  
L54 ANSWER 28 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN  
AN 1993-134388 [16] WPIX  
DNC C1993-059997  
TI Oligoside derived from antigenic polyoside of a pathogen - useful for treating and preventing e.g. bacterial infections and mycoses.  
DC B04

IN MOREAU, M  
PA (AVET) AVENTIS PASTEUR; (INMR) PASTEUR MERIEUX SERUMS & VACCINS  
SA; (INMR) PASTEUR MERIEUX SERUMS & VACCINS  
CYC 24  
PI WO 9307178 A1 19930415 (199316)\* EN 38 C08B037-00  
RW: AT BE CH DE DK ES FR GB GR IE IT LU MC NL SE  
W: AU CA FI HU JP KR NO US  
FR 2682388 A1 19930416 (199328) 29 C08B037-00  
AU 9229469 A 19930503 (199334)  
FI 9302626 A 19930609 (199334) C08B000-00  
EP 562107 A1 19930929 (199339) FR  
R: AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL SE  
NO 9302102 A 19930805 (199343)  
JP 06506233 W 19940714 (199432) 10 A61K039-02 <--  
AU 661071 B 19950713 (199535) C08B037-00  
HU 70298 T 19950928 (199546)  
US 6007818 A 19991228 (200007) A61K039-00 <--  
NO 306905 B1 20000110 (200009) C08B037-00  
US 6045805 A 20000404 (200024) A61K039-085 <--  
KR 249709 B1 20000315 (200122) C08B037-00  
HU 219672 B 20010628 (200143) C08B037-00  
EP 562107 B1 20020502 (200230) FR C08B037-00  
R: AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL SE  
DE 69232585 E 20020606 (200245) C08B037-00  
ES 2174839 T3 20021116 (200302) C08B037-00  
FI 110164 B1 20021213 (200306) A61K039-02 <--  
CA 2098105 C 20030429 (200337) FR C08B037-00  
ADT WO 9307178 A1 WO 1992-FR955 19921009; FR 2682388 A1 FR 1991-12478  
19911010; AU 9229469 A AU 1992-29469 19921009; FI 9302626 A WO 1992-FR955  
19921009, FI 1993-2626 19930609; EP 562107 A1 EP 1992-923831 19921009, WO  
1992-FR955 19921009; NO 9302102 A WO 1992-FR955 19921009, NO 1993-2102  
19930609; JP 06506233 W WO 1992-FR955 19921009, JP 1993-506690 19921009;  
AU 661071 B AU 1992-29469 19921009; HU 70298 T WO 1992-FR955 19921009, HU  
1993-1682 19921009; US 6007818 A Div ex WO 1992-FR955 19921009, Div ex US  
1993-70446 19931007, US 1995-474190 19950607; NO 306905 B1 WO 1992-FR955  
19921009, NO 1993-2102 19930609; US 6045805 A Cont of WO 1992-FR955  
19921009, Cont of US 1993-70446 19931007, US 1995-474194 19950607; KR  
249709 B1 WO 1992-FR955 19921009, KR 1993-701727 19930609; HU 219672 B WO  
1992-FR955 19921009, HU 1993-1682 19921009; EP 562107 B1 EP 1992-923831  
19921009, WO 1992-FR955 19921009; DE 69232585 E DE 1992-632585 19921009,  
EP 1992-923831 19921009, WO 1992-FR955 19921009; ES 2174839 T3 EP  
1992-923831 19921009; FI 110164 B1 WO 1992-FR955 19921009, FI 1993-2626  
19930609; CA 2098105 C CA 1992-2098105 19921009, WO 1992-FR955 19921009  
FDT AU 9229469 A Based on WO 9307178; EP 562107 A1 Based on WO 9307178; JP  
06506233 W Based on WO 9307178; AU 661071 B Previous Publ. AU 9229469,  
Based on WO 9307178; HU 70298 T Based on WO 9307178; NO 306905 B1 Previous  
Publ. NO 9302102; HU 219672 B Previous Publ. HU 70298, Based on WO  
9307178; EP 562107 B1 Based on WO 9307178; DE 69232585 E Based on EP  
562107, Based on WO 9307178; ES 2174839 T3 Based on EP 562107; FI 110164  
B1 Previous Publ. FI 9302626; CA 2098105 C Based on WO 9307178  
PRAI FR 1991-12478 19911010  
REP 3.Jnl.Ref; BE 1000118; EP 245045; EP 97407  
IC ICM A61K039-00; A61K039-02; A61K039-085;  
C08B000-00; C08B037-00  
ICS A61K031-70; A61K031-715; A61K035-74; A61K038-00; A61K039-08  
; A61K039-09; A61K039-102; A61K039-106;  
A61K039-112; C07H003-06; C07K001-00; C07K014-00  
AB WO 9307178 A UPAB: 19930924  
New oligoside (I), retaining at least one antigenic determinant of an  
antigenic polyside (A) derived from a pathogen, is prepared by (1)  
oxido-reductive depolymerisation of (A); (2) recovering (I) and opt. (3)  
coupling it to a conjugation partner or to a carrier to form a conjugate.  
Pref. (I) have mean elution constant on Sepharose 4BCL of 0.2-1, best  
0.6-0.7 (equivalent to mol.weight 30000-60000, dextran equivalent) and is derived  
from the capsular polysaccharides of a pathogenic Staphylococcus,  
Streptococcus, Klebsiella, Salmonella, Escherichia, Neisseria or  
Haemophilus, especially Salmonella typhi, Strep. pneumoniae, N.meningitidis or  
H. influenzae.  
(I) is pref. conjugated to a peptide, protein or organic polymer,  
most pref. pertussis, cholera, tetanus or diphtheria toxin.4  
Alternatively, (I) is incorporated into a vector which stimulates  
immunogenicity in mammals, or into a liposome.  
USE/ADVANTAGE - (I), especially in conjugated form, are useful in vaccines  
to protect against (or reduce the effects of) bacterial infections or  
mycoses. The usual dose (by any standard route) is 1-200 microg in 0.5ml.  
This method of (I) production produces fragments of homogeneous size which

retain the essential structural determinants; is simple and inexpensive, and can be applied to any polyoside structure.

0/6

FS CPI

FA AB

MC CPI: B02-V02; B04-B02B1; B04-C02F; B12-A01; B12-A02C

L54 ANSWER 29 OF 29 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 1991-178107 [24] WPIX

DNC C1991-076917

TI New mammalian cytokine interleukin-11 - for use in treating immune system disorders, e.g. deficiencies in haematopoietic progenitor or stem cells, and cancer.

DC B04 D16

IN BENNETT, F K; PAUL, S R; YANG, Y; STEPHEN, R P; YANG, Y C

PA (CHIL-N) CHILDRENS MEDICAL CENT; (GEMY) GENETICS INST INC; (CHIL-N)

CHILDRENS MED CENTER CORO

CYC 21

PI WO 9107495 A 19910530 (199124)\*

RW: AT BE CH DE DK ES FR GB GR IT LU NL SE

W: AU CA HU JP KR US

AU 9067578 A 19910613 (199137)

EP 504177 A1 19920923 (199239) EN 69 C12N015-24

R: AT BE CH DE DK ES FR GB GR IT LI LU NL SE

DK 9200666 A 19920722 (199242) C07K013-00

US 5215895 A 19930601 (199323) 17 C12P021-02

JP 05504560 W 19930715 (199333) 69 C07K013-00

AU 644389 B 19931209 (199405) C07K013-00

HU 64595 T 19940128 (199409) C12N015-24

US 5371193 A 19941206 (199503) 18 C07K013-00

JP 2688539 B2 19971210 (199803) 20 C07K014-54

US 5700664 A 19971223 (199806) 20 C12N015-24

JP 10004982 A 19980113 (199812) 21 C12N015-09

MX 184567 B 19970430 (199821) C12N015-024

JP 2783361 B2 19980806 (199836) 21 C12N015-09

HU 215233 B 19981130 (199903) C12N015-24

US 5854028 A 19981229 (199908) C12N015-63

KR 9705050 B1 19970411 (199938) C12N015-24

US 6066317 A 20000523 (200032) A61K038-19

EP 504177 B1 20010207 (200109) EN C12N015-24

R: AT BE CH DE DK ES FR GB GR IT LI LU NL SE

DE 69033700 E 20010315 (200122) C12N015-24

CA 2069428 C 20010731 (200147) EN C12N015-24

ES 2156852 T3 20010801 (200149) C12N015-24

ADT EP 504177 A1 EP 1990-917548 19901120, WO 1990-US6803 19901120; DK 9200666

A WO 1990-US6803 19901120, DK 1992-666 19920521; US 5215895 A CIP of US

1989-441100 19891122, US 1990-526474 19900521; JP 05504560 W WO

1990-US6803 19901120, JP 1991-500597 19901120; AU 644389 B AU 1990-67578

19901120; HU 64595 T WO 1990-US6803 19901120, HU 1992-1712 19901120; US

5371193 A Div ex US 1990-526474 19900521, US 1993-17522 19930212; JP

2688539 B2 WO 1990-US6803 19901120, JP 1991-500597 19901120; US 5700664 A

CIP of US 1989-441100 19891122, CIP of US 1990-526474 19900521, WO

1990-US6803 19901120, US 1992-949516 19921119; JP 10004982 A Div ex JP

1991-500597 19901120, JP 1997-51468 19901120; MX 184567 B MX 1992-3439

19920626; JP 2783361 B2 Div ex JP 1991-500597 19901120, JP 1997-51468

19901120; HU 215233 B WO 1990-US6803 19901120, HU 1992-1712 19901120; US

5854028 A CIP of US 1989-441100 19891122, CIP of US 1990-526474 19900521,

Div ex US 1992-949516 19921119, US 1997-814459 19970310; KR 9705050 B1 WO

1990-US6803 19901120, KR 1992-701213 19920522; US 6066317 A CIP of US

1989-441100 19891122, Cont of US 1990-526474 19900521, Div ex US

1992-949516 19921119, Cont of US 1997-814459 19970310, US 1998-122525

19980724; EP 504177 B1 EP 1990-917548 19901120, WO 1990-US6803 19901120;

DE 69033700 E DE 1990-633700 19901120, EP 1990-917548 19901120, WO

1990-US6803 19901120; CA 2069428 C CA 1990-2069428 19901120, WO

1990-US6803 19901120; ES 2156852 T3 EP 1990-917548 19901120

FDT EP 504177 A1 Based on WO 9107495; JP 05504560 W Based on WO 9107495; AU 644389 B Previous Publ. AU 9067578, Based on WO 9107495; HU 64595 T Based on WO 9107495; US 5371193 A Div ex US 5215895; JP 2688539 B2 Previous Publ. JP 05504560, Based on WO 9107495; US 5700664 A CIP of US 5215895, Based on WO 9107495; JP 2783361 B2 Previous Publ. JP 10004982; HU 215233 B Previous Publ. HU 64595, Based on WO 9107495; US 5854028 A CIP of US 5215895, Div ex US 5700664; US 6066317 A Cont of US 5215895, Div ex US 5700664, Cont of US 5854028; EP 504177 B1 Based on WO 9107495; DE 69033700 E Based on EP 504177, Based on WO 9107495; CA 2069428 C Based on WO 9107495; ES 2156852 T3 Based on EP 504177

PRAI WO 1990-US6803 19901120; US 1989-441100 19891122;



WO 1990-US6803U 19901120; US 1993-17522 19930212;  
 US 1992-949516 19921119; US 1997-814459 19970310;  
 US 1998-122525 19980724

REP 2.Jnl.Ref

IC ICM A61K038-19; C07K013-00; C12N015-024; C12N015-09; C12N015-24;  
 C12N015-63; C12P021-02

ICS A61K037-02; A61K037-66; A61K038-20; A61K039-395;  
 C07H015-12; C07K013-000; C07K014-00; C07K015-06; C12N001-21;  
 C12N015-19; C12P021-002

ICA A61K038-00; C07H021-04; C07K014-54; C12N005-10

ICI A61K038-00; A61K037-02; A61K037-66; C12P021-02, C12R001:91; C12P021-02,  
 C12R001:91

AB WO 9107495 A UPAB: 19940329  
 Mammalian IL-11 (I), free from other proteins, is new. (I) has a molecular weight of 20 kD (SDS-PAGE and calculations) and biological activity in T1165 assays, megakaryocyte colony forming assays with IL-3 and B cell plaque forming assays. Also new are a process for producing (I) recombinantly, DNA encoding (I), a cell transformed with this DNA, a plasmid vector containing the DNA and homogeneous mammalian (I) having biological activity in the T1165 assay without IL-6. Also present in the compsn. may be other cytokine e.g. IL-1 to IL-9, GM-CSF, G-CSF, M-CSF, the interferons, Meg-CSF, MIF, LIF, TNF and erythropoietin, haematopoietins e.g. IL-3 or IL-6, growth factors or antibodies. Dosage of (I) is 1-1000 micro-g or 50-5000 units, where a unit is the concentration leading to half maximal stimulation in the T1165 assay.

USE/ADVANTAGE - (I) is used to stimulate or treat disorders of the immune system e.g. deficiencies in haematopoietic progenitor stem cells (e.g. following bone marrow transplantation), and to treat cancer and other pathological states caused by disease, exposure to radiation or drugs and e.g. leukopenia, bacterial and viral infections, anaemia and B or T cells deficiencies. (I) is also used to prolong the effects of vaccines. Use of (I) does not create undesirable side effects. @69pp  
 Dwg.No.0/2)

0/2

FS CPI

FA AB

MC CPI: B04-B02B1; B04-B04A1; B04-C01G; B12-A01; B12-A06; B12-A07;  
 B12-G05; B12-G07; B12-H01; D05-C12; D05-H12

=> d all 156 tot

L56 ANSWER 1 OF 4 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 2000-594517 [56] WPIX

CR 2000-594515 [56]; 2000-594516 [56]; 2000-679550 [66]; 2001-006956 [01]

DNC C2000-177617

TI A Streptococcus pneumoniae vaccine for preventing pneumonia and meningitis comprises a polysaccharide antigen conjugated to protein D from Haemophilus influenzae.

DC B04 D16

IN CAPIAU, C; DESCHAMPS, M; DESMONS, P M; LAFERRIERE, C A J; POOLMAN, J; PRIEELS, J; POOLMAN, J P J

PA (SMIK) SMITHKLINE BEECHAM BIOLOGICALS

CYC 93

PI WO 2000056360 A2 20000928 (200056)\* EN 77 A61K039-385 <--  
 RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
 OA PT SD SE SL SZ TZ UG ZW  
 W: AE AG AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM DZ  
 EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK  
 LR LS LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI  
 SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2000034307 A 20001009 (200103) A61K039-385 <--

BR 2000009163 A 20011226 (200206) A61K039-385 <--

EP 1163000 A2 20011219 (200206) EN A61K039-385 <--  
 R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
 RO SE SI

CZ 2001003380 A3 20020313 (200223) A61K039-385 <--

KR 2002000549 A 20020105 (200244) A61K039-385 <--

HU 2002000367 B 20020528 (200249) A61K039-385 <--

CN 1351503 A 20020529 (200258) A61K039-385 <--

AU 750913 B 20020801 (200261) A61K039-385 <--

ZA 2001007637 A 20020828 (200264) 97 A61K000-00

JP 2002540075 W 20021126 (200307) 96 A61K039-09 <--

MX 2001009455 A1 20020301 (200362) A61K039-005 <--

NZ 513840 A 20040227 (200418) A61K039-385 <--

ADT WO 2000056360 A2 WO 2000-EP2468 20000317; AU 2000034307 A AU 2000-34307 20000317; BR 2000009163 A BR 2000-9163 20000317, WO 2000-EP2468 20000317; EP 1163000 A2 EP 2000-912626 20000317, WO 2000-EP2468 20000317; CZ 2001003380 A3 WO 2000-EP2468 20000317, CZ 2001-3380 20000317; KR 2002000549 A WO 2000-EP2468 20000317, KR 2001-711939 20010919; HU 2002000367 B WO 2000-EP2468 20000317, HU 2002-367 20000317; CN 1351503 A CN 2000-807528 20000317; AU 750913 B AU 2000-34307 20000317; ZA 2001007637 A ZA 2001-7637 20010917; JP 2002540075 W JP 2000-606264 20000317, WO 2000-EP2468 20000317; MX 2001009455 A1 WO 2000-EP2468 20000317, MX 2001-9455 20010919; NZ 513840 A NZ 2000-513840 20000317, WO 2000-EP2468 20000317

FDT AU 2000034307 A Based on WO 2000056360; BR 2000009163 A Based on WO 2000056360; EP 1163000 A2 Based on WO 2000056360; CZ 2001003380 A3 Based on WO 2000056360; KR 2002000549 A Based on WO 2000056360; HU 2002000367 B Based on WO 2000056360; AU 750913 B Previous Publ. AU 2000034307, Based on WO 2000056360; JP 2002540075 W Based on WO 2000056360; MX 2001009455 A1 Based on WO 2000056360; NZ 513840 A Based on WO 2000056360

PRAI GB 1999-16677 19990715; GB 1999-6437 19990319; GB 1999-9077 19990420; GB 1999-9466 19990423

IC ICM A61K000-00; A61K039-005; A61K039-09; A61K039-385

ICS A61K035-74; A61K039-02; A61K039-04; A61K039-085; A61K039-095; A61K039-102; A61K039-112; A61K039-116; A61K039-39; A61P011-00; A61P031-04; C07K014-285

ICA C12N015-09

AB WO 2000056360 A UPAB: 20040316

NOVELTY - A polysaccharide conjugate antigen (I) comprising a polysaccharide antigen derived from a pathogenic bacterium conjugated to protein D (or a fragment) from *Haemophilus influenzae*, is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:

- (1) an immunogenic composition comprising (I);
- (2) an immunogenic composition comprising *Neisseria meningitidis* protein D polysaccharide conjugate antigen;
- (3) an immunogenic composition comprising *Haemophilus influenzae* b protein D polysaccharide conjugate antigen;
- (4) an immunogenic composition comprising conjugated capsular polysaccharides of *Streptococcus pneumoniae*, *Haemophilus influenzae* b, meningococcus C and meningococcus Y, the carrier protein for at least one of the polysaccharides is protein D from *H. influenzae*;
- (5) a vaccine comprising (1)-(4); and
- (6) a method for producing an immunogenic composition to a pathogenic bacterium comprising:
  - (a) isolating a polysaccharide antigen from a pathogenic bacterium;
  - (b) activating the polysaccharide; and
  - (c) conjugating the polysaccharide to protein D.

ACTIVITY - Antibacterial. No biological data given

MECHANISM OF ACTION - Vaccine.

USE - The bacterial polysaccharide antigen vaccines are used to induce an immune response to *Streptococcus pneumoniae* and is used to prevent pneumonia, bacteremia, meningitis and acute otitis media.

ADVANTAGE - The conjugation of the antigen to a larger immunogenic protein increases the induced immune response, especially in children less than two years old.

Dwg.0/3

FS CPI

FA AB; DCN

MC CPI: B04-B04C1; B04-C02F; B04-C02V; B04-F10A; B04-F10B; B04-N03; B04-N05; B04-N06; B12-M07; B14-A01B2; B14-S11B; D05-H07

L56 ANSWER 2 OF 4 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 1994-025891 [03] WPIX

CR 1992-249778 [30]

DNC C1994-011927

TI New adhesion-oligosaccharide conjugate - useful as vaccine for *Haemophilus influenzae*, and new synthetic poly ribosyl ribitol phosphate oligosaccharide(s).

DC B04 D16

IN KRIVAN, H C; NORBERG, N T; SAMUELS, J E; SAMUEL, J E

PA (MICR-N) MICROCARB INC; (ANTE-N) ANTEX BIOLOGICS FORMERLY MICROCARB INC; (ANTE-N) ANTEX BIOLOGICS INC; (ANTE-N) ANTEXBIOLOGICS INC

CYC 20

PI WO 9400149 A1 19940106 (199403)\* EN 124 A61K039-00 <--

RW: AT BE CH DE DK ES FR GB GR IE IT LU MC NL PT SE

W: CA JP US

EP 647139 A1 19950412 (199519) EN A61K039-00 <--  
 R: AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL PT SE  
 JP 07509693 W 19951026 (199551) 33 C07K014-11  
 US 5679547 A 19971021 (199748) 35 C12P021-00  
 EP 647139 A4 19970903 (199815) A61K039-00 <--  
 US 5721115 A 19980224 (199815) 36 C12P021-06  
 JP 2805174 B2 19980930 (199844) 42 C07K014-11  
 US 5843463 A 19981201 (199904) A61K039-102 <--  
 CA 2138765 C 19990907 (200003) EN C12N015-31  
 ADT WO 9400149 A1 WO 1993-US6016 19930622; EP 647139 A1 EP 1993-916717  
 19930622, WO 1993-US6016 19930622; JP 07509693 W WO 1993-US6016 19930622,  
 JP 1994-502559 19930622; US 5679547 A CIP of WO 1986-DK14 19860206, CIP of  
 US 1990-631698 19901221, CIP of US 1991-810966 19911220, Div ex US  
 1992-903079 19920622, US 1995-485569 19950607; EP 647139 A4 EP 1993-916717  
 19930622; US 5721115 A CIP of US 1990-631698 19901221, CIP of US  
 1991-810966 19911220, Div ex US 1992-903079 19920622, US 1995-480993  
 19950607; JP 2805174 B2 WO 1993-US6016 19930622, JP 1994-502559 19930622;  
 US 5843463 A CIP of US 1990-631698 19901221, CIP of US 1991-810966  
 19911220, US 1992-903079 19920622; CA 2138765 C CA 1993-2138765 19930622,  
 WO 1993-US6016 19930622  
 FDT EP 647139 A1 Based on WO 9400149; JP 07509693 W Based on WO 9400149; JP  
 2805174 B2 Previous Publ. JP 07509693, Based on WO 9400149; CA 2138765 C  
 Based on WO 9400149  
 PRAI US 1992-903079 19920622; WO 1986-DK14 19860206;  
 US 1990-631698 19901221; US 1991-810966 19911220;  
 US 1995-485569 19950607; US 1995-480993 19950607  
 REP 2.Jnl.Ref; EP 276516; EP 320942; US 4455296; 3.Jnl.Ref; EP 338265; WO  
 9210936  
 IC ICM A61K039-00; A61K039-102; C07K014-11; C12N015-31;  
 C12P021-00; C12P021-06  
 ICS A61K037-02; A61K038-00; A61K039-02; C07H001-04; C07H013-00;  
 C07H015-02; C07H021-02; C07H021-04; C07K003-28; C07K014-00;  
 C07K014-285; C07K015-00; C07K019-00; C12N001-21; C12N015-06;  
 C12N015-09; C12P017-00; C12P021-02; C12P021-04; C12Q001-34;  
 G01N033-53  
 AB WO 9400149 A UPAB: 20000118  
 An immunogenic oligosaccharide-protein conjugate (I) comprising a  
 polyribosyl-ribitol phosphate (PRP) fragment coupled to an Haemophilus  
 influenzae adhesin protein (II) is new.  
 (II) is pref. an H. influenzae outer membrane protein with a mol.weight  
 of about 47000 daltons, and purified (II) is claimed per se.  
 USE - (I), as well as their protein components, may be used in  
 vaccines against both invasive and non-invasive strains of H. influenzae.  
 (I), (II) and oligomers are also useful as reagents for scientific  
 research on the properties of pathogenicity, virulence and infectivity of  
 H. influenzae, as well as host defence mechanisms. E.g. the novel DNA can  
 be used in an oligonucleotide probe to identify the DNA of other  
 microorganisms which might encode an adhesion for such organism. (I) can  
 be used to prepare a monoclonal antibody useful to further purity compsns.  
 containing (II) by affinity chromatography. (II) could also be applied to  
 standard immunoassays to screen for the presence of antibodies to H.  
 influenza in a sample. (IV) are intermediates in the synthesis of (III),  
 which may be used to prepare (Ia).  
 Dwg.0/9  
 FS CPI  
 FA AB; DCN  
 MC CPI: B04-C02; B04-C02X; B04-E02F; B04-E03F; B04-E08; B04-N03; B05-B01M;  
 B12-K04; B14-A01A; B14-S11B; D05-H07; D05-H12;  
 D05-H12E  
 L56 ANSWER 3 OF 4 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN  
 AN 1990-132245 [17] WPIX  
 DNC C1990-058101  
 TI Capsular polysaccharide adhesion antigen - from coagulase negative  
 bacteria used to prevent or treat infection caused by staphylococcal  
 strains.  
 DC A96 B04 D16 D22  
 IN PIER, G B  
 PA (BGHM) BRIGHAM & WOMENS HOSPITAL; (BGHM) BRIGHAM & WOMENS HOSPITAL INC;  
 (BRIG-N) BRIGHAM & WOMENS HO; (PIER-I) PIER G B  
 CYC 15  
 PI WO 9003398 A 19900405 (199017)\* 41  
 RW: AT BE CH DE FR GB IT LU NL SE  
 W: AU JP  
 AU 8943430 A 19900418 (199027)  
 EP 436648 A 19910717 (199129)

R: AT BE CH DE FR GB IT LI LU NL SE

US 5055455 A 19911008 (199143)

JP 04501718 W 19920326 (199219) 15

CA 1317288 C 19930504 (199323) C12P019-04

EP 436648 A4 19911113 (199520)

US 5980910 A 19911109 (19954) A61K039-09 <--

US 6399066 B1 20020604 (200242) A61K039-40 <--

US 2002136730 A1 20020926 (200265) A61K039-40 <--

US 6743431 B2 20040601 (200436) A61K039-085 <--

ADT EP 436648 A EP 1989-911517 19890928; US 5055455 A US 1988-250417 19880928;  
 JP 04501718 W JP 1989-510684 19890928; CA 1317288 C CA 1989-614255  
 19890928; EP 436648 A4 EP 1989-911517 ; US 5980910 A Div ex US  
 1988-250417 19880928, Cont of US 1991-727982 19910710, Cont of US  
 1993-33756 19930318, US 1994-336688 19941107; US 6399066 B1 Div ex US  
 1988-250417 19880928, Cont of US 1991-727982 19910710, Cont of US  
 1993-33756 19930318, Div ex US 1994-336688 19941107, US 1999-393832  
 19990910; US 2002136730 A1 Div ex US 1988-250417 19880928, Cont of US  
 1991-727982 19910710, Cont of US 1993-33756 19930318, Div ex US  
 1994-336688 19941107, Div ex US 1999-393832 19990910, US 2002-93582  
 20020308; US 6743431 B2 Div ex US 1988-250417 19880928, Cont of US  
 1991-727982 19910710, Cont of US 1993-33756 19930318, Div ex US  
 1994-336688 19941107, Div ex US 1999-393832 19990910, US 2002-93582  
 20020308

FDT US 5980910 A Div ex US 5055455; US 6399066 B1 Div ex US 5055455, Div ex US  
 5980910; US 2002136730 A1 Div ex US 5055455, Div ex US 5980910, Div ex US  
 6399066; US 6743431 B2 Div ex US 5055455, Div ex US 5980910, Div ex US  
 6399066

PRAI US 1988-250417 19880928; US 1991-727982 19910710;  
 US 1993-33756 19930318; US 1994-336688 19941107;  
 US 1999-393832 19990910; US 2002-93582 20020308

REP US 4789735; US 4830852; 2.Jnl.Ref; EP 302781; FR 2410043

IC A61K037-00; A61K039-02; A61K039-08; C07K015-04;  
 C07K015-14; C12P021-00  
 ICM A61K039-085; A61K039-09; A61K039-40;  
 C12P019-04

ICS A61K037-00; A61K039-02; A61K039-08; C07K015-04;  
 C07K015-14; C08B037-00; C12P021-00

AB WO 9003398 A UPAB: 19991122  
 The following are m (A) a capsular polysaccharide adhesion from  
 coagulase-negative bacteria (e.g. Staphylococcus epidermidis or hominus  
 strains) in pure form; (B) a vaccine against coagulase-negative  
 staphylococci comprising a vehicle containing the pure capsular polysaccharide  
 adhesionm antigen specific to the staphylococci; the vehicle may be e.g.  
 Freund's complete or incomplete adjuvant, saline, serum albumin or  
 saponin; (C) monovalent antibody (MAb) against capsular polysaccharide  
 adhesin of coagulase-negative bacteria.  
 USE/ADVANTAGE - The polysaccharide adhesin can produce  
 antibodies which prevent the adherence of adhesin-bearing  
 pathogenic bacteria to the recipients tissue cells or polymeric medical  
 prostheses or catheters and can therefore be used for preventing or  
 treating diseases and infections due to staphylococci. The adhesin  
 can also be used to screen polymeric materials for resistance to  
 attachment by bacteria. The MAbs can be administered to prevent or reduce  
 infections by coagulase-negative staphylococci. The adhesin  
 -specific antibodies can also be used in affinity chromatography and in  
 diagnosis and assays.  
 Dwg.0/5

FS CPI

FA AB

MC CPI: A09-C; A12-V01; A12-V02; B02-V02; B04-B04C5;  
 B04-C02; B12-A01; D05-H07; D05-H11

L56 ANSWER 4 OF 4 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 1983-42112K [18] WPIX

DNN N1983-076352 DNC C1983-041031

TI Mono clonal antibodies against bacterial adhesion(s) - useful for treating  
 diarrhoea in neonates, respiratory diseases and burns.

DC B04 D16 S03

IN SADOWKI, P L

PA (MOLE-N) MOLECULAR GENETICS INC

CYC 12

PI EP 77734 A 19830427 (198318)\* EN 37  
 R: BE DE FR GB IT LU NL  
 AU 8289454 A 19830428 (198324)  
 JP 58099423 A 19830613 (198329)  
 DK 8204621 A 19830620 (198331)

US 4443549 A 19840417 (198418)  
 CA 1187822 A 19850528 (198526)  
 US 4652448 A 19870324 (198714)  
 ADT US 4443549 A US 1982-428622 19821007; US 4652448 A US 1983-558518 19831206  
 PRAI US 1981-312993 19811019; US 1982-428622 19821007;  
 US 1983-558518 19831206  
 REP 4.Jnl.Ref; No-SR.Pub  
 IC A61K039-40; C12N005-00; C12N015-00; C12P001-00; C12R001-91;  
 G01N033-54  
 AB EP 77734 A UPAB: 19930925  
 Production of anti-adhesion antibodies comprises fusion of a cell producing the antibodies with a myeloma to provide a fused cell hybrid, followed by propagation of the hybrid and collection of the antibodies.  
 Production of antipilus antibodies comprises injection of a BALB/C mouse with a bacterial pilus to induce formation of antibacterial pilus antibody-producing cells of the mouse. Then a fused cell hybrid of the cells is produced with P3/NSI/1-Ag4-1 myeloma cells, and the hybrid is cultured in vitro in selective HAT medium, isolated and propagated and the resulting antibodies are harvested.  
 Continuous cell line producing anti-adhesion antibodies and comprising a fused cell hybrid of an anti-adhesion antibody-producing cell and a myeloma cell is new, and cell line 2BD4E4 (ATCC HB8178) is new.  
 Monoclonal antibodies against Escherichia coli adhesions are useful for admin. to animals and humans, especially for the prophylaxis and treatment of enterotoxigenic diarrhoeal diseases in neonatal calves, lambs and piglets. The antibodies are obtainable in large amts. and are useful as highly sensitive and specific probes in medicinal and veterinary diagnosis, etc. The anti-adhesion antibodies may also be useful against respiratory diseases and burn infections and against other bacterial diseases. The antibodies are also useful in affinity chromatography systems and in the assay of adhesions.  
 FS CPI EPI  
 FA AB  
 MC CPI: B04-B04A; B04-B04C; B12-A01; B12-J04; B12-K04;  
 D05-H  
 EPI: S03-E14H4

=> d all 161 tot

L61 ANSWER 1 OF 5 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN  
 AN 2000-038242 [03] WPIX  
 CR 1993-093726 [11]; 2000-012250 [01]  
 DNC C2000-009691  
 TI Purified *Moraxella catarrhalis* outer membrane proteins useful for vaccinating against chronic otitis media, acute maxillary sinusitis and other bronchopulmonary and lower respiratory tract infections.  
 DC B04 D16  
 IN HANSEN, E J; HELMINEN, M E; MACIVER, I  
 PA (TEXA) UNIV TEXAS  
 CYC 1  
 PI US 5993826 A 19991130 (200003)\* 50 A61K039-102 <--  
 ADT US 5993826 A CIP of US 1991-745591 19910815, CIP of WO 1992-US6869 19920814, US 1993-25363 19930302  
 FDT US 5993826 A CIP of US 5552146  
 PRAI US 1993-25363 19930302; US 1991-745591 19910815;  
 WO 1992-US6869 19920814  
 IC ICM A61K039-102  
 ICS A61K039-02; C07K014-285; C07K016-102  
 AB US 5993826 A UPAB: 20000925  
 NOVELTY - A purified *Moraxella catarrhalis* (also called *Branhamella catarrhalis* and *Neisseria catarrhalis*) 80 kiloDalton (kD) CopB outer membrane protein (I), is new.  
 DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for:  
 (i) an antigen composition (II) prepared by:  
 (1) introducing a recombinant expression vector including a DNA segment encoding (I) into a recombinant host cell;  
 (2) culturing the host cell under suitable conditions for the expression of (I); and  
 (3) collecting the expressed antigen; and  
 (ii) a method (III) for inducing an antibody response to M. *catarrhalis* 80 kD CopB antigens in an animal, comprising administering (I).  
 ACTIVITY - Auditory; Respiratory active.  
 MECHANISM OF ACTION - Vaccine, administration of (I) stimulates an immune response against M. *catarrhalis* antigens in a patient.

Search done by Noble Jarrell

Groups of mice were immunized with the 8B6 monoclonal antibody, specific for the 80 kD outer membrane protein of *M. catarrhalis*. Control mice were immunized with an irrelevant antibody, 2H11 which is specific for *Haemophilus ducreyi*. Doses of 150 micrograms were used 18 hours prior to bacterial challenge. 5 Microliter doses of bacterial suspension, containing *M. catarrhalis* strain 035E, were inoculated into the lungs of the mice. 6 Hours after inoculation, the mice were sacrificed and the number of bacteria remaining in the lungs was determined. It was found that where the 2H11 antibody was used, 97% of the initial bacterial population remained. However, just 38% remained when the 8B6 antibody was used.

USE - (I) may be used to vaccinate against *M. catarrhalis*, a pathogen implicating in causing chronic otitis media, acute maxillary sinusitis and other bronchopulmonary and lower respiratory tract infections.

Dwg.0/13

FS CPI  
FA AB; DCN  
MC CPI: B04-B04C1; B04-C01G; B04-E03F; B04-F0100E; B04-F10A5;  
B04-G09; B04-N03A; B11-C07A; B11-C08E1; B11-C09; B12-M07; B12-M08;  
B14-A01A5; B14-K01; B14-N02; B14-N04; B14-N05;  
B14-S11B; D05-C12; D05-H04; D05-H07; D05-H08; D05-H11;  
D05-H12A; D05-H14; D05-H17A5; D05-H18

L61 ANSWER 2 OF 5 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 1999-560502 [47] WPIX

CR 1990-304862 [40]; 1997-131755 [12]; 1998-413151 [35]; 2002-664558 [71]

DNC C1999-163303

TI Isolated *Haemophilus influenzae* protein e useful for producing vaccines against meningitis, pneumonia, bacteremia, postpartum sepsis, acute febrile tracheo-bronchitis and otitis media.

DC B04 D16

IN GREEN, B A; ZLOTNICK, G W

PA (PRAX-N) PRAXIS BIOLOGICS INC

CYC 1

PI US 5955580 A 19990921 (199947)\* 23 C07K001-00

ADT US 5955580 A CIP of US 1989-320971 19890309, Div ex US 1990-491466

19900309, US 1995-449406 19950523

FDT US 5955580 A Div ex US 5601831

PRAI US 1990-491466 19900309; US 1989-320971 19890309;

US 1995-449406 19950523

IC ICM C07K001-00

ICS A61K039-102; C07K014-285

AB US 5955580 A UPAB: 20021108

NOVELTY - Isolated *Haemophilus influenzae* protein e, purified of endotoxins, is new.

DETAILED DESCRIPTION - An isolated protein e from *Haemophilus influenzae* (free from endotoxic contamination). The protein (when administered to a mammal) is capable of raising antibodies in the mammal which are protective in the infant rat passive immunization model.

USE - The proteins can be used for vaccination against nontypable and typable *H. influenzae*. They can be used to immunize against diseases including meningitis, pneumonia, bacteremia, postpartum sepsis, acute febrile tracheo-bronchitis or otitis media. The bactericidal antibodies induced by protein e epitopes can be used to passively immunize an individual against *H. influenzae*. The antibody products can also be used for the detection of e proteins (e.g. via enzyme linked immunoabsorbant assay (ELISA)) and for the diagnosis of *H. influenzae* disease.

Dwg.0/8

FS CPI

FA AB; DCN

MC CPI: B04-B04C1; B04-B04M; B04-C01; B04-E03F; B04-F0100E;  
B04-F10A; B04-G07; B04-N0300E; B11-A; B11-C07A; B11-C08D; B11-C08E1;  
B11-C09; B12-K04A4; B12-K04E; B12-M05; B14-A01A; B14-C03;  
B14-G01; B14-G03; B14-K01; B14-L06; B14-N02; B14-N16;  
B14-P01; B14-S11B; D05-A01A4; D05-A01B; D05-C12; D05-H04;  
D05-H07; D05-H09; D05-H11; D05-H12A; D05-H17A5; D05-H18

L61 ANSWER 3 OF 5 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 1999-517930 [43] WPIX

CR 1996-010692 [01]

DNC C1999-151167

TI Antigenic peptide, oligopeptide and protein useful as vaccine for *Moraxella catarrhalis*.

DC B04 D16

IN MURPHY, T F

PA (UYNV) UNIV NEW YORK STATE RES FOUND

CYC 1  
 PI US 5948412 A 19990907 (199943)\* 20 A61K039-02 <--  
 ADT US 5948412 A CIP of US 1994-245758 19940517, US 1997-810655 19970303  
 FDT US 5948412 A CIP of US 5607846  
 PRAI US 1997-810655 19970303; US 1994-245758 19940517  
 IC ICM A61K039-02  
 ICS C07K014-00  
 AB US 5948412 A UPAB: 19991124  
 NOVELTY - Pure antigenic peptide, oligopeptide or protein (I) with one or more epitopes of E, an outer membrane protein of *Moraxella catarrhalis* is new.

DETAILED DESCRIPTION - E has an apparent molecular weight of 35000-50000 daltons by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and amino acid residues 26-429 of a defined sequence of 459 amino acids given in the specification. E is a heat-modifiable protein.

An INDEPENDENT CLAIM is also included for an antigenic formulation comprising a pure peptide, oligopeptide or protein with one or more epitopes of E.

ACTIVITY - Antibacterial.

MECHANISM OF ACTION - Vaccine.

USE - (I) can be used as immunogens in prophylactic and/or therapeutic vaccine formulations for active immunization and for generating protein-specific and peptide-specific antisera useful for passive immunization.

Antigenic formulations comprising (I) can be used to prevent otitis media, sinusitis, conjunctivitis and lower respiratory tract infections caused by *Moraxella catarrhalis*. (I) can be used as antigens for diagnostic immunoassays.

Dwg.0/3

FS

CPI

FA

MC

AB; DCN  
 CPI: B04-B04C2; B04-B04D5; B04-C01G; B04-E03F; B04-E08;  
 B04-F0100E; B04-N03A; B04-N04B0E; B11-C07A4; B12-K04A4;  
 B14-A01A; B14-K01; B14-N02; B14-N03;  
 B14-S11B; D05-C12; D05-H07; D05-H09; D05-H11; D05-H12A;  
 D05-H12D5; D05-H12E; D05-H14; D05-H17A5; D05-H18; D05-H19

L61 ANSWER 4 OF 5 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 1986-208380 [32] WPIX

DNC C1986-089596

TI Oral vaccine for prophylaxis of periodontitis - comprises vaccine as antigen containing whole cell, pilus or extract of periodontitis causing bacterium.

DC B04 D16 D21

PA (LIOY) LION CORP

CYC 1

PI

JP 61140527 A 19860627 (198632)\* 4  
 JP 06062431 B2 19940817 (199431) 5 A61K039-02 <--

ADT JP 61140527 A JP 1984-263874 19841214; JP 06062431 B2 JP 1984-263874 19841214

FDT JP 06062431 B2 Based on JP 61140527

PRAI JP 1984-263874 19841214

IC

A61K039-02

ICM A61K039-02

ICS A61K039-114

AB

JP 61140527 A UPAB: 19930922

Oral vaccine for prophylaxis of periodontitis, where the vaccine is made as antigen, which is whole cell, pilus or extract of periodontitis causative bacterium. Bacterium is eq. *Bacterioides gingivalis* or *Actinomyces viscosus*.

USE/ADVANTAGE - By oral administration of this vaccine, local immunity mechanism is stimulated, by antibody eq. IgA, IgM, injection by periodontitis causative bacterium is protected specifically. Especially inoculation of vaccine at juvenile age, gives long period of immunological competence, it is effective for prophylaxis of adult periodontitis. It is more safety than injective administration.

Gingivalis 381 strain is cultured on hemin and menadione added Todd-Hewlett broth for 2 days. Cell is collected by centrifugation (8,000 r.p.m 15 min), washed by phosphate buffer (5 mM, pH 7.4), treated by 0.5% formalin over night, and inactivated vaccine of cell antigen is obtd. The antigen is stored in refrigerator (at -80 deg.C), and used by thawing. For administration, in the case of antigen, cell containing solution adjusted at 10 power 4 - 10 power 10 /ml is administered (p.o) at 0.1-10 ml/day for 3-15 days. In the case of pilus or extract antigen, these containing solution adjusted at 0.01-10 mg/ml is administered (p.o) at 0.1-10 ml for 3-15 days

continuously.

0/0

FS CPI

FA AB

MC CPI: B02-V02; B04-B02B1; B04-B04C1; B12-A01;  
B12-L03; B12-L04; D05-H07

L61 ANSWER 5 OF 5 WPIX COPYRIGHT 2004 THE THOMSON CORP on STN

AN 1985-236083 [38] WPIX

DNC C1985-102472

TI Vaccine against infectious bovine keratoconjunctivitis - comprises  
Neisseria or Branhamella gram negative cocci.

DC B04 C03 D16

PA (GWIN-I) GWIN R M

CYC 1

PI US 4539201 A 19850903 (198538)\* 10

ADT US 4539201 A US 1983-546600 19831028

PRAI US 1983-546600 19831028

IC A61K039-09; C12R001-36

AB US 4539201 A UPAB: 19930925

Medicament inducing immunity to infections bovine keratoconjunctivitis  
(IBK) in cattle comprises an effective amount of gram -ve cocci from  
Neisseria and Branhamella sp., pref. those which are nonpathogenic in  
cattle, and not N gonorrhoeae or N. meningitidis. Admin. is pref.  
topically to the eye.

ADVANTAGE - The microorganisms used are effective immune stimulators,  
producing antibodies effective to produce immunity against Moraxella  
bovis, vaccines containing which do not produce practical protection against  
IBK (pinkeye).

0/7

FS CPI

FA AB

MC CPI: B02-V; B12-A01; B12-L04; B12-L09; C02-V;  
C12-A01; C12-L04; C12-L09; D05-H07

=> b home

FILE 'HOME' ENTERED AT 13:11:19 ON 16 DEC 2004

=>